

US008063182B1

(12) United States Patent

Brockhaus et al.

(54) HUMAN TNF RECEPTOR FUSION PROTEIN

- (75) Inventors: Manfred Brockhaus, Bettingen (CH); Reiner Gentz, Rheinfelden (DE); Dembic Zlatko, Basel (CH); Werner Lesslauer, Basel (CH); Hansruedi Lotscher, Mohlin (CH); Ernst-Jurgen Schlaeger, Efringen-Kirchen (DE)
- (73) Assignee: Hoffman-LaRoche Inc., Nutley, NJ (US)
- (*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.
- (21) Appl. No.: 08/444,790
- (22) Filed: May 19, 1995

Related U.S. Application Data

(60) Division of application No. 08/095,640, filed on Jul. 21, 1993, now Pat. No. 5,610,279, which is a continuation of application No. 07/580,013, filed on Sep. 10, 1990, now abandoned.

(30) Foreign Application Priority Data

Sep. 12, 1989	(CH)	
Mar. 8, 1990	(CH)	
Apr. 20, 1990	(CH)	1347/90
Aug. 31, 1990	(EP)	

(51) Int. Cl.

C07K 14/715	(2006.01)
A61P 29/00	(2006.01)
A61K 38/17	(2006.01)
C07K 19/00	(2006.01)
C07H 21/04	(2006.01)

- (52) **U.S. Cl.** **530/350**; 514/12.2; 530/387.3; 536/23.5; 930/144
- (58) Field of Classification Search 530/324,

530/350; 424/192.1; 536/23.5 See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,593,002 A	6/1986	Dulbecco	
4,675,285 A	6/1987	Clark et al.	
4,729,326 A	3/1988	Richter	
4,769,326 A	9/1988	Rutler	
4,770,995 A	9/1988	Rubin et al.	
4,816,567 A	3/1989	Cabilly et al.	
4,894,439 A	1/1990	Dorin et al.	
4,912,044 A	3/1990	Jacob et al.	
4,935,233 A	6/1990	Bell et al.	
4,948,875 A	8/1990	Tanaka et al.	
4,963,354 A	10/1990	Shepard et al.	
4,965,271 A	10/1990	Mandell et al.	
5,055,447 A	10/1991	Palladino et al.	
5,073,627 A	12/1991	Curtis et al.	
5,075,222 A	12/1991	Hannum et al.	
5,098,702 A	3/1992	Zimmerman et al.	
5,098,833 A	3/1992	Lasky et al.	
5,116,964 A	* 5/1992	Capon et al.	536/23.5

(10) Patent No.: US 8,063,182 B1

(45) **Date of Patent:** Nov. 22, 2011

Hanel et al.
5 11 11 1
Dembinski et al.
Sledziewski et al.
Wallach et al.
Gero
Capon et al.
Huston et al.
Park et al.
Klimpel et al.
Capon et al.
LeMaire et al 530/350
Sims et al.
Dayer et al.
Wallach et al 530/388.22
Smith et al 435/365
Capon et al.
Beutler et al.
Capon et al.
Wallach et al.
Wallach et al.
Capon et al.
Mosley et al.
Jacobs et al.
Brockhaus et al.
Feldmann et al.
· 1)

(Continued)

FOREIGN PATENT DOCUMENTS

58976 1/1991

(Continued)

OTHER PUBLICATIONS

Miles JS, et al. Principles of DNA cloning. British Medical Journal. 1989 299:1019-1022.*

Olsson, I, et al., Isolation and charaterization of a tumor necrosis factor binding protein from urine. Eur J Haematol. 1989;42:270-275.*

Dembic et al, 1990. Cytokine. 2(4): 231-7.*

AU

Chan et al. 2000, Science, 288: 2351-2354.*

Smith et al, 1989. Journal of Biological Chemistry. 14646-15652.* Heller et al., JBC 265(10): 5708-5717 (1990).

(Continued)

Primary Examiner — Gary Nickol

Assistant Examiner - Zachary Howard

(74) Attorney, Agent, or Firm — Marshall, Gerstein & Borun LLP

(57) **ABSTRACT**

The present invention is concerned with non-soluble proteins and soluble or insoluble fragments thereof, which bind TNF, in homogeneous form, as well as their physiologically compatible salts, especially those proteins having a molecular weight of about 55 or 75 kD (non-reducing SDS-PAGE conditions), a process for the isolation of such proteins, antibodies against such proteins, DNA sequences which code for non-soluble proteins and soluble or non-soluble fragments thereof, which bind TNF, as well as those which code for proteins comprising partly of a soluble fragment, which binds TNF, and partly of all domains except the first of the constant region of the heavy chain of human immunoglobulins and the recombinant proteins coded thereby as well as a process for their manufacture using transformed pro- and eukaryotic host cells.

36 Claims, 6 Drawing Sheets

U.S. PATENT DOCUMENTS

5,639,597	Α	6/1997	Lauffer et al.
5,695,953	Α	12/1997	Wallach et al.
5,705,364	Α	1/1998	Etcheverry et al.
5,712,155	Α	1/1998	Smith et al 735/320.1
5,721,121	Α	2/1998	Etcheverry et al.
5,808,029	Α	9/1998	Brockhaus et al.
5,811,261	A *	9/1998	Wallach et al 435/69.1
5,863,786	Α	1/1999	Feldmann et al.
5,447,851	Α	7/1999	Beutler et al.
5,945,397	Α	8/1999	Smith et al.
5,981,701	A *	11/1999	Wallach et al 530/350
RE36,755	Е	6/2000	Smith et al.
6,143,866	Α	11/2000	Brewer et al.
6,165,476	Α	12/2000	Strom et al.
6,201,105	B1	3/2001	Smith et al.
6,221,675	B1	4/2001	Hauptmann et al.
6,271,346	B1	8/2001	Hauptmann et al.
6,294,352	B1	9/2001	Hauptmann et al.
6,417,158	B1	7/2002	Hauptmann et al.
6,541,610	B1	4/2003	Smith
6,541,620	B1	4/2003	Brewer et al.
6,572,852	B2	6/2003	Smith et al.
6,858,409	B1	2/2005	Thompson et al.
7,253,264	B1	8/2007	Lauffer et al.
2003/0064480	A1	4/2003	Lauffler et al.

FOREIGN PATENT DOCUMENTS

EP	120694	10/1984
EP	227110	7/1987
EP	230574	8/1987
EP	269455	6/1988
EP	315 062	10/1988
EP	308 378	3/1989
EP	0 314 317	5/1989
EP	325 224	7/1989
EP	325262	7/1989
EP	0334165	9/1989
EP	393 438	4/1990
EP	0 394 827	10/1990
EP	398 327	11/1990
EP	412 486	2/1991
EP	414178	2/1991
EP	418 014	3/1991
EP	417563	3/1991
EP	422 339	4/1991
EP	433 900	6/1991
EP	460846	12/1991
EP	464 533	1/1992
EP	471701	2/1992
EP	526452	2/1993
EP	526905	2/1993
EP	0 567 566 B1	11/1993
EP	568925	11/1993
EP	606869	7/1994
GB	2218101	11/1989
GB	2 246 569	2/1992
ЛЪ	61-293924	12/1986
JP	61-293924 A2	12/1986
JP	02-154695	6/1990
WO	WO 89/02922 *	^k 4/1989
WO	WO 89 09622	10/1989
WO	9013575	11/1990
WO	WO 91/02078	2/1991
WO	91/03553	3/1991
WO	WO 91/17184	11/1991
WO	WO 91/08298	12/1991
WO	WO 92/08495	5/1992
WO	WO 92/13095	8/1992
WO	WO 93/07863	4/1993
WO	WO 93/19777	10/1993
WO	WO 94/06476	3/1994

OTHER PUBLICATIONS

Abstract by Heller et al., Napa Valley Conference (1989). Olsson et al., Eur. J. Haematol., 42: 270-275 (1989). Nophar et al., EMBO J., 9:3269-3278 (1990). Evans, T.J. et al., J. Exp. Med. 180, pp. 2173-2179 (1994). Traunecker, A. et al., Nature, 339, pp. 68-70 (1989).

Dembie, Z., et al., Cytokine, 2, pp. 231-237 (1990).

Byrn, et al., Nature, 344, pp. 667-670 (1990).

Peppel, K., et al., J. Exp. Med., 174, pp. 1483-1489 (1991).

Zettlmeissl, G., et al., DNA & Cell Biology, 9, pp. 347-353 (1990). Loetscher et al, *J. of Bio. Chem.*, 266(27), pp. 18324-18329, (1991). Nature Biotechnology, 15 (1997) p. 13.

Fisher, et al., N. Engl. J. of Med. 334, pp. 1697-1702 (1996).

Hsu, et al., J. Biol. Chem. 268, pp. 16430-16436 (1993).

Mohler et al., Soluble Tumor NecrosisFfactor (TNF) Receptors are Effective Therapeutic Agents in Lethal Endotoxemia and Function Simultaneously as Both TNF Carriers and TNF Antagonists. J. Immunol., 151:1548-1561 (1993).

Sell, Immunology, Immunopathology and Immunity, 4th Edition, Elsevier Science Publishing Co., New York, 1987, at pp. 85-91.

Bringman, Monoclonal antibodies to human tumor necrosis factors alpha and beta: application for affinity purification, immunoassays, and as structural probes. Hybridoma, 6(5):489-507 (1987).

Paul et al., Fundamental Immunology, 2nd Edition, Paul, ed., Raven Press, New York, 1989, at pp. 679-701 and pp. 735-764.

Smith et al., A Receptor for Tumor Necrosis Factor Defines an Unusual Family of Cellular and Viral Proteins, *Science*, 248:1019-1023, (1990).

Wingfield et al., Tumour Necrosis Factor is a Compact Trimer, FEBS Lett. 211: 179-84 (1987).

Smith and Baglioni, The ActiveFform of Tumor Necrosis Factor is a Trimer J. Biological Chemistry, 262:6951-4 (1987).

Larsson and Mosbach, Affintiy Precipitation of Enzymes FEBS Lett. 98(2):333-338 (1979).

Irwin and Tipton, Chapter 22, "Affinity Precipitation Methods" in Methods in Mol. Bio., 59:217 (1996).

Kohno et al., Presentation 1495, poster 271 presented at American College of Rheumatology Annual Meeting, Nov. 13-17, 2005, San Diego, CA.

Klein, Immunology, 1st Edition, Klein ed., Blackwell Scientific Publications, Cambridge, MA 1990 at pp. 446-447.

Byrn et al., Biological properties of a CD4 Immunoadhesin, Nature, 344:667-670 (Apr. 1990).

Traunecker et al., Highly Efficient Neutralization of HIV with Recombinant CD-4-Immunoglobulin Molecules, Nature 339:68-70, 1989.

Capon, Designing CD4 Immunoadhesions for AIDS Therapy, Nature, 337:525-531 (1989).

Barone et al., Comparative Analysis of the Ability of Entanercept and Inflixamab to Lyse TNF-Expressing Cells in a Complement-Dependent fashion, Arthritis Rheum., v42(9) supplement, Sep. 1999.

Khare et al, Poster 715 presented at the Annual Meeting of the Society for Investigative Dermatology (SID), May 3-5, 2006, Philadelphia, PA.

Nesbitt, et al., "Mechanism of Action of Certolizumab Pegol (Cdp870): In Vitro Comparison With Other Anti-Tumor Necrosis Factor α Agents", *Inflamm Bowel Dis*, 13: 1323-1332 (Nov. 2007).

Aruffo et al., "CD44 Is the Principal Cell Surface Receptor for Hyaluronate," Cell 61:1303-1313 (1990).

By enabled at al., "Inhibitory Effects of TNF α Antibodies on Synovial Cell Interleukin-1 Production in Rheumatoid Arthritis," The Lancet 2(8657):244-247 (1989).

Monnat, "Molecular Analysis of Spontaneous Hypoxanthine Phosphoribosyltransferase Mutations in Thioguanine-resistant HL-60 Human Leukemia Cells," Cancer Res. 49:81-87 (1989).

Shalaby et al., "The Involvement of Human Tumor Necrosis Factors- α and - β in the Mixed Lymphocyte Reaction," J. Immunol. 141:449-503 (1988).

Watson et al., "A Homing Receptor—IgG Chimera as a Probe for Adhesive Ligands of Lymph Node High Endothelial Venules," J. Cell Biol. 110:2221-2229 (1990).

Gen Seq DBase Printout.

Stauber et al., "Human tumor necrosis factor- α receptor—purification by immunoaffinity chromatography and initial characterization", J. Bio. Chem. 263:19098-19104 (1988).

Seckinger et al., "Purification and biologic characterization of a sp cific tum r necrosis fact r α inhibitor", J. Bio. Chem. 264:11966-11973 (1989).

Engelmann et al., "A tumor necrosis factor-binding protein purified to homogeneity from human urine protects cells from tumor necrosis factor toxicity", J. Bio. Chem. 264:11974-11980 (1989).

Hohmann et al., "Two different cell types have different major r cept rs f r human tumor necr sis factor ($TNF\alpha$)", J. Bio. Chem. 264:14927-14934 (1989).

Smith t al., "A receptor for tum r necrosis factor defines an unusual family of cellular and viral proteins", Scienc 248:1019-1023 (1990). Heller et al., "Complementary DNA cloning of a receptor for tumor necrosis factor and demonstration of a shed form of the receptor", Proc. Natl. Acad. Sci. U.S.A. 87:6151-6155 (1990).

Novick et al., "Soluble cytokine receptors are present in normal human urine", J. Exp. Med. 170:1409-1414 (1989).

Engelmann et al., "Two tum r necr sis factor-binding proteins purified from human urine", J. Bio. Chem. 265:1531-1536 (1990).

Schall et al., "Molecular cloning and expression of a receptor for human tumor necrosis factor", Cell 61:361-370 (1990).

Seckinger et al., "Tumor necrosis factor inhibitor: purification, NH₃terminal amino acid sequence and evidence for anti-inflammatory and immunomodulatory activities", Eur. J. Immunol. 20:1167-1174 (1990).

Hohmann et al., "Expression of the types A and B tumor necrosis factor (TNT) recept rs is independently r gulated, and both receptors mediate activation of the transcription factor NF-kB", J. Bi . Chem. 265:22409-22417 (1990).

Espevik et al., "Characterization of binding and biological effects of monoclonal antibodies against a human tumor necrosis factor receptor", J. Exp. Med. 171:415-426 (1990).

Porteu and Nathan, "Shedding of tumor necrosis factor receptors by activated human neutrophils", J. Exp. Med. 172:599-607 (1990).

Engelmann et al., "Antibodies to a s luble f rm f a tumor necrosis factor (TNF) receptor have TNF-like activity", J. Bio. Chem. 265:14497-14504 (1990).

Seckinger et al., "Characterization of a tumor necrosis factor α (TNF- α) inhibitor: evidence of immunological cross-activity with the TNF receptor", Proc. Natl. Sci. USA 87:5188-5192 (1990).

Gray et al., "Cloning of human tumor necrosis factor (TNF) receptor cDNA and expression of recombinant soluble TNF-binding protein", Proc. Natl. Sci. USA 87:7380-7384 (1990).

Loetscher et al., "Molecular cloning and expression of the human 55 kd tum r necrosis fact r rec ptor", Cell 61:351-359 (1990).

Peppel et al., "Chimaeric TNF-receptor—IgG molecule acts as soluble inhibitor of TNF mediated cytotoxicity", Journal of Cellular Biochem., Abstract, 20th Annual Meetings, Keystone Symposia on Molecular and Cellular Biology, p. 118, Supplement 15F (1991).

Olsson et al., "Isolation and characterization of a tumor necrosis factor binding pr tein fr m urin", Eur. J. Ha matol. 42:270-275 (1989). Capon et al., "Desiging CD4 immunoadhesins for AIDS therapy", Nature 337:525-530 (Feb. 9, 1989).

Abstract 92-009794/02 (1992) for EP 464 533.

U.S. Appl. No. 08/444,791-Office Action mailed Jun. 8, 2010.

U.S. Appl. No. 08/444,791-Office Action mailed Oct. 15, 2010.

Declaration of Taruna Arora, PH.D. Under 37 C.F.R. 1.132 with Exhibits A-D, Dec. 16, 2010.

Heller et al., "Amplified Expression of Tumor Necrosis Factor Receptor in Cells Transfected with Epstein-Barr Virus Shuttle Vector cDNA Libraries," J. Biol. Chem. 265(10): 5708-5717 (1990).

Arora et al., "Differences in Binding and Effector Functions Between Classes of TNF Antagonists," Cytokine 45: 124-131 (2009).

Furst et al., "Tumor Necrosis Factor Antagonists: Different Kinetics and/or Mechanisms of Action May Explain Differences in the Risk for Developing Granulomatous Infection," Semin. Arthritis Rheum. 36: 159-167 (2006).

Mitoma et al., "Mechanisms for Cytotoxic Effects of Anti-Tumor Necrosis Factor Agents on Transmembrane Tumor Necrosis Factor α -Expressing Cells," Arthr. & Rheum. 58(5): 1248-1257 (2008).

Strangfeld et al., "Risk of Herpes Zoster in Patients with Rheumatoid Arthritis Treated with Anti-TNF-α Agents," JAMA 301(7): 737-744 (2009).

Wallis et al., "Granulomatous Infectious Diseases Associated with Tumor Necrosis Factor Antagonists," Clin. Inf. Dis. 38: 1261-1265 (2004). Wallis et al., "Reactivation of Latent Granulomatous Infections by Infliximab," Clin. Inf. Dis. 41(Suppl 2): S1-S5 (2005).

Winzor et al., "Evaluation of Equilibrium Constants from Precipitin Curves: Interaction of α -Crystallin with an Elicited Monoclonal Antibody," Arch. Biochem. Biophys. 268(1): 221-226 (1989).

Avis, Kenneth, Parental Preparations, Remington Pharmaceutical Sciences, Chapter 85, pp. 1518-1541 (1985).

Center for Drug Evaluation and Research, Guidelines on Sterile Drug Products Produced by Aseptic Processing, Published by the FDA Jun. 1987.

Feldmann et al., "Cytokine production in the rheumatoid joint: implications for treatment," Ann. Rheum. Dis. 49: 480-486 (1990).

Hoogenboom et al., "Construction and Expression of Antibody-Tumor Necrosis Factor Fusion Proteins," Molecular Immunol. 28(9): 1027-1037 (1991).

Smitii, "cDNA Expression: Cloning of the Receptor for Human Tumor Necrosis Factor," Presentation Programme, 29th Midwinter Conference of Immunologists (Jan. 27-30, 1990).

U.S. Appl. No. 08/484,783, filed Jun. 7, 1995, Kohno et al.

U.S. Appl. No. 07/555,274, filed Jul. 19, 1990, Kohno et al.

Final Office action dated Jun. 24, 2011, in U.S. Appl. No. 08/444,791 (Brockhaus, et al.).

U.S. Appl. No. 08/478,995, Lauffler, Leander et al.

Abraham et al., p55 Tumor Necrosis Factor Receptor Fusion Protein in the Treatment of Patients With Severe Sepsis and Septic Shock: JAMA, 19:1531-1538 (1997).

Abraham et al., Lenercept (p55 Tumor Necrosis Factor Receptor Fusion Protein) In Severe Sepsis and Early Septic Shock: A Randomized, Double-Blind, Placebo-Controlled, Multicenter Phase III Trial With 1,342 Patients, Crit Care Med, 29:503-510 (2001).

Aggarwal et al., Characterization of Receptors for Human Tumour Necrosis Factor and Their Regulation by γ -Interferon, Nature, 318:665-667 (1985).

Aggarwal et al., Induction of Receptors for Tumor Necrosis Factor- α by Interferons Is Not a Major Mechanism for Their Synergistic Cytotoxic Response, J. Biol. Chem., 262:10000-10007 (1987).

Aggarwal et al., Human tumour necrosis factors: structure and receptor interactions, in Tumor necrosis factor and related cytotoxins, pp. 39-51, (Ciba Foundation symposium 131), Wiley, Chichester (1987). Arenzana-Seisdedos et al., Immunoregulatory Mediators in the Pathogenesis of Rheumotoid Arthritis, Scand. J. Rheumatol., Supplement 66:13-17 (1987).

Aruffo et al., Molecular Cloning of a CD28 cDNA by a High-Efficiency COS Cell Expression System, Proc. Natl. Acad. Sci. USA, 84:8573-8577 (1987).

Ashkenazi et al., Protection Against Endotoxic Shock by a Tumor Necrosis Factor Receptor Immunoadhesin, Proc. Natl. Acad. Sci., U.S.A. 88:10535-10539 (1991).

Ayala, Modem Genetics, Benjamin/Cummings, Publ. Co., Menlo Park CA, p. 45, (1980).

Baglioni et al., Binding of Human Tumor Necrosis Factor to High Affinity Receptors on HeLa and Lymphoblastoid Cells Sensitive to Growth Inhibition, J. Biol. Chem., 260:13395-13397 (1985).

Benjamini et al., Antibody Structure, in Immunology: A Short Course, 3rd ed., Wiley-Liss New York, 61-69 (1996).

Branellec et al., TNF: Antitumoral Agent at the Border Lines of Immunity and Inflammation, Path. Biol., 39:230-239 (1991).

Brockhaus et al., Identification of Two Types of Tumor Necrosis Factor Receptors on Human Cell Lines by Monoclonal Antibodies, Proc. Natl. Acad. Sci. USA, 87:3127-3131 (1990).

Carter et al., Purification, Cloning, Expression and Biological Characterization of an Interleukin-1 Receptor Antagonist Protein, Nature, 344:633-638 (1990).

Carpenter et al., Epidermal Growth Factor, J. Biol. Chem., 265:7709-7712 (1990).

Carpenter, Receptors for Epidermal Growth Factor and Other Polypeptide Mitogens, Ann. Rev, Biochem., 56:881-914 (1987).

Casadei et al., Expression and Secretion of Aequorin as a Chimeric Antibody by Means of a Mammalian Expression Vector, Proc. Natl. Acad. Sci., U.S.A. 87:2047-2051 (1990).

Coffman et al., The Role of Helper T Cell Products in Mouse B Cell Differentiation and Isotype Regulation, Immunol. Rev., 102:5-28 (1988).

Creasey et al., A High Molecular Weight Component of the Human Tumor Necrosis Factor Receptor is Associated With Cytotoxicity, Proc. Natl. Acad. Sci. USA, 84:3293-3297 (1987).

Dayer, Chronic Inflammatory Joint Diseases: Natural Inhibitors of Interleukin 1 and Tumor Necrosis Factor α , J. Rheumatol, 18 (Suppl. 27): 71-75 (1991).

Dower et al., Human Cytokine Receptors, J. Clin. Immunol., 10:289-299 (1990).

Eisenberg et al., Primary Structure and Functional Expression From Complementary DNA of a Human Interleukin-1 Receptor Antagonist, Nature, 343:341-346 (1990).

Ellison et al., The Nucleotide Sequence of a Human Immunoglobulin C γ 1 Gene, Nucleic Acids Res. 10(13): 4071-79 (1982).

Esmon, The Roles of Protein C and Thrombomodulin in the Regulation of Blood Coagulation, J. Biol. Chem., 264:4743-4746 (1989). European Search Report for EP 97 12 0664, dated Mar. 9, 1998.

Fell et al., Genetic Construction and Characterization of a Fusion

Protein Consisting of a Chimeric F(ab') With Specificity for Carcinomas and Human IL-2, J. Immunol., 146:2446-2452 (1991).

Fernandez-Botran et al., A Soluble, High-Affinity, Interleukin-4-Binding Protein is Present in the Biological Fluids of Mice, Proc. Natl. Acad. Sci., 87:4202-4206 (1990).

Fernandez-Botran, Soluble Cytokine Receptors: Their Role in Immunoregulation, The FASEB Journal, 5:2567-2574 (1991).

Ferrante et al., Inhibition of Tumour Necrosis Factor Alpha (TNF- α)-Induced Neutrophil Respiratory Burst by a TNF Inhibitor, Immunology, 72:440-442 (1991).

Fisher et al., Cloning and Expression of Human Tissue Factor cDNA, Thrombosis Research, 48:89-99 (1987).

Foley et al., An Inhibitor of the Toxicity of Tumour Nectrosis Factor in the Serum of Patients With Sarcoidosis, Tuberculosis and Crohn's Disease, Clin. Exp. Immunol, 80:395-399 (1990).

Fomsgaard et al., Preliminary Study on Treatment of Septic Shock Patients With Antilipopolysaccharide IgG from Blood Donors, Scand. J. Infect. Dis., 21:697-708 (1989).

Garcia et al., High Sensitivity of Transgenic Mice Expressing Soluble TNFR1 Fusion Protein to Mycobacterial Infections: Synergistic Action of TNF and IFN-γ in the Differentiation of Protective Granulomas, Eur. J. Immunol., 27:3182-3190 (1997).

Gascoigne et al., Secretion of a Chimeric T-Cell Receptor-Immunoglobulin Protein, Proc. Natl. Acad. Sci USA, 84:2936-2940 (1987).

Gehr et al., Both Tumor Necrosis Factor Receptor Types Mediate Proliferative Signals in Human Mononuclear Cell Activation, J. Immunol., 149:911-917 (1992).

Gillies et al., Targeting Human Cytotoxic T Lymphocytes to Kill Heterologous Epidermal Growth Factor Receptor-Bearing Tumor Cells, J. Immunol., 144:1067-1071 (1991).

Goodman, Identification of Antigenic Determinants, in Basic & Clinical Immunol., 24-25 (1982).

Goodman, Immunogenicity & Antigenic Specificity, in Basic & Clinical Immunol., 101-108 (1991).

Goodwin et al., Molecular cloning and Expression of the Type 1 and Type 2 Murine Receptors for Tumor Necrosis Factor, Molecular and Cellular Biology, 11:3020-3026 (1991).

Gray et al, Cloning and Expression of cDNA for Human Lymphotoxin, a Lymphokine With Tumour Necrosis Activity, Nature, 312:721-724 (1984).

Grundmann et al., Characterization of cDNA Coding for Human Factor XIIIa, Proc. Natl. Acad. Sci. USA, 83:8024-8028 (1986).

Haak-Frendscho et al., Inhibition of TNF by a TNF Receptor Immunoadhesin, J. Immunol., 152:1347-1353 (1994).

Hannum et al., Interleukin-1 Receptor Antagonist Activity of a Human Interleukin-1 Inhibitor, Nature, 343:336-340 (1990).

Heflin et al., Prevention by Granulocyte Depletion of Increased Vascular Permeability of Sheep Lung Following Endotoxemia, J. Clin. Invest., 68:1253-1260 (1981).

Himmler et al., Molecular Cloning and Expression of Human and Rat Tumor Necrosis Factor Receptor Chain (p60) and Its Soluble Derivative, Tumor Necrosis Factor-Binding Protein, DNA and Cell Biology, 9:705-715 (1990).

Hobart, The Immune System: A Course on the Molecular and Cellular Basis of immunity, Blackwell Scientific Pubs, p. 7 (1975). Holtmann et al., Down Regulation of the Receptors for Tumor Necrosis Factor by Intereukin 1 and 4β -Phorbol-12-Myristate-13-Acetate, J. Immunol., 139:1161-1167 (1987).

Idzerda et al., Human Interleukin 4 Receptor Confers Biological Responsiveness and Defines a Novel Receptor Superfamily, J. Exp. Med., 171:861-873 (1990).

Imamura et al., Expression of Tumor Necrosis Factor Receptors on Human Monocytes and Internalization of Receptor Bound Ligand, J. Immunol., 139:2989-2992 (1987).

Ishikura et al., Differential Biologic Effects Resulting From Bimodal Binding of Recombinant Human Tumor Necrosis Factor to Myeloid Leukemia Cells, Blood, 73:419-424 (1989).

Israel et al., Binding of Human TNF- α to High-Affinity Cell Surface Receptors: Effect of IFN, Immunology Letters, 12:217-224 (1986). Jacobs et al., Pharmacokinetic Parameters and Biodistribution of Soluble Catabine Receptors. International Review of Experimental

Soluble Cytokine Receptors, International Review of Experimental Pathology, 34B:123-135 (1993).

Jones et al, Structure of Tumour Necrosis Factor, Nature, 338:225-228 (1989).

Kaczmarski et al., The Cytokine Receptor Superfamily, Blood Reviews, 5:193-203 (1991).

Kaushansky, Structure-Function Relationships of the Hematopoietic Growth Factors, Proteins: Structure, Function & Genetics, 12:1-9 (1992).

Keegan et al., The Interleukin-4 Receptor. Signal Transduction by a Hematopoietin Receptor, Journal of Leukocyte Biology, 55:272-279 (1994).

Keegan et al., Interleukin 4 Receptor: Signaling Mechanisms, Immunology Today, 15:423-432 (1994).

Kleinau et al., Importance of CD23 for Collagen-Induced Arthritis: Delayed Onset and Reduced Severity in CD23-Deficient Mice, J. Immunol., 162:4266-4270 (1999).

Klinkert et al., TNF- α Receptor Fusion Protein Prevents Experimental Auto-Immune Encephalomyelitis and Demyelination in Lewis Rats: an Overview, The Journal of Neuroimmunology, 72:163-168 (1997).

Kohno et al., A Second Tumor Necrosis Factor Receptor Gene Product Can Shed a Naturally Occurring Tumor Necrosis Factor Inhibitor, Proc. Natl. Acad. Sci. USA, 87:8331-8335 (1990).

Kruse et al., Conversion of Human Interleukin-4 Into a High Affinity Antagonist by a Single Amino Acid Replacement, The EMBO Journal, 11:3237-3244 (1992).

Kull et al., Cellular Receptor for ¹²⁵I-Labeled Tumor Necrosis Factor: Specific Binding, Affinity Labeling, and Relationship to Sensitivity, Proc. Natl. Acad. Sci. USA, 82:5756-5760 (1985).

Landolfi, A Chimeric IL-2/Ig Molecule Possesses the Functional Activity of Both Proteins, J. Immunol., 146:915-919 (1991).

Langner et al., Structural and Functional Analysis of a TNF Receptor-Immunoglobulin Fusion Protein, New Advances on Cytokines, 349-354 (1992).

Leberthon et al., Enhanced Tumor Uptake of Macromolecules Induced by a Novel Vasoactive Interleukin 2 Immunoconjugate, Cancer Research, 51:2694-2698 (1991).

Lesslauer et al., Recombinant Soluble Tumor Necrosis Factor Receptor Proteins Protect Mice From Lipopolysaccharide-Induced Lethality, Eur. J. Immunol., 21:2883-2886 (1991).

Liabakk et al., A Rapid and Sensitive Immunoassay for Tumor Necrosis Factor Using Magnetic Monodisperse Polymer Particles, Journal of Immunological Methods, 134:253-259 (1990).

Loetscher et al., Efficacy of a Chimeric TNFR-IgG Fusion Protein to Inhibit TNF Activity in Animal Models of Septic Shock, Endotoxin Research Series, 2:455-462•1993).

Loetscher et al., Two distinct human TNF receptors: purification, molecular cloning and expression, in Tumor Necrosis Factor: Structure-Function Relationship and Clinical Application, (3rd International Conference, 1992.

Maliszewski et al., Cytokine Receptors and B Cell Functions: Recombinant Soluble Receptors Specifically Inhibit IL-1 and IL-4 Induced Cell Activities In Vitro, J. Immunol., 144:3028-3033 (1990). Mohler et al., Soluble Tumor Necrosis Factor (TNF) Receptors Are Effective Therapeutic Agents in Lethal Endotoxemia and Function Simultaneously as Both TNF Carriers and TNF Antagonists, J. Immunol., 151:1548-1561 (1993). Mori et al., Attenuation of Collagen-Induced Arthritis in 55-kDa TNF Receptor Type 1 (TNFR1)-IgG1-Treated and TNFR1-Deficient Mice, J. Immunol., 157:3178-3182 (1996).

Morrissey et al., Molecular Cloning of the cDNA for Tissue Factor, the Cellular Receptor for the Initiation of the Coagulation Protease Cascade, Cell: 50:129-135 (1987).

Morrison, In Vitro Antibodies: Strategies for Production and Application, Annu. Rev. Immunol., 10:239-265 (1992).

Mosley et al., The Murine Interleukin-4 Receptor: Molecular Cloning and Characterization of Secreted and Membrane Bound Forms, Cell: 59:335-348 (1989).

Novotny et al., A Soluble, Single-Chain T-Cell Receptor Fragment Endowed With Antigen-Combining Properties, Proc. Natl. Acad. Sci. USA, 88:8646-8650 (1991).

Okayama et al., High-Efficiency Cloning of Full-Length cDNA, Molecular and Cellular Biology, 2:161-170 (1982).

Okayama et al., A cDNA Cloning Vector That Permits Expression of cDNA Inserts in Mammalian Cells, Molecular and Cellular Biology, 3:280-289 (1983).

Old, Tumor Necrosis Factor, 2nd Intl Conference on Tumor Necrosis Factor & Related Cytokines, Napa, CA, 1-30 (1989).

Paborsky et al., Purification of Recombinant Human Tissue Factor, Biochemistry, 28:8072-8077 (1989).

Parrillo, Pathogenetic Mechanisms of Septic Shock, New Eng. J. Med., 328:1471-1477 (1993).

Peetre et al., A Tumor Necrosis Factor Binding Protein is Present in Human Biological Fluids, Eur. J. Haematol. 41:414-419 (1988).

Pennica et al., Human Tumour Necrosis Factor: Precursor Structure, Expression and Homology to Lymphotoxin, Nature, 312:724-729 (1984).

Piguet et al., Evolution of Collagen Arthritis in Mice is Arrested by Treatment With Anti-Tumor Necrosis (TNF) Antibody or a Recombinant Soluble TNF Receptor, Immunology, 77 (4):510-514 (1992). Redfield'et al., Secondary Structure and Topology of Human Interleukin 4 in Solution, Biochemistry, 30:11029-11035 (1991).

Rubin, Binding Receptor Characters Zako and Expression, and Intracellular Events. 2nd Intl Conference on Tumor Necrosis Factor & Related Cytokines, Napa, CA, 94-96 (1989).

Ruddle et al., An Antibody to Lymphotoxin and Tumor Necrosis Factor Prevents Transfer of Experimental Allergic Encephalomyelitis, J. Exp. Med., 172:1193-1200 (1990).

Rutka et al., The Effects of Human Recombinant Tumor Necrosis Factor on Glioma-Derived Cell Lines: Cellular Proliferation, Cytotoxicity, Morphological and Radioreceptor Studies, Int. J. Cancer., 41:573-582 (1988).

Saxne et al., Detection of Tumor Necrosis Factor α But Not Tumor Necrosis Factor β in Rheumatoid Arthritis Synovial Fluid and Serum, Arthritis & Rheumatism, 31:1041-1045 (1988).

Scallon et al., Functional Comparisons of Different Tumour Necrosis Factor Receptor/IgG Fusion Proteins, Cytokine, 7:759-770 (1995). Scarpati et al., Human Tissue Factor: cDNA Sequence and Chromosome Localization of the Gene, Biochemistry, 26:5234-5238 (1987).

Schleiffenbaum et al., The Tumor Necrosis Factor Receptor and Human Neutrophil Function, J. Clin. Invest., 86:184-195 (1990).

Schnee et al., Construction and Expression of a Recombinant Antibody-Targeted Plasminogen Activator, Proc. Natl. Acad. Sci. USA, 84:6904-6908 (1987).

Seckinger et al., A Human Inhibitor of Tumor Necrosis Factor α , J. Exp. Med. 167:1511-1516 (1988).

Shalaby et al., Receptor Binding and Activation of Polmorphonuclear Neutrophils by Tumor Necrosis Factor-Alpha, Journal of Leukocyte Biology, 41:196-204 (1987).

Shalaby et al., Binding and Regulation of Cellular Function by Monoclonal antibodies Against Human Tumor Necrosis Factor Receptors, J. Exp. Med. 172: 1517-1520 (1990).

Sheehan et al., Generation and Characterization of Hamster Monoclonal Antibodies That Neutralize Murine Tumor Necrosis Factors, Journal of Immunology, 142:3884-3893 (1989).

Shin et al., Expression and Characterization of an Antibody Binding Specificity Joined to Insulin-Like Growth Factor 1: Potential Applications for Cellular Targeting, Proc. Natl. Acad. Sci., 87:5322-5326 (1990). Sims et al., cDNA Expression Cloning of the IL-1 Receptor, a Member of the Immunoglobulin Superfamily, Science, 241:585-589 (1988).

Sims et al., Cloning the Interleukin 1 Receptor From Human T Cells, Proc. Natl. Acad. Sci., 86:8946-8950 (1989).

Smith et al., The Active Form of Tumor Necrosis Factor is a Trimer, J. Biol. Chem., 262:6951-6954 (1987).

Smith et al., Blocking of HIV-1 Infectivity by a Soluble, Secreted Form of the CD4 Antigen, Science, 238:1704-1707 (1987).

Smith et al., Multimeric Structure of the Tumor Necrosis Factor Receptor of HeLa Cells, J. Biol. Chem., 264:14646-14652 (1989).

Spicer et al., Isolation of cDNA Clones Coding for Human Tissue Factor: Primary Structure of the Protein and cDNA, Proc. Natl. Acad. Sci., 84:5148-5152 (1987).

Staines et al., Collagen Arthritis—What Can It Teach Us?, British Journal of Rheumatology, 33:798-807 (1994).

Strader et al., Structural Basis of β -Adrenergic Receptor Function, The FASEB Journal, 3:1825-1832 (1989).

Suggs et al., Use of Synthetic Oligonucleotides as Hybridization Probes: Isolation of Cloned cDNA Sequences for Human $\beta_{2,}$ -Microglobulin, Proc. Natl. Acad. Sci. U.S.A., 78:6613-6617 (1981). Tauber et al., Toxicity in Neuronal Cells Caused by Cererospinal Fluid fFom Pneumococcal and Gram-Negative Meningitis, The Journal of Infectious Diseases, 166:1045-1050 (1992).

Thoma et al., Identification of a 60-kD Tumor Necosis Factor (TNF) Receptor as the Major Signal Transducing Component in TNF Responses, J. Exp. Med. 172: 1019-23 (1990).

Tsujimoto et al., Characterization and Affinity Crosslinking of Receptors for Tumor Necrosis Factor on Human Cells, Archives of Biochemistry and Biophysics, 249:563-568 (1986).

Tsujimoto et al, Interferon-y Enhances Expression of Cellular Receptors for Tumor Necrosis Factor, J. Immunol., 136:2441-2444 (1986). Tsujimoto et al, Tumor necrosis factor: specific binding and internalization in sensitive and resistant cells, Proc. Natl. Acad. Sci. 82: 7626-30 (1985).

Ulich et al., Intratracheal Administration of Endotoxin and Cytokines, Clinical Immunology & Immunopathology., 72:137-140 (1994).

Unglaub et al., Downregulation of Tumor Necrosis Factor (TNF) Sensitivity Via Modulation of TNF Binding Capacity by Protein Kinase C Activators, J. Exp. Med. 166:1788-1797 (1987).

Van Der Poll et al., Pretreatment with a 55-kDa Tumor Necrosis Factor Receptor-Immunoglobulin Fusion Protein Attenuates Activation of Coagulation, but not of Fibrinolysis, during Lethal Bacteremia in Baboons, The Journal of Infectious Diseases., 176:296-299 (1997).

Van Zee et al., Protection Against Lethal *Escherichia coli* Bacteremia in Baboons (*Papio anubis*) by Pretreatment With a 55-kDa TNF Receptor (CD120a)-Ig Fusion Protein, Ro 45-2081, J. Immunol., 156:2221-2230 (1996).

Wallach et al., Soluble and Cell Surface Receptors for Tumor Necrosis Factor, Progress, Inflammation Research & Therapy, 51-57 (1991).

Wallach et al., Cell surface and soluble TNF receptors, in Tumor Necrosis Factor: Structure-Function Relationship and Clinical Application, (3rd International Conference on Tumor Necrosis Factor and Related Cytokines, Makuhari, Chiba, Nov. 21-25, 1990), Osawa and Bonavida, eds., Basel, Karger, pp. 47-57 (1992).

Wilks, The CD4 Receptor: Post Binding Events, Conformational Change and the Second Site, Molec. Aspects Med., 12:255-265 (1991).

Yamasaki et al., Cloning and Expression of the Human Interleukin-6 (BSF-2/IFNβ 2) Receptor, Science, 241:825-282 (1988).

Yonehara et al., A Cell-Killing Monoclonal Antibody (Anti-Fas) to a Cell Surface Antigen Co-Downregulated With the Receptor of Tumor Necrosis Factor, J. Exp. Med., 169:1747-1765 (1989).

Yoshie et al., Binding and Crosslinking of ¹²⁵I-Labeled Recombinant Human Tumor Necrosis Factor to Cell Surface Receptors, J. Biochem., 100: 531-541(1986).

Official Communication relating to an Opposition in EP Application No. 99 100 703.0, Jan. 31, 2006.

H. Engelmann et al., Jour. of Biological Chemistry, vol. 264, Nr. 20, p. 11974-11980, (Jul. 1989).

P.W. Gray et al, Proc. of the Nat'l Acad. of Sci., vol. 87, p. 7380-7884, (Oct. 1990).

H. Engelmann et al., Jour. of Biological Chemistry, vol. 264, p. 531-1536, (Jan. 1990).

H. Loetscher et al,. Cell, vol. 61, p. 351-359 (Apr. 1990).

T.J. Schall et al., Cell, vol. 61, p. 361-370 (Apr. 1990).

R. A. Heller et al., Proc. of the Nat'l Acad. of Sci., vol. 87, pp. 6151-6155, (Aug. 1990).

C. A. Smith et al, Science, vol. 248, pp. 1019-1023, (May 1990).

P. Seckinger et al, Eur. J. Immunology, vol. 20, pp. 1167-1174 (1990). D. Novick et al, J. Exp. Med., vol. 170, pp. 1409-1414, (1988).

P. Seckinger et al., Jour. of Biological Chem., vol. 264, pp. 22966-11973 (Jul. 1989).

Folks et al., Tumor necrosis factor α induces expression of human immunodeficiency virus in a chronically infected T-cell clone, PNAS, 86:2365-8 (1989).

Lee, Generation of cDNA probes directed by amino acid sequence: cloning of urate oxidase, Science, 239:1288-1291 (1988).

Moreland et al., Treatment of rheumatoid arthritis with a recombinant human tumor necrosis factor receptor (p75)-Fc fusion protein, New England Journal of Medicine, 337:141-147 (1997).

Sandborn et al., Etanercept for active Crohn's disease: a randomized, double-blind, placebo-controlled trial, Gastroenterol., 121:1088-94 (2001).

Wozney, Using purified protein to clone its genes, Methods in Enzymology, 182:738-751 (1990).

Inflammation & Infection: Rheumatoid Arthritis, Drug & Market Development, 67-70 (2003).

Letter to the EPO with new Auxiliary request 3 in appeal proc. against EP 0 471 701, Dec. 31, 2001.

Notification of the Examiner during the prosecution of EP 99100703. 0, Oct. 24, 2000.

Zavoico, Drug & Market Development, 10(7):235-243 (1999).

Heller et al., Abstract at the 2nd International Conference on Tumor Necrosis Factor and Related Cytokines, Napa Valley, California, Jan. 15-20, 1989.

Loetscher et al., Two Distinct Human TNF Receptors: Purification, Molecular Cloning and Expression, 3rd International Conference on Tumor Necrosis Factor and Related Cytokines, Makuhari, Chiba, Japan, Nov. 21-25, 1990.

Inflammation & Infection: Rheumatoid Arthritis, Drug & Market Development, 67-70 (Mar. 2003).

Barone et al., Arthritis Rheum., 42(9)Suppl, Sep. 1999, Abstract 116. IPS Notice of Opposition to EP 0 939 121 and Statement in Support of the Opposition against EP 0939121, Dec. 30, 2003.

Patentee's Response to Notice of Opposition Request for Oral Hearing, dated Aug. 3, 2004 (with translation).

Answer of the Opponent on the Admissibility of the Oppositions, dated Nov. 3, 2004.

Response to Patentee's Written Submission, dated Jul. 22, 2005.

Opponent's Observations Under R 71a EPC, dated Sep. 22, 2006. AHP Comments on Oral Hearing and Preliminary Opinion, dated Sep. 22, 2006.

AHP Response to Opponent's Observations, dated Oct. 6, 2006. US 6,224,867, 05/2001, Smith et al. (withdrawn)

* cited by examiner

<u>Figure 1</u>

	GAATTEGEGEGEGTTEAAGATEAETEGEAECEAGECESTEATETETATGEEEGAGTETEAA
-125	CCCTCAACTGTCACCCCAAGGCACTTGGGACGTCCTGGACAGACCGAGTCCCGGGAAGCC
-65	CCAGCACTGCCGCTGCCACACTGCCCTGAGCCCAAATGGGGGAGTGAGAGGCCATAGCTG
	-28
-30	Not Club ou Sor Thria) Prodent out ou Protente 2 Leuteu Cluber
-50	
-3	
-10	LeuValGlyIleTyrProSerGlyValIleGlyLeuValProHisLeuGlyAspArgGlu
55	TTGGTGGGAATATACCCCTCAGGGGTTATTGGACTGGTCCCTCACCTAGGGGACAGGGAG
10	LysArcAspSerValCysProGlnGlyLysTyrIleHisProGlnAsnAsnSerIleCys
115	AAGAGAGATAGTGTGTGTGTCCCCAAGGAAAATATATCCACCCTCAAAATAATTCGATTTGC
20	Cuember up Cuelli at up Clumber up comund and an Cuellan Clubban Clubban
175	
112	TO TALLAAD TO LALAAAG MALLAATTO TALAATGALTO TLAGOLLUGOOGAGAT
50	ThrAspCysArgGluCysGluSerGlySerPheThrAlaSerGluAsnHisLeuArgHis
235	ACEGACTECAGEGAGTETEAGAGCEGCTCCTTCACCECTTCAGAAAACCACCTCAGACAC
70	CvsLeuSerCvsSerLvsCvsArgLvsGluMetGlvGlnValGluIleSerSerCvsThr
295	TECCTCACCTECTAATECCEAAAGCAAATEGETCAGETEGAGATETETTECTECACA
20	
20	
300	
	*** ***
110	AsnLeuPheGlnCysPheAsnCysSerLeuCysLeuAsnGlyThrValHisLeuSerCys
415	AACCTTTTCCAGTGCTTCAATTGCAGCCTCTGCCTCAATGGGACCGTGCACCTCTCCTGC
130	GInGluLysGInAsnThrValCysThrCysHisAlaGlyPhePheLeuArcGluAsnGlu
475	CAGGAGAAACAGAACACGTGTGCACCTGCCATGCAGGTTTCTTTC
150	
100	CueVal SarCueSarAenCuel uel ueSart auClucueThri uel auCuel auProcla
575	CysValSerCysSerAsnCysLysLysSerLeuGluCysThrLysLeuCysLeuProGln
-535	CysValSerCysSerAsnCysLysLysSerLeuGluCysThrLysLeuCysLeuProGln TGTGTCTCCTGTAGTAACTGTAAGAAAAGCCTGGAGTGCACGAAGTTGTGCCTACCCCAG
-535	CysValSerCysSerAsnCysLysLysSerLeuGluCysThrLysLeuCysLeuProGln TGTGTCTCCTGTAGTAACTGTAAGAAAAGCCTGGAGTGCACGAAGTTGTGCCTACCCCAG
535 170	CysValSerCysSerAsnCysLysLysSerLeuGluCysThrLysLeuCysLeuProGln TGTGTCTCCTGTAGTAACTGTAAGAAAAGCCTGGAGTGCACGAAGTTGTGCCTACCCCAG IleGluAsnValLysGlyThrGluAspSerGlyThrThr <u>ValLeuLeuProLeuValLle</u>
-535 170 595	CysValSerCysSerAsnCysLysLysSerLeuGluCysThrLysLeuCysLeuProGln TGTGTCTCCTGTAGTAACTGTAAGAAAAGCCTGGAGTGCACGAAGTTGTGCCTACCCCAG IleGluAsnValLysGlyThrGluAspSerGlyThrThr <u>ValLeuLeuProLeuValLle</u> ATTGAGAATGTTAAGGGCACTGAGGACTCAGGCACCACAGTGCTGTCGCCCCTGGTCATT
-535 170 595	CysValSerCysSerAsnCysLysLysSerLeuGluCysThrLysLeuCysLeuProGln TGTGTCTCCTGTAGTAACTGTAAGAAAAGCCTGGAGTGCACGAAGTTGTGCCTACCCCAG IleGluAsnValLysGlyThrGluAspSerGlyThrThr <u>ValLeuLeuProLeuValLle</u> ATTGAGAATGTTAAGGGCACTGAGGACTCAGGCACCACAGTGCTGTTGCCCCTGGTCATT
-535 170 595 190	CysValSerCysSerAsnCysLysLysSerLeuGluCysThrLysLeuCysLeuProGln TGTGTCTCCTGTAGTAACTGTAAGAAAAAGCCTGGAGTGCACGAAGTTGTGCCTACCCCAG IleGluAsnValLysGlyThrGluAspSerGlyThrThr <u>ValLeuLeuProLeuValLle</u> ATTGAGAATGTTAAGGGCACTGAGGACTCAGGCACCACAGTGCTGTTGCCCCTGGTCATT <u>PhePheGlyLeuCysLeuLeuSerLeuLeuPheIleGlyLeuMetTyrArgTyrGlnArg</u>
-535 170 595 190 655	CysValSerCysSerAsnCysLysLysSerLeuGluCysThrLysLeuCysLeuProGln TGTGTCTCCTGTAGTAACTGTAAGAAAAGCCTGGAGTGCACGAAGTTGTGCCTACCCCAG IleGluAsnValLysGlyThrGluAspSerGlyThrThr <u>ValLeuLeuProLeuValLe</u> ATTGAGAATGTTAAGGGCACTGAGGACTCAGGCACCACAGTGCTGTTGCCCCTGGTCATT <u>PhePheGlyLeuCysLeuLeuSerLeuLeuPheIle</u> GlyLeuMetTyrArgTyrGlnArg
-535 170 595 190 655	CysValSerCysSerAsnCysLysLysSerLeuGluCysThrLysLeuCysLeuProGln TGTGTCTCCTGTAGTAACTGTAAGAAAAGCCTGGAGTGCACGAAGTTGTGCCTACCCCAG IleGluAsnValLysGlyThrGluAspSerGlyThrThr <u>ValLeuLeuProLeuValLle</u> ATTGAGAATGTTAAGGGCACTGAGGACTCAGGCACCACAGTGCTGTTGCCCCTGGTCATT <u>PhePheGlyLeuCysLeuLeuSerLeuLeuPheIle</u> GlyLeuMetTyrArgTyrGlnArg TTCTTTGGTCTTTGCCTTTTATCCCTCCTCTTCATTGGTTTAATGTATCGCTACCCAACGG
-535 170 595 190 655	CysValSerCysSerAsnCysLysLysSerLeuGluCysThrLysLeuCysLeuProGln TGTGTCTCCTGTAGTAACTGTAAGAAAAGCCTGGAGTGCACGAAGTTGTGCCTACCCCAG IleGluAsnValLysGlyThrGluAspSerGlyThrThr <u>ValLeuProLeuValLe</u> ATTGAGAATGTTAAGGGCACTGAGGACTCAGGCACCACAGTGCTGTTGCCCCTGGTCATT <u>PhePheGlyLeuCysLeuLeuSerLeuLeuPheIle</u> GlyLeuMetTyrArgTyrGlnArg TTCTTTGGTCTTTGCCTTTTATCCCTCCTCTTCATTGGTTTAATGTATCGCTACCAACGG
-535 170 595 190 655 210	CysValSerCysSerAsnCysLysLysSerLeuGluCysThrLysLeuCysLeuProGln TGTGTCTCCTGTAGTAACTGTAAGAAAAGCCTGGAGTGCACGAAGTTGTGCCTACCCCAG IleGluAsnValLysGlyThrGluAspSerGlyThrThr <u>ValLeuLeuProLeuValLe</u> ATTGAGAATGTTAAGGGCACTGAGGACTCAGGCACCACAGTGCTGTCGCCCTGGTCATT <u>PhePheGlyLeuCysLeuLeuSerLeuLeuPheIle</u> GlyLeuMetTyrArgTyrGlnArg TTCTTTGGTCTTTGCCTTTTATCCCTCCTCTTCATTGGTTTAATGTATCGCTACCAACGG
-535 170 595 190 655 210 715	CysValSerCysSerAsnCysLysLysSerLeuGluCysThrLysLeuCysLeuProGln TGTGTCTCCTGTAGTAACTGTAAGAAAAGCCTGGAGTGCACGAAGTTGTGCCTACCCCAG IleGluAsnValLysGlyThrGluAspSerGlyThrThr <u>ValLeuLeuProLeuValLe</u> ATTGAGAATGTTAAGGGCACTGAGGACTCAGGCACCACAGTGCTGTCGCCCCTGGTCATT <u>PhePheGlyLeuCysLeuLeuSerLeuLeuPheIle</u> GlyLeuMetTyrArgTyrGlnArg TTCTTTGGTCTTTGCCTTTTATCCCTCCTCTTCATTGGTTTAATGTATCGCTACCAACGG TrpLysSerLysLeuTyrSerIleValCysGlyLysSerThrProGluLysGluGlyGlu TGGAAGTCCAAGCTCTACTCCATTGTTTGTGGGGAAATCGACACCTGAAAAAGAGGGGGGAG
-535 170 595 190 655 210 715	CysValSerCysSerAsnCysLysLysSerLeuGluCysThrLysLeuCysLeuProGln TGTGTCTCCTGTAGTAACTGTAAGAAAAGCCTGGAGTGCACGAAGTTGTGCCTACCCCAG IleGluAsnValLysGlyThrGluAspSerGlyThrThr <u>ValLeuLeuProLeuValLle</u> ATTGAGAATGTTAAGGGCACTGAGGACTCAGGCACCACAGTGCTGTCGCCCCTGGTCATT <u>PhePheGlyLeuCysLeuLeuSerLeuLeuPheIle</u> GlyLeuMetTyrArgTyrGlnArg TTCTTTGGTCTTTGCCTTTTATCCCTCCTCTTCATTGGTTTAATGTATCGCTACCAACGG TrpLysSerLysLeuTyrSerIleValCysGlyLysSerThrProGluLysGluGlyGlu TGGAAGTCCAAGCTCTACTCCATTGTTTGTGGGGAAATCGACACCTGAAAAAGAGGGGGGAG
-535 170 595 190 655 210 715 230	CysValSerCysSerAsnCysLysLysSerLeuGluCysThrLysLeuCysLeuProGln TGTGTCTCCTGTAGTAACTGTAAGAAAAGCCTGGAGTGCACGAAGTTGTGCCTACCCCAG IleGluAsnValLysGlyThrGluAspSerGlyThrThr <u>ValLeuLeuProLeuValLle</u> ATTGAGAATGTTAAGGGCACTGAGGACTCAGGCACCACAGTGCTGTCGCCCCTGGTCATT <u>PhePheGlyLeuCysLeuLeuSerLeuLeuPheIle</u> GlyLeuMetTyrArgTyrGlnArg TTCTTTGGTCTTTGCCTTTTATCCCTCCTCTTCATTGGTTTAATGTATCGCTACCAACGG TrpLysSerLysLeuTyrSerIleValCysGlyLysSerThrProGluLysGluGlyGlu TGGAAGTCCAAGCTCTACTCCATTGTTGTGGGAAATCGACACCTGAAAAAGAGGGGGAG ***
535 170 595 190 555 210 715 230 775	CysValSerCysSerAsnCysLysLysSerLeuGluCysThrLysLeuCysLeuProGln TGTGTCTCCTGTAGTAACTGTAAGAAAAGCCTGGAGTGCACGAAGTTGTGCCTACCCCAG IleGluAsnValLysGlyThrGluAspSerGlyThrThr <u>ValLeuLeuProLeuValLle</u> ATTGAGAATGTTAAGGGCACTGAGGACTCAGGCACCACAGTGCTGTTGCCCCTGGTCATT <u>PhePheGlyLeuCysLeuLeuSerLeuLeuPheIle</u> GlyLeuMetTyrArgTyrGlnArg TTCTTTGGTCTTTGCCTTTTATCCCTCCTCTTCATTGGTTTAATGTATCGCTACCAACGG TrpLysSerLysLeuTyrSerIleValCysGlyLysSerThrProGluLysGluGlyGlu TGGAAGTCCAAGCTCTACTCCATTGTTTGTGGGGAAATCGACACCTGAAAAAGAGGGGGGAG *** LeuGluGlyThrThrThrLysProLeuAlaPrcAsnProSerPheSerProThrProGly CTTGAAGGAACTACTACTAAGCCCCTGGCCCCAAACCCAAGCTTCAGTCCCACTCCAGGC
535 170 595 190 555 210 715 230 775	CysValSerCysSerAsnCysLysLysSerLeuGluCysThrLysLeuCysLeuProGln TGTGTCTCCTGTAGTAACTGTAAGAAAAGCCTGGAGTGCACGAAGTTGTGCCTACCCCAG IleGluAsnValLysGlyThrGluAspSerGlyThrThr <u>ValLeuLeuProLeuValLle</u> ATTGAGAATGTTAAGGGCACTGAGGACTCAGGCACCACAGTGCTGTTGCCCCTGGTCATT <u>PhePheGlyLeuCysLeuLeuSerLeuLeuPheIle</u> GlyLeuMetTyrArgTyrGlnArg TTCTTTGGTCTTTGCCTTTTATCCCTCCTCTTCATTGGTTTAATGTATCGCTACCAACGG TrpLysSerLysLeuTyrSerIleValCysGlyLysSerThrProGluLysGluGlyGlu TGGAAGTCCAAGCTCTACTCCATTGTTTGTGGGGAAATCGACACCTGAAAAAGAGGGGGGAG ***
535 170 595 190 555 210 715 230 775 250	CysValSerCysSerAsnCysLysLysSerLeuGluCysThrLysLeuCysLeuProGln TGTGTCTCCTGTAGTAACTGTAAGAAAAGCCTGGAGTGCACGAAGTTGTGCCTACCCCAG IleGluAsnValLysGlyThrGluAspSerGlyThrThr <u>ValLeuLeuProLeuValLie</u> ATTGAGAATGTTAAGGGCACTGAGGACTCAGGCACCACAGTGCTGTTGCCCCTGGTCATT <u>PhePheGlyLeuCysLeuLeuSerLeuLeuPheIle</u> GlyLeuMetTyrArgTyrGlnArg TTCTTTGGTCTTTGCCTTTTATCCCTCCTCTTCATTGGTTTAATGTATCGCTACCAACGG TrpLysSerLysLeuTyrSerIleValCysGlyLysSerThrProGluLysGluGlyGlu TGGAAGTCCAAGCTCTACTCCATTGTTTGTGGGGAAATCGACACCTGAAAAAGAGGGGGGAG *** LeuGluGlyThrThrThrLysProLeuAlaPrcAsnProSerPheSerProThrProGly CTTGAAGGAACTACTACTAAGCCCCTGGCCCCAAACCCAAGCTTCAGTCCCACTCCAGGC
535 170 595 190 655 210 715 230 775 250 835	CysValSerCysSerAsnCysLysLysSerLeuGluCysThrLysLeuCysLeuProGln TGTGTCTCCTGTAGTAACTGTAAGAAAAGCCTGGAGTGCACGAAGTTGTGCCTACCCCAG IleGluAsnValLysGlyThrGluAspSerGlyThrThr <u>ValLeuLeuProLeuValLe</u> ATTGAGAATGTTAAGGGCACTGAGGACTCAGGCACCACAGTGCTGTTGCCCCTGGTCATT <u>PhePheGlyLeuCysLeuLeuSerLeuLeuPheIleG</u> lyLeuMetTyrArgTyrGlnArg TTCTTTGGTCTTTGCCTTTTATCCCTCCTCTTCATTGGTTTAATGTATCGCTACCAACGG TrpLysSerLysLeuTyrSerIleValCysGlyLysSerThrProGluLysGluGlyGlu TGGAAGTCCAAGCTCTACTCCATTGTTTGTGGGAAATCGACACCTGAAAAAGAGGGGGAG *** LeuGluGlyThrThrThrLysProLeuAlaPrcAsnProSerPheSerProThrProGly CTTGAAGGAACTACTACTACTGCCTGGCCCCAAACCCAAGCTTCAGTCCCACTCCAGGC PheThrProThrLeuGlyPheSerPrcValProSerSerThrPheThrSerSerSerThr
535 170 595 190 655 210 715 230 715 230 775 250 835	CysValSerCysSerAsnCysLysLysSerLeuGluCysThrLysLeuCysLeuProGln TGTGTCTCCTGTAGTAACTGTAAGAAAAGCCTGGAGTGCACGAAGTTGTGCCTACCCCAG IleGluAsnValLysGlyThrGluAspSerGlyThrThr <u>ValLeuLeuProLeuValLe</u> ATTGAGAATGTTAAGGGCACTGAGGACTCAGGCACCACAGTGCTGTTGCCCCTGGTCATT <u>PhePheGlyLeuCysLeuLeuSerLeuLeuPheIle</u> GlyLeuMetTyrArgTyrGlnArg TTCTTTGGTCTTTGCCTTTTATCCCTCCTCTTCATTGGTTTAATGTATCGCTACCAACGG TrpLysSerLysLeuTyrSerIleValCysGlyLysSerThrProGluLysGluGlyGlu TGGAAGTCCAAGCTCTACTCCATTGTTTGTGGGAAATCGACACCTGAAAAAGAGGGGGAG *** LeuGluGlyThrThrThrLysProLeuAlaPrcAsnProSerPheSerProThrProGly CTTGAAGGAACTACTACTACAGCCCTGGCCCCAAACCCAAGCTTCAGTCCCACTCCAGGC PheThrProThrLeuGlyPheSerProValProSerSerThrPheThrSerSerSerThr TTCACCCCCACCTGGGCTTCAGTCCCGTGCCCAGTTCCACCTCCAGCTCCACC
535 170 595 190 555 210 715 230 775 230 775 250 835	CysValSerCysSerAsnCysLysLysSerLeuGluCysThrLysLeuCysLeuProGln TGTGTCTCCTGTAGTAACTGTAAGAAAAGCCTGGAGTGCACGAAGTTGTGCCTACCCCAG IleGluAsnValLysGlyThrGluAspSerGlyThrThr <u>ValLeuLeuProLeuValIle</u> ATTGAGAATGTTAAGGGCACTGAGGACTCAGGCACCACAGTGCTGTTGCCCCTGGTCATT <u>PhePheGlyLeuCysLeuLeuSerLeuLeuPheIle</u> GlyLeuMetTyrArgTyrGlnArg TTCTTTGGTCTTTGCCTTTTATCCCTCCTCTTCATTGGTTTAATGTATCGCTACCAACGG TrpLysSerLysLeuTyrSerIleValCysGlyLysSerThrProGluLysGluGlyGlu TGGAAGTCCAAGCTCTACTCCATTGTTGTGGGGAAATCGACACCTGAAAAAGAGGGGGAG *** LeuGluGlyThrThrThrLysProLeuAlaPrcAsnProSerPheSerProThrProGly CTTGAAGGAACTACTACTACAGCCCTGGCCCCAAACCCAAGCTTCAGTCCCACTCCAGGC PheThrProThrLeuGlyPheSerProValProSerSerThrPheThrSerSerSerThr TTCACCCCCACCCTGGGCTTCAGTCCCGTGCCCAGTTCCACCTCCAGCTCCACC
535 170 595 190 555 210 715 230 775 250 835 270	CysValSerCysSerAsnCysLysLysSerLeuGluCysThrLysLeuCysLeuProGin TGTGTCTCCTGTAGTAACTGTAAGAAAAGCCTGGAGTGCACGAAGTTGTGCCTACCCCAG IleGluAsnValLysGlyThrGluAspSerGlyThrThr <u>ValLeuLeuProLeuValLe</u> ATTGAGAATGTTAAGGGCACTGAGGACTCAGGCACCACAGTGCTGTTGCCCCTGGTCATT <u>PhePheGlyLeuCysLeuLeuSerLeuLeuPheIle</u> GlyLeuMetTyrArgTyrGlnArg TTCTTTGGTCTTTGCCTTTTATCCCTCCTCTTCATTGGTTTAATGTATCGCTACCAACGG TrpLysSerLysLeuTyrSerIleValCysGlyLysSerThrProGluLysGluGlyGlu TGGAAGTCCAAGCTCTACTCCATTGTTGTGGGGAAATCGACACCTGAAAAAGAGGGGGAG *** LeuGluGlyThrThrThrLysProLeuAlaProAsnProSerPheSerProThrProGly CTTGAAGGAACTACTACTAAGCCCCTGGCCCCAAACCCAAGCTTCAGTCCCACTCCAGGC PheThrProThrLeuGlyPheSerProValProSerSerThrPheThrSerSerSerThr TTCACCCCCCACCCTGGGCTTCAGTCCCGTGCCCCAGTTCCACCTTCACCTCCAGCTCCACC
535 170 595 190 555 210 715 230 775 250 835 270 895	CysValSerCysSerAsnCysLysLysSerLeuGluCysThrLysLeuCysLeuProGln TGTGTCTCCTGTAGTAACTGTAAGAAAAGCCTGGAGTGCACGAAGTTGTGCCTACCCCAG IleGluAsnValLysGlyThrGluAspSerGlyThrThr <u>ValLeuLeuProLeuValIle</u> ATTGAGAATGTTAAGGGCACTGAGGACTCAGGCACCACAGTGCTGTGCCCCTGGTCATT <u>PhePheGlyLeuCysLeuLeuSerLeuLeuPheIle</u> GlyLeuMetTyrArgTyrGlnArg TTCTTTGGTCTTTGCCTTTTATCCCTCCTCTTCATTGGTTTAATGTATCGCTACCAACGG TrpLysSerLysLeuTyrSerIleValCysGlyLysSerThrProGluLysGluGlyGlu TGGAAGTCCAAGCTCTACTCCATTGTTTGTGGGGAAATCGACACCTGAAAAAGAGGGGGGAG *** LeuGluGlyThrThrThrLysProLeuAlaPrcAsnProSerPheSerProThrProGly CTTGAAGGAACTACTACTACAGCCCTGGCCCCAAACCCAAGCTTCAGTCCCACTCCAGGC PheThrProThrLeuGlyPheSerProValProSerSerThrPheThrSerSerSerThr TTCACCCCCACCCTGGGCTTCAGTCCCGTGCCCAGTTCCACCTCCAGCTCCACC
535 170 595 190 555 210 715 230 775 250 835 270 895	CysValSerCysSerAsnCysLysLysSerLeuGluCysThrLysLeuCysLeuProGln TGTGTCTCCTGTAGTAACTGTAAGAAAAGCCTGGAGTGCACGAAGTTGTGCCTACCCCAG IleGluAsnValLysGlyThrGluAspSerGlyThrThr <u>ValLeuLeuProLeuValLie</u> ATTGAGAATGTTAAGGGCACTGAGGACTCAGGCACCACAGTGCTGTTGCCCCTGGTCATT <u>PhePheGlyLeuCysLeuLeuSerLeuLeuPheIleG</u> lyLeuMetTyrArgTyrGlnArg TTCTTTGGTCTTTGCCTTTTATCCCTCCTCTTCATTGGTTTAATGTATCGCTACCAACGG TrpLysSerLysLeuTyrSerIleValCysGlyLysSerThrProGluLysGluGlyGlu TGGAAGTCCAAGCTCTACTCCATTGTTTGTGGGGAAATCGACACCTGAAAAAGAGGGGGGAG *** LeuGluGlyThrThrThrLysProLeuAlaProAsnProSerPheSerProThrProGly CTTGAAGGAACTACTACTAAGCCCCTGGCCCCAAACCCAAGCTTCAGTCCCACTCCAGGC PheThrProThrLeuGlyPheSerProValProSerSerThrPheThrSerSerSerThr TTCACCCCCACCCTGGGCTTCAGTCCCGTGCCCAGTTCCACCTTCACCTCCAGGC CyrThrProGlyAspCysProAsnPheAlaAlaProArgArgGluValAlaProProTyr TATACCCCCGGTGACTGTCCCCACTTTGCGGCTCCCCGCAGAGGGGGGCACCACCCTAT
535 170 595 190 555 210 715 230 715 230 775 250 835 270 895 290	CysValSerCysSerAsnCysLysLysSerLeuGluCysThrLysLeuCysLeuProGln TGTGTCTCCTGTAGTAACTGTAAGAAAAAGCCTGGAGTGCACGAAGTTGTGCCTACCCCAG IleGluAsnValLysGlyThrGluAspSerGlyThrThr <u>ValLeuLeuProLeuValLle</u> ATTGAGAATGTTAAGGGCACTGAGGACTCAGGCACCACAGTGCTGTTGCCCCTGGTCATT <u>PhePheGlyLeuCysLeuLeuSerLeuLeuPheIle</u> GlyLeuMetTyrArgTyrGlnArg TTCTTTGGTCTTTGCCTTTTATCCCTCCTCTTCATTGGTTTAATGTATCGCTACCAACGG TrpLysSerLysLeuTyrSerIleValCysGlyLysSerThrProGluLysGluGlyGlu TGGAAGTCCAAGCTCTACTCCATTGTTGTGGGGAAATCGACACCTGAAAAAGAGGGGGGAG *** LeuGluGlyThrThrThrLysProLeuAlaProAsnProSerPheSerProThrProGly CTTGAAGGAACTACTACTACTAAGCCCCTGGCCCCAAACCCAAGCTTCAGTCCCACTCCAGGC PheThrProThrLeuGlyPheSerProValProSerSerThrPheThrSerSerSerThr TTCACCCCCACCCTGGGCTTCAGTCCCGTGCCCCAGTTCCACCTCCAGCTCCACC TyrThrProGlyAspCysProAsnPheAlaAlaProArgArgGluValAlaProProTyr TATACCCCCGGTGACTGTCCCAACTTTGCGGGCTCCCCGCAGAGAGGTGGCACCACCCTAT GlnGlyAlaAspProIleLeuAlaThrAlaLeuAlaSerAspProIleProAsnProLeu

<u>Figure 1 (cont.</u>

310	GlnLysTrpGluAspSerAlaHisLysProGlnSerLeuAspThrAspAspProAlaThr
1015	CAGAAGTGGGAGGACAGCGCCCACAAGCCACAGAGCCTAGACACTGATGACCCCGCGACG
330	LeuTyrAlaValValGluAsnValProProLeuArgTrpLysGluPheValArgArgLeu
1075	CTGTACGCCGTGGTGGAGAACGTGCCCCCGTTGCGCTGGAAGGAA
350	GlyLeuSerAspHisGluIleAspArgLeuGluLeuGlnAsnGlyArgCysLeuArgGlu
1 135	GGGCTGAGCGACCACGAGATCGATCGGCTGGAGCTGCAGAACGGGCGCTGCCTGC
370	AlaGInTyrSerMetLeuAlaThrTrpArgArgArgThrProArgArgGluAlaThrLeu
1195	GCGCAATACAGCATGCTGGCGACCTGGAGGCGGCGCGCGC
390	GluLeuLeuGlyArgValLeuArgAspMetAspLeuLeuGlyCysLeuGluAspIleGlu
1255	GAGCTGCTGGGACGCGTGCTCCGCGACATGGACCTGCTGGGCTGCCTGGAGGACATCGAG
410	GlualaLeuCysGlyProAlaAlaLeuProProAlaProSerLeuLeuArg
1315	GAGGGGGTTTGCGGGCCCGGCGGCCCGCGGGGCCCAGTCTTCTCAGATGAGGCTGC
1375	GCCCCTGCGGGCAGCTCTAAGGACCGTCCTGCGAGATCGCCTTCCAACCCCACTTTTTTC
1435	TGGAAAGGAGGGGTCCTGCAGGGGCAAGCAGGAGCTAGCAGCCGCCTACTTGGTGCTAAC
1495	CCCTCGATGTACATAGCTTTTCTCAGCTGCCTGCGCGCCGCCGACAGTCAGCGCTGTGCG
1555	CGCGGAGAGAGGTGCGCCGTGGGCTCAAGAGCCTGAGTGGGTGG
1615 1675 1735	ACGCTATGCCTCATGCCCGTTTTGGGTGTCCTCACCAGCAAGGCTGCTCGGGGGGCCCCTG GTTCGTCCCTGAGCCTTTTTCACAGTGCATAAGCAGTTTTTTTT
1795	CCTGGACAAGCACATAGCAAGCTGAACTGTCCTAAGGCAGGGGGGGAGCACGGAACAATGG
1855	GGCCTTCAGCTGGAGCTGTGGACTTTTGTACATACACTAAAATTCTGAAGTTAAAAAAA
1915	AACCCGAATTC





Binding/Cell

Figure 3



Sandwich - Assay

Figure 4

1	SerfispSerValCysAspSerCysGluAspSerThrTyrThrGlnLeuTrpAsnTrpVal
1	TCGGACTCCGTGTGTGACTCCTGTGAGGACAGCACATACACCCAGCTCTGGAACTGGGTT
21	ProüluCysLeuSerCysGlySerArgCysSerSerAspülnValGluThrGlnAlaCys
61	CCCGRGTGCTTGAGCTGTGGCTCCCGCTGTAGCTCTGACCAGGTGGAAACTCRAGCCTGC
41	ThrArgGluGlnAsnArglleCysThrCysArgProGlyTrpTyrCysAlaLeuSerLys
121	ACTCGGGAACAGAACCGCATCTGCACCTGCAGGCCCGGCTGGTACTGCGCGCTGAGCAAG
61	GInGIuGIyCysArgLeuCysAlaProLeuProLysCysArgProGIyPheGIyValAla
181	CAGGAGGGGGGGCGGCGGCGCGCGCGGGGGGCTTCGGCGGGGCGGGCC
81	ArgProOlyThrOluThrSerRspValValCysLysProCysAlaProOlyThrPheSer
241	,AGACCAGGAACTGAAACATCAGACGTGGTGTGCAAGCCCTGTGCCCCGGGGACGTTCTCC
101	AsnThrThrSerSerThrAspileCysArgProHisGinlleCysAsnUalUalAlaile
301	AACACGACTTCATCCACGGATATTTGCAGGCCCCACCAGATCTGTAACGTGGTGGCCATC
121	ProGlyAsnAlaSerArgAspAlaUalCysThrSerThrSerProThrArgSerMetAla
361	CCTGGGRATGCAAGCAGGGATGCAGTCTGCACGTCCACGTCCCCCACCCGGAGTATGGCC
141	ProGlyAlaUalHisLeuProGlnProUalSerThrArgSerGlnHisThrGlnProSer
421	CCAGGGGGCAGTACACTTACCCCAGCCAGTGTCCACACGATCCCAACACACGCAGCCAAGT
161	ProGluProSerThrAlaProSerThrSerPheLeuLeuProMetGlyProSerProPro
481	CCRGRACCCRGCACTGCTCCRAGCACCTCCTTCCTGCTCCCARTGGGCCCCRGCCCCCCA
181	AlaGluGlySerThrGlyAspPheAlaLeuPraValGlyLeulleValGlyValThrAla
541	GCTGAAGGGAGCACTGGCGACTTCGCTCTTCCAGTTGGACTGATTGTGGGTGTGACAGCC
201	LeuGlyLeuLeulleileGlyValValAsnCysVallleNetThrGinValLysLysLys
601	TTGGGTCTACTAATAATAGGAGTGGTGGAACTGTGTCATCATGACCCAGGTGAAAAAGAAG
221	ProLeuCysLeuGlnArgGluAlaLysUalProHisLeuProAlaAspLysAlaArgGly
661	CCCTTGTGCCTGCAGAGAGAGAGCCAAGGTGCCTCACTTGCCTGCC
241	ThrGinGlyProGluGinGinHisLeuLeulieThrAlaProSerSerSerSerSer
721	ACACAGGGGCCCCGAGCAGCAGCAGCACCTGCTGATCACAGCGCGCCGAGCTCCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAG
261	LeuGluSerSerAlaSerAlaLeuAspArgArgAlaProThrArgAsnGlnProGlnAla
781	CTGGRGAGCTCGGCCAGTGCGTTGGACAGARGGGCGCCCACTCGGAACCAGCCACAGGCA

Figure 4 (cont.)

281	ProûlyValGluAlaSerûlyAlaGlyGluAlaArgAlaSerThrGlySerSerAlaA so
841	CCAGGCGTGGAGGCCAGTGGGGGCCGGGGAGGCCCGGGCCAGCACCGGGAGCTCAGCAGAT
301	SerSerProGlyGlyHisGlyThrGInValAsnValThrCysIleValAsnValCysSer
901	TCTTCCCCTGGTGGCCRTGGGACCCAGGTCAATGTCACCTGCATCGTGAACGTCTGTAGC
321	SerSerAspHisSerSerGlnCysSerSerGlnAlaSerSerThrMetGlyAspThrAsp
961	AGCTCTGACCACAGCTCACAGTGCTCCTCCCARGCCAGCTCCACAATGGGAGACACAGAT
341	SerSerProSerGluSerProLysAspGluGlnUalProPheSerLysGluGluCysAla
1021	TCCAGCCCCTCGGAGTCCCCGAAGGACGAGGAGGAGGTCCCCTTCTCCAAGGAGGAATGTGCC
361	PhefingSerûlnLeuGluThrProGluThrLeuLeuGlySerThrûluGluLysProLeu
1081	TTTCGGTCRCRGCTG5AGAC6CCAGAGACCCTGCTGG5GGGGCACCGAAGAGAGCCCCTG
381 1141 1201 1261 1321 1381 1441 1501 1561 1681 1741 1801 1861	ProLeuGI yUa I ProAspA I aGI yMetLysProSer CCCCTTGGAGTGCCTGATGCTGGGATGAAGCCCAGTTAACCAGGCCGGTGTGGGGCTGTGT CGTAGCCAAGGTGGCTGAGGCCTGGCAGGATGACCCTGCGAAGGGGGCCCTGGTCCTTCCA GGCCCCCACCACTAGGACTCTGAGGCTCTTTCTGGGGCCAAGTTCCTCTAGTGCCCTCCAC AGCCGCAGCCTCCCTCTGACCTGCAGGCCAAGAGCAGAGGCAGCGAGTTGTGGAAAGCCT CTGCTGCCATGGCGTGTCCCTCTGGGAAGGCTGGCCGGGCAGGGGAGGTGGGGGGGG
1921 1981 2041 2101 2161 2221 2281	GCCGGGCGTGGTGGCGGGCACCTATRGTCCCAGCTACTCAGAAGCCTGAGGCTGGGAAAAAAAA CGTTTGAACCCGGGAAGCGGAGGTTGCAGGGGAGCCGAGATCACGCCACTGCACTCCAGCC TGGGCGACAGAGCGAGAGTCTGTCTCAAAAAAAAAA

HUMAN TNF RECEPTOR FUSION PROTEIN

This is a division of application Ser. No. 08/095,640, filed Jul. 21, 1993; now U.S. Pat. No. 5,610,279, which is a continuation application of Ser. No. 07/580,013, filed Sep. 10, 5 1990, now abandoned. This application claims priority under 35 U.S.C. §119 to application Serial Numbers 3319/89, 746/ 90 and 1347/90, filed on Sep. 12, 1989, Mar. 8, 1990 and Apr. 20, 1990, respectively, all in Switzerland. This application also claims priority under 35 U.S.C. §119 to European Patent 10 Application Number 90116707.2—(now Patent Number EP 0417563), filed Aug. 31, 1990.

BACKGROUND OF THE INVENTION

Tumor necrosis factor α (TNF α , also cachectin), discovered as a result of its hemorragic-necrotizing activity on certain tumors, and lymphotoxin (TNF β) are two closely related peptide factors [3] from the class of lymphokines/cytokines which are both referred to herein-after as TNF [see references 20 2 and 3]. TNF possesses a broad cellular spectrum of activity. For example, TNF has inhibitory or cytotoxic activity on a series of tumor cell lines [2,3], stimulates the proliferation of fibroblasts and the phagocytic/cytotoxic activity of myeloic cells [4, 5, 6], induces adhesion molecules in endothelial cells 25 or exerts an inhibitory activity on the endothelium [7, 8, 9, 10], inhibits the synthesis of specific enzymes in adipocytes [11] and induces the expression of histo-compatibility antigens [12]. Many of these TNF activities are produced via induction of other factors or by synergistic effects with other 30 factors such as interferons or interleukins [13-16].

TNF is involved in pathological conditions such as shock states in meningococcal sepsis [17], the development of autoimmune glomerulonephritis in mice [18] and cerebral malaria in mice [19] and human beings [41]. The toxic effects 35 of endotoxin appear to be mediated by TNF [20]. Furthermore, TNF can trigger interleukin-1 fever [39]. On the basis of its pleiotropic functional properties, TNF in interaction with other cytokines is involved in additional pathological conditions as a mediator of immune response, inflammation, 40 and other processes.

These biological effects are mediated by TNF via specific receptors. According to present knowledge not only $TNF\alpha$, but also TNF β bind to the same receptors [21]. Different cell types differ in their number of TNF receptors [22, 23, 24]. 45 Generally known TNF-binding proteins (TNF-BP) have been detected by covalent bonding to radioactively labelled TNF [24-29], and the following apparent molecular weights of the TNF/TNF-BP complexes obtained have been determined to be: $95/100\,kD$ and $75\,kD$ [24], $95\,kD$ and $75\,kD$ [25], $138\,kD,~50$ 90 kD, 75 kD and 54 kD [26], 100±5 kD [27], 97 kD and 70 kD [28] and 145 kD [29]. One such TNF/TNF-BP complex was isolated by anti-TNF-antibody immune affinity chromatography and preparative SDS-polyacrylamide gel electrophoreses (SDS-PAGE) [27]. The reductive cleavage of this 55 complex and subsequent SDS-PAGE analysis gave several bands which were not tested for TNF-binding activity. Since the specific conditions which must be used for the cleavage of the complex lead to inactivation of the binding protein [31], the latter has also not been possible. The separation of soluble 60 TNF-BP from human serum or urine by ion exchange chromatography and gel filtration (molecular weight in the region of 50 kD) was described by Olsson et al. [30]

Brockhaus et al. [32] obtained an enriched TNF-BP preparation from membrane extracts of HL_{60} cells by TNF α -ligand 65 affinity chromatography and HPLC which, in turn, was used as an antigen preparation for the production of monoclonal

antibodies against TNF-BP. Using such an immobilized antibody (immune affinity chromatography) Loetscher and Brockhaus obtained an enriched preparation of TNF-BP [31] from an extract of human placenta using TNF α -ligand affinity chromatography and HPLC, which gave a strong broad band at 35 kD, a weak band at about 40 kD and a very weak band in the region between 55 kD and 60 kD on SDS-PAGE analysis. Moreover, the gel showed a protein background smear in the region of 33 kD to 40 kD. The significance of these protein bands was, however, not clear due to the heterogenicity of the starting material which was used (placenta tissue; combined material from several placentas). In the state of the art TNF-BP have already been characterized by a N-terminal partial sequence [European Patent Application, Publication No. 308 378], whereby this sequence differs from the N-terminal partial sequence according to formula (IA) in accordance with the invention. Moreover, the TNF-binding proteins described in the state of the art are soluble, i.e. non-membrane bound, TNF-BP and not membrane-bound, i.e. insoluble, TNF-BP isolated from urine.

SUMMARY OF THE INVENTION

This invention comprises insoluble, homogenous proteins or soluble or insoluble fragments thereof, capable of binding tumor necrosis factor-(TNF).

This invention also comprises TNF-binding proteins containing amino acid sequences of FIG. 1 or FIG. 4, proteins containing fragments of these sequences, and proteins analagous to the sequences of FIG. 1 or FIG. 4 or to fragments thereof.

This invention % further comprises DNA sequences encoding the proteins: described above, proteins encoded by these sequences, and antibodies to any of these proteins.

This invention comprises. DNA sequences which combine two partial DNA sequences, one sequence encoding soluble fragments of TNF binding proteins and the other partial sequence encoding all domains except the first domain of the constant region of the heavy chain of human immunoglobulin IgG, IgA, IgM, or IgE, and the recombinant proteins encoded by these sequences.

This invention additionally comprises vectors containing the above DNA sequences, and host systems transfected with such vectors.

This invention finally comprises a process for the isolation of an insoluble homogenous protein capable of binding TNF.

BRIEF DESCRIPTION OF THE FIGURES

FIG. 1. Nucleotide sequence (SEQ ID NO: 1) and deduced amino acid sequence (SEQ ID NO: 2) for cDNA clone derived from 55 kD TNF-BP. The 19 amino acid transmembrane region is underlined. Hypothetical glycosylation sites are identified by asterisks.

FIG. 2. Binding analysis of COS cells transfected with plasmid pN123. Panel 2A—binding of transfected cells to ¹²⁵I-TNF α . Panel 2B—Scatchard plot of binding data.

FIG. **3**. Sandwich assays of cells transfected with plasmid pK19. Culture supernatants of cells transfected 30 with pK19 were incubated with anti-55 kD TNF-BP antibody followed by 125 I-TNF α . Columns 1, 5, and 8 are controls. Columns 2, 3, 4, 5, and 6 are five parallel transfections.

FIG. 4. Nucleotide sequence (SEQ ID NO: 3) and deduced amino acid sequence (SEQ ID NO: 4) for cDNA clones derived from 75/6510 TNF-BP.

DETAILED DESCRIPTION OF THE INVENTION

The TNF-binding proteins of the present invention are homogenous, insoluble proteins and soluble or insoluble

fragments of such proteins which are capable of binding TNF. These proteins have the ability to bind TNF as measured by standard assays.

The TNF-binding proteins of the present invention include homogenous proteins containing the amino acid sequence depicted in FIG. 1 (SEQ ID NO: 2) or in FIG. 4 (SEQ ID NO: 4), proteins containing fragments of either sequence, and analogues of any such proteins for example proteins containing amino acid sequences analogous to the amino acid sequences of FIG. 1 (SEQ ID NO: 2) or FIG. 4 (SEQ ID NO: 4) or to fragments thereof. An analogue is a protein in which one or more amino acids of the sequences depicted in FIG. 1 (SEQ ID NO: 2) or in FIG. 4 (SEQ ID NO: 4) have had their side-groups chemically modified in a known manner, or those 15in which one or more amino acids have been replaced or deleted, without thereby eliminating TNF-binding ability. Such analogues may be produced by known methods of peptide chemistry, or by known methods of recombinant DNA technology, such as planned mutagenesis. 20

The TNF binding activity of the proteins of the present invention may be determined using the assay described in Example 1.

TNF-binding proteins of this invention are obtained as follows:

TNF binding proteins may be isolated from tissues and purified to homogeneity, or isolated from cells which contain membrane-bound TNF binding protein, and purified to homogeneity. One possible method for growing cells and isolating cell extract is described in Example 2, however, 30 other cells types and other growth and isolation methods are well known in the art. Purification of TNF-binding proteins from cell extracts may be performed using the methods described in Examples 4, 5, and 6 in combination with the assay described in Example 1. TNF-binding proteins isolated 35 with the invention is also an object of the present invention. and purified by these methods were sequenced by wellknown methods, as described in Example 7. From these amino acid sequences, DNA probes were produced and used to obtain mRNA encoding TNF binding proteins from which cDNA was made, all by known methods described in 40 Examples 8 and 11. Other well-known methods for producing cDNA are known in the art and may effectively be used. In general, any TNF-binding protein can be isolated from any cell or tissue expressing such proteins using a cDNA probe such as the probe described above, isolating mRNA and tran- 45 scribing the mRNA into cDNA. Thereafter, the protein can be produced by inserting the cDNA into an expression vector as described in Example 9, such as a virus, plasmid, cosmid, or other vector, inserting the expression vector into a cell, such as the COS cell-described in Example 9 or the insect cell 50 described in Example 10, proliferating the resulting cells, and isolating the expressed TNF-binding protein from the medium or from cell extract as described above. Alternatively, TNF-binding proteins may be chemically synthesized using the sequence described above and an amino acid syn- 55 thesizer, or manual synthesis using chemical conditions well known to form peptide bonds between selected amino acids. Analogues and fragments of TNF-binding proteins may be produced by the above methods. In the case of analogues, the proteins may be chemically modified, or modified by genetic 60 engineering as described above. These fragments and analogues may then be tested for TNF-binding activity using methods such as the assay of Example 1.

Finally, monoclonal antibodies directed against TNF-binding proteins, such as the antibodies described in Example 3, 65 may be produced by known techniques, and used to isolate TNF-binding proteins.

4

In more detail, the proteins of the present invention are non-soluble proteins, i.e. for example membrane proteins or so-called receptors, and soluble or non-soluble, fragments thereof, which bind TNF (TNF-BP), in homogeneous form, as well as their physiologically compatible salts. Preferred proteins are those which according to SDS-PAGE under nonreducing conditions are characterized by apparent molecular weights of about 55 kD, 51 kD, 38 kD, 36 kD and 34 kD or 75 kD and 65 kD, especially those with about 55 kD and 75 kD. Furthermore, there are preferred those proteins which are characterized by containing at least one of the following amino acid partial sequences:

- (IA) Leu-Val-Pro-His-Leu-Gly-Asp-Arg-Glu-Lys-Arg-Asp-Ser-Val-Cys-Pro-Gln-Gly-Lys-Tyr-Ile-His-Pro-Gln-X-Asn-Ser-Ile (SEQ ID NO: 5)
- (IB) Ser-Thr-Pro-Glu-Lys-Glu-Gly-Glu-Leu-Glu-Gly-Thr-Thr-Thr-Lys (SEQ ID NO: 6)
- (IIA) Ser-Gln-Leu-Glu-Thr-Pro-Glu-Thr-Leu-Leu-Gly-Ser-Thr-Glu-Glu-Lys-Pro-Leu (SEQ ID NO: 7)
- (IIB) Val-Phe-Cys-Thr (SEQ ID NO: 8)
- (IIC) Asn-Gln-Pro-Gln-Ala-Pro-Gly-Val-Glu-Ala-Ser-Gly-Ala-Gly-Glu-Ala (SEQ ID NO: 9)
- (IID) Leu-pro-Ala-Gln-Val-Ala-Phe-X-Pro-Tyr-Ala-Pro-Glu-Pro-Gly-Ser-Thr-Cys (SEQ ID NO: 10)
- (IIE) Ile-X-Pro-Gly-Phe-Gly-Val-Ala-Tyr-Pro-Ala-Leu-Glu (SEQ ID NO: 11)
- (IIF) Leu-Cys-Ala-Pro (SEQ ID NO: 12)
- (IIG) Val-Pro-His-Leu-Pro-Ala-Asp (SEQ ID NO: 13)
- (IIH) Gly-Ser-Gln-Gly-Pro-Glu-Gln-Gln-X-X-Leu-Ile-X-Ala-Pro (SEQ ID NO: 14)
 - in which X stands for an amino acid residue which could not be unequivocally determined.

A process for the isolation of the TNF-BP in accordance This process comprises carrying out essentially the following purification steps in sequence: production of a cell or tissue extract, immune affinity chromatography and/or single or multiple ligand affinity chromatography, high resolution liquid chromatography (HPLC) and preparative SDS-polyacrylamide gel electro phoresis (SDS-PAGE). The combination of the individual purification steps, which are known from the state of the art, is essential to the success of the process in accordance with the invention, whereby individual steps have been modified and improved having regard to the problem to be solved. Thus, for example, the original combined immune affinity chromatography/TNF\alpha-ligand affinity chromatography step originally used for the enrichment of TNF-BP from human placenta [31] has been altered by using a BSA-Sepharose 4B pre-column. For the application of the cell or membrane extract, this pre-column was connected in series with the immune affinity column followed by the ligand affinity column. After the application of the extract the two aforementioned columns were coupled, each eluted and the TNF-BP-active fractions were purified again via a ligand affinity column. The use of a detergent-containing solvent mixture for the performance of the reversed-phase HPLC step is essential to the invention.

Further, an industrial process for the production of high cell densities of mammalian cells from which TNF-BP can be isolated is also an object of the present invention. Such a process comprises using a medium, which has been developed for the specific growth requirements of the cell line used, in combination with a perfusion apparatus as described e.g. in detail in Example 2. By means of such a process there can be produced, for example, in the case of HL-60 cells up to more than 20-fold higher cell densities than usual.

In addition thereto, the present invention is also concerned with DNA sequences coding for proteins and soluble or nonsoluble fragments thereof, which bind TNF. Thereunder there are to be understood; for example, DNA sequences coding for non-soluble proteins or soluble as well as non-soluble fragments thereof, which bind TNF, such DNA sequences being selected from the following:

(a) DNA sequences as given FIG. 1 or FIG. 4 as well as their complementary strands, or those which include these sequences;

(b) DNA sequences which hybridize with sequences defined under (a) or fragments thereof;

(c) DNA sequences which, because of the degeneracy of the genetic code, do not hybridize with sequences as defined under (a) and (b), but which code for polypeptides, having 15 exactly the same amino acid sequence.

That is to say, the present invention embraces not only allelic variants, but also those DNA sequences which result from deletions, substitutions and additions from one or more nucleotides of the sequences given in FIG. 1 or FIG. 4, 20 whereby in the case of the proteins coded thereby there come into consideration, just as before, TNF-BP. One sequence which results from such a deletion is described, for example, in Science 248, 1019-1023, (1990).

There are preferred first of all those DNA sequences which 25 code for such a protein having an apparent molecular weight of about 55 kD, whereby the sequence given in FIG. 1 is especially preferred, and sequences which code for nonsoluble as well as soluble fragments of such proteins. A DNA sequence which codes, for example, for such a non-soluble 30 protein fragment extends from nucleotide -185 to 1122 of the sequence given in FIG. 1. DNA sequences which code for soluble protein fragments are, for example, those which extend from nucleotide -185 to 633 or from nucleotide -14 to 633 of the sequence given in FIG. 1. There are also preferred 35 DNA sequences which code for a protein of about 75/65 kD, whereby those which contain the partial cDNA sequences shown in FIG. 4 are preferred. Especially preferred DNA sequences in this case are the sequences of the open reading frame of nucleotide 2 to 1,177. The peptides IIA, IIC, IIE, IIF, 40 IIG and IIH are coded by the partial cDNA sequence in FIG. 4, whereby the insignificant deviations in the experimentally determined amino acid sequences are based on the cDNAderived sequence with highest probability from the limited resolution of the gas phase sequencing. DNA sequences 45 which code for insoluble (deposited on Oct. 17, 2006 with the American Type Culture Collection under Accession No. PTA 7942) as well as soluble fractions of TNF-binding proteins having an apparent molecular weight of 65 kD/75 kD are also preferred. DNA sequences for such soluble fragments can be 50 determined on the basis of the amino acid sequences derived from the nucleic acid sequences coding for such non-soluble TNF-BP.

The invention is also concerned with DNA sequences which comprise a combination of two partial DNA 55 sequences, with one of the partial sequences coding for those soluble fragments of non-soluble proteins which bind TNF (see above) and the other partial sequence coding for all domains other than the first domain of the constant region of the heavy chain of human immunoglobulins such as IgG, IgA, 60 IgM or IgE, in particular IgG₁ or IgG₃ subtypes.

The present invention is also concerned with the recombinant proteins coded by any of DNA sequences described above. Of course, there are thereby also included such proteins in whose amino acid sequences amino acids have been 65 exchanged, for example by planned mutagenesis, so that the activity of the TNF-BP or fragments thereof, namely the 6

binding of TNF or the interaction with other membrane components participating in the signal transfer, have been altered or maintained in a desirable manner. Amino acid exchanges in proteins and peptides which do not generally alter the activity of such molecules are known in the state of the art and are described, for example, by H. Neurath and R. L. Hill in "The'Proteins" (Academic Press, New York, 1979, see especially FIG. 6, page 14). The most commonly occurring exchanges are: Ala/Ser, Val/Ile, Asp/Glu, Thr/Ser, Ala/Gly, Ala/Thr, Ser/Asn, Ala/Val, Ser/Gly, Tyr/Phe, Ala/Pro, Lys/ Arg, Asp/Asn, Leu/Ile, Leu/Val, Ala/Glu, Asp/Gly as well as these in reverse. The present invention is also concerned with vectors which contain any of the DNA sequences described above in accordance with the invention and which are suitable for the transformation of suitable pro- and eukaryotic host systems, whereby there are preferred those vectors whose use leads to the expression of the proteins which are coded by any of the DNA sequences described above in accordance with the invention. Finally, the present invention is also concerned with pro- and eukarvotic host systems transformed with such vectors, as well as a process for the production of recombinant compounds in accordance with the invention by cultivating such host systems and subsequently isolating these compounds from the host systems themselves or their culture supernatants.

An object of the present invention are also pharmaceutical preparations which contain at least one of these TNF-BPs or fragments thereof, if desired in combination with other pharmaceutically active substances and/or non-toxic, inert, therapeutically compatible carrier materials.

Finally, the present invention is concerned with the use of such a TNF-BP on the one hand for the production of pharmaceutical preparations and on the other hand for the treatment of illnesses, preferably those in which TNF is involved in their course.

Starting materials for the TNF-BP in accordance with the invention are quite generally cells which contain such TNF-BP [in membrane-bound form] and which are generally accessible without restrictions to a person skilled in the art, such as, for example, HL60 [ATCC No. CCL 240], U 937 [ATCC No. CRL 1593], SW 480 [ATCC No. CCL 228] and HEp2 cells [ATCC No. CCL 23]. These cells can be cultivated according to known methods of the state of the art [40] or, in order to produce high cell densities, according to the procedure already described generally and described in detail in. Example 2 for HL60 cells. TNF-BP can then be extracted from the cells, which are centrifuged-off from the medium and washed, according to known methods of the state of the art using suitable detergents, for example Triton X-114, 1-0n-octyl-β-D-glucopyranoside (octylglucoside) or 3-[(3cholylamido-propyl)-dimethylammonio]-1-propane sulphonate (CHAPS), especially using Triton X-100. For the detection of such TNF-BP there can be Used the usually used detection methods for TNF-BP, for example a polyethylene glycol-1-induced precipitation of the ¹²⁵I-TNF/TNF-BP complex [27], especially filter-binding tests with radioactively labelled TNF according to Example 1. In order to produce the TNF-BP in accordance with the invention, the general methods of the state of the art used for the purification of proteins, especially of membrane proteins, such as, for example, ion exchange chromatography, gel filtration, affinity chromatography, HPLC and SDS-PAGE can be used. Especially preferred methods for the production of TNF-BP in accordance with the invention are affinity chromatography, especially with TNF-a as the ligand bound to the solid phase, and immune-affinity chromatography, HPLC and SDS-PAGE. The elution of TNF-BP bands which are separated

using SDS-PAGE can be effected according to known methods of protein chemistry, for example using electroelution according to Hunkapiller et al. [34], whereby according to present knowledge the electro-dialysis times given there generally have to be doubled. Thereafter, traces of SDS which 5 still remain can then be removed in accordance with Bosserhoff et al. [50]

The thus-purified TNF-BP can be characterized by methods of peptide chemistry which are known in the state of the art, such as, for example, N-terminal amino acid sequencing of enzymatic well as chemical peptide cleavage. Fragments obtained by enzymatic or chemical cleavage can be separated according to usual methods such as, for example, HPLC and can themselves be subjected to further N-terminal sequencing. Such fragments which themselves bind TNF can be iden-15 tified using the afore-mentioned detection methods for TNF-BP and are likewise objects of the present invention.

Starting from the thus-obtained amino acid sequence information or the DNA and amino acid sequences given in FIG. 1 as well as in FIG. 4 there can be produced, taking into con- 20 sideration the degeneracy of the genetic code, according to methods known in the state of the art suitable oligonucleotides [51]. By means of these, again according to known methods of molecular biology [42,43], cDNA or genomic DNA banks can be searched for clones which contain nucleic 25 acid sequences coding for TNF-BP. More-over, using the polymerase chain reaction (PCR) [49] cDNA fragments can be cloned by completely degenerating the amino acid sequence of two spaced apart relatively short segments while taking into consideration the genetic code and introducing into their complementarity suitable oligo-nucleotides as a "primer", whereby the fragment lying between these two sequences can be amplified and identified. The determination of the nucleotide sequence of a such a fragment permits an independent determination of the amino acid sequence of the 35 protein fragment for which it codes. The cDNA fragments obtainable by PCR can also, as already described for the oligonucleotides themselves, be used according to known methods to search for clones containing nucleic acid sequences coding for TNF-BP from cDNA or genomic DNA 40 e.g. cells of the human cell lines Hela [ATCC CCL2] and 293 banks. Such nucleic acid sequences can then be sequenced according to known methods [42]. On the basis of the thusdetermined sequences and of the already known sequences for certain receptors, those partial sequences which code for soluble TNF-BP fragments can be determined and cut out 45 from the complete sequence using known methods [42].

The complete sequence or such partial sequences can then be integrated using known methods into vectors described in the state of the art for their multiplication and expression in prokaryotes [42]. Suitable prokaryotic host organisms are, for 50 example, gram-negative and gram-positive bacteria such as, for example, E. coli strains such as E. coli HB101 [ATCC No. 33 694] or E. coli W3110 [ATCC No. 27 325] or B. subtilis strains.

Furthermore, nucleic acid sequences in accordance with 55 the invention which code for TNF-BP as well as for TNF-BP fragments can be integrated using known methods into suitable vectors for reproduction and expression in eukaryotic host cells, such as, for example, yeast, insect cells and mammalian cells. Expression of such sequences is preferably 60 effected in mammalian and insect cells.

A typical expression vector for mammalian cells contains an efficient promoter element in order to produce a good transcription rate, the DNA sequence to be expressed and signals for an efficient termination and polyadenylation of the 65 transcript. Additional elements which can be used are "enhancers" which lead to again intensified transcription and

sequences which e.g. can bring about a longer half life of the mRNA. For the expression of nucleic acid sequences in which the endogenous sequence fragment coding for a signal peptide is missing, there can be used vectors which contain such suitable sequences which code for signal peptides of other known proteins. See, for example, the vector pLJ268 described by Cullen, B. R. in Cell 46, 973-982 (1986) as well as Sharma, S. et al. in "Current Communications in Molecular Biology", edt. by Gething, M. J., Cold Spring Harbor Lab. (1985), pages 73-78.

Most of these vectors which are used for a transient expression of a particular DNA sequence in mammalian cells contain the replication source of the SV40 virus. In cells which express the T-antigen of the virus (e.g. COS cells), these vectors are reproduced abundantly. A transient expression is, however, not limited to COS cells. In principle any transfectable mammalian cell line can be used for this purpose. Signals which can bring about a strong transcription are e.g. the early and late promoters of SV40, the promoter and enhancer of the "major immediate-early" gene of HCMV (human cytomegalovirus), the LTR's ("long terminal repeats") of retroviruses such as, for example, RSV, HIV and MMTV. There can, however, also be used signals of cellular genes such as e.g. the promoters of the actin and collagenase genes.

Alternatively, however, stable cell lines which have the specific DNA sequence integrated into the genome (chromosome) can also be obtained. For this, the DNA sequence is cotransfected together with a selectable marker, e.g. neomycin, hygromycin, dihydrofolate reductase (dhfr) or hypoxanthin guanine phosphoribosyl transferase (hgpt). The DNA sequence stably incorporated in the chromosome can also be reproduced abundantly. A suitable selection marker for this is, for example, dihydrofolate reductase (dhfr). Mammalian cells (e.g. CHO cells), which contain no intact dhfr gene, are thereby incubated with increasing amounts of methotrexate after transinfection has been effected. In this manner cell lines which contain more than a thousand copies of the desired DNA sequence can be obtained.

Mammalian cells which can be used for the expression are [ATCC CRL 1573] as well as 3T3 [ATCC CCL 163] and L cells, e.g. [ATCC CCL 149], (CHO) cells [ATCC CCL 61], BHK [ATCC CCL 10] cells as well as the CV 1 [ATCC CCL 70] and the COS cell lines [ATCC CRL 1650, CRL 1651].

Suitable expression vectors include, for example, vectors such as pBC12MI [ATCC 67 109], pSV2dhfr [ATCC 37 146], pSVL [Pharmacia, Uppsala, Sweden], pRSVcat [ATCC 37 152] and pMSG [Pharmacia, Uppsala, Sweden]. The vectors "pK19" and "pN123" used in Example 9 are especially preferred vectors. These can be isolated according to known methods from E. coli strains HB101(pK19) and HB101 (pN123) transformed with them [42]. These E. coli strains have been deposited on the 26 Jan. 1990 at the Deutschen Sammlung von Mikroorganismen and Zellkulturen GmbH (DSM) in Braunschweig, FRG, under DSM 5761 for HB101 (pK19) and DMS 5764 for HB101(pN123). For the expression of proteins which consist of a soluble fragment of nonsoluble TNF-BP and an immunoglobulin fragment, i.e. all domains except the first of the constant region of the heavy chain, there are especially suitable pSV2-derived vectors as described, for example, by German, C. in "DNA Cloning" [Vol. II., edt. by Glover, D. M., IRL Press, Oxford, 1985]. The vectors pCD4-Hp (DSM 5315, deposited on 21 Apr. 1989), pDC4-Hy1 (DSM 5314, deposited on 21 Apr. 1989) and pCD4-Hy3 (DSM 5523, deposited on 14 Sep. 1989) which have been deposited at the Deutschen Sammlung von Mikroorganismen and Zellkulturen GmbH (DSM) in Braunsch-

55

weig, FRG, and which are described in detail in European Patent Application No. 90107393.2 are especially preferred vectors. This European Patent Specification and the equivalent Applications referred to in Example 11 also contain data with respect to the further use of these vectors for the expres- 5 sion of chimeric proteins (see also Example 11) and for the construction of vectors for the expression of such chimeric proteins with other immunoglobulin fragments.

The manner in which these cells are transfected depends on the chosen expression system and vector system. An over-10 view of these methods is to be found e.g. in Pollard et al., "DNA Transformation of Mammalian Cells" in "Methods in Molecular Biology" [Nucleic Acids Vol. 2, 1984, Walker, J. M., ed, Humana, Clifton, N.J.]. Further methods are to be found in Chen and Okayama ["High-Efficiency Transforma- 15 tion of Mammalian Cells by Plasmid DNA", Molecular and Cell Biology 7, 2745-2752, 1987] and in Feigner [Feigner et al., "Lipofectin: A highly efficient, lipid-mediated. DNAtransfection procedure", Proc. Nat. Acad. Sci. USA 84, 7413-7417, 1987].

The baculovirus expression system, which has already been used successfully for the expression of a series of proteins (for an overview see Luckow and Summers, Bio/Technology 6, 47-55, 1988), can be used for the expression in insect cells. Recombinant proteins can be produced in authen-25 tic form or as fusion proteins. The thus-produced proteins can also be modified such as, for example, glycosylated (Smith et al., Proc. Nat. Acad. Sci. USA 82, 8404-8408, 1987). For the production of a recombinant baculovirus which expresses the desired protein there is used a so-called "transfer vector". 30 Under this there is to be understood a plasmid which contains the heterologous DNA sequence under the control of a strong promoter, e.g. that of the polyhedron gene, whereby this is surrounded on both sides by viral sequences. The vectors "pN113", "pN119" and "pN124" used in Example 10 are 35 especially preferred vectors. These can be isolated according to known methods from E. coli strains HB101(pN113), HB101(pN119) and HB101(pN124) transformed with them. These E. coli strains have been deposited on the 26 Jan. 1990 at the Deutschen Sammlung von Mikroorganismen and 40 Zellkulturen GmbH (DSM) in Braunschweig, FRG, under DSM 5762 for HB101(pN113), DSM 5763 for HB101 (pN119) and DSM 5765 for HB101(pN124). The transfer vector is then transfected into the insect cells together with DNA of the wild type baculovirus. The recombinant viruses 45 which result in the cells by homologous recombination can then be identified and isolated according to known methods. An overview of the baculovirus expression system and the methods used therein is to be found in Luckow and Summers [52].

Expressed TNF-BP as well as its non-soluble or soluble fractions can then be purified from the cell mass or the culture supernatants according to methods of protein chemistry which are known in the state of the art, such as, for example, the procedure already described on pages 5-6.

The TNF-BP obtained in accordance with the invention can also be used as antigens to produce polyclonal and monoclonal antibodies according to known-techniques [44, 45] or according to the procedure described in Example 3. Such antibodies, especially monoclonal antibodies against the 75 60 kD TNF-BP species, are also an object of the present invention. Those antibodies which are directed against the 75 kD TNF-BP can be used for the isolation of TNF-BP by modifications of the purification procedure described in detail in Examples 4-6 which are familiar to a person skilled in the art. 65

On the basis of the high binding affinity of TNF-BP in accordance with the invention for TNF (K_d value in the order

of 10^{-9} - 10^{-10} M), these or fragments thereof can be used as diagnostics for the detection of TNF in serum or other body fluids according to methods known in the state of the art, for example in solid phase binding tests or in combination with anti-TNF-BP antibodies in so-called "sandwich" tests.

Moreover, TNF-BP in accordance with the invention can be used on the one hand for the purification of TNF and on the other hand for the detection of TNF agonists and TNF antagonists according to procedures which are known in the state of the art

The TNF-BP in accordance with the invention as well as their physiologically compatible salts, which can be manufactured according to methods which are known in the state of the art, can also be used for the production of pharmaceutical preparations, primarily those for the treatment of illnesses in which TNF is involved in their course. For this purpose, one or more of the said compounds, where desired or required in combination with other pharmaceutically active substances, can be processed in a known manner with the usually used solid or liquid carrier. materials. The dosage of such preparations can be effected having regard to the usual criteria in analogy to already used preparations of similar activity and structure.

Since the invention has been described hereinbefore in general terms, the following Examples are intended to illustrate details of the invention, but they are not intended to limit its scope in any manner.

Example 1

Detection of TNF-Binding Proteins

The TNF-BP were detected in a filter test with human radioiodinated ¹²⁵I-TNF. TNF (46, 47) was radioactively labelled with Na¹²⁵I (IMS40, Amersham, Amersham, England) and iodo gene (#28600, Pierce Eurochemie, Oud-Beijerland, Netherlands) according to Fraker and Speck [48]. For the detection of the TNF-BP, isolated membranes of the cells or their solubilized, enriched and purified fractions were applied to moist nitrocellulose filter (0.45µ, BioRad, Richmond, Calif., USA). The filters were then blocked in buffer solution with 1% skimmed milk powder and subsequently incubated with $5 \cdot 10^5$ cpm/ml of ¹²⁵I-TNFa (0.3-1.0·10⁸ cpm/ μ g) in two batches with and without the addition of 5 μ g/ml of non-labelled TNF α , washed and dried in the air. The bound radio-activity was detected semiquantitatively by autoradiography or counted in a gamma-counter. The specific ¹²⁵I-TNF- α binding was determined after correction for unspecific binding in the presence of unlabelled TNF- α in excess. The specific TNF-binding in the filter test was measured at various TNF concentrations and analyzed according to Scatchard, whereby a K_d value of $\cdot 10^{-9} \cdot 10^{-10}$ M was determined.

Example 2

Cell Extracts of HL-60 Cells

HL60 cells [ATCC No. CCL 240] were cultivated on an experimental laboratory scale in a RPMI 1640 medium [GIBCO catalogue No. 074-01800], which contained 2 g/l NaHCO₃ and 5% foetal calf serum, in a 5% CO₂ atmosphere, and subsequently centrifuged.

The following procedure was used to produce high cell densities on an industrial scale. The cultivation was carried out in a 751 Airlift fermenter (Fa. Chemap, Switzerland) with a working volume of 581. For this there was used the cassette membrane system "PROSTAK" (Millipore, Switzerland)

40

55

with a membrane surface of $0.32 \text{ m}^2(1 \text{ cassette})$ integrated into the external circulation circuit. The culture medium (see Table 1) was pumped around with a Watson-Marlow pump, Type 603U, with 5 l/min. After a steam sterilization of the installation, whereby the "PROSTAK" system was sterilized 5 separately in autoclaves, the fermentation was started with growing HL-60 cells from a 201 Airlift fermenter (Chemap). The cell cultivation in the inoculation fermenter was effected in a conventional batch process in the medium according to Table 1 and an initial cell titre of 2×10^5 cells/ml. After 4 days the HL60 batch was transferred with a titre of 4.9×10⁶ cells/ ml into the 751 fermenter. The pH value was held at 7.1 and the pO₂ value was held at 25% saturation, whereby the oxygen introduction was effected through a microporous frit. 15 After initial batch fermentation, on the 2nd day the perfusion at a cell titre of 4×10^6 cells/ml was started with 301 of medium exchange per day. On the filtrate side of the medium the conditioned medium was removed and replaced by the addition of fresh medium. The added medium was fortified as 20 follows: Primatone from 0.25% to 0.35%, glutamine from 5 mM to 6 mM and glucose from 4 g/l to 6 g/l. The perfusion rate was then increased on the 3rd and 4th day to 72 l of medium/day and on the 5th day to 100 l of medium/day. The fermentation had finished after 120 hours of continuous cul-25 tivation. Exponential cell growth up to 40×10^6 cells/ml took place under the given fermentation conditions. The duplication time of the cell population was 20-22 hours to 10×10^6 cells/ml and then increased to 30-36 hours with increasing cell density. The proportion of living cells lay at 90-95% during the entire fermentation period. The HL-60 batch was then cooled down in the fermenter to about 12° C. and the cells were harvested by centrifugation (Beckman centrifuge [Model J-6B, Rotor JS], 3000 rpm, 10 min., 4° C.).

TABLE 1

HL-60 medium		
Components	Concentrations mg/l	
CaCl ₂ (anhydrous)	112.644	
$Ca(NO_3)_2 \bullet 4H_2O$	20	
$CuSO_4 \bullet 5H_2O$	$0.498 \cdot 10^{-3}$	
$Fe(NO_3)_3 \bullet 9H_2O$	0.02	
$FeSO_4 \bullet 7H_2O$	0.1668	
KCl	336.72	
KNO3	0.0309	
MgCl ₂ (anhydrous)	11.444	
MgSO ₄ (anhydrous)	68.37	
NaCl	5801.8	
Na ₂ HPO ₄ (anhydrous)	188.408	
$NaH_2PO_4 \bullet H_2O$	75	
$Na_2SeO_3 \bullet 5H_2O$	$9.6 \cdot 10^{-3}$	
$ZnSO_4 \bullet 7H_2O$	0.1726	
D-Glucose	4000	
Glutathion (red.)	0.2	
Hepes buffer	2383.2	
Hypoxanthin	0.954	
Linoleic acid	0.0168	
Lipoic acid	0.042	
Phenol Red	10.24	
Putrescine 2HCl	0.0322	
Na pyruvate	88	
Thymidine	0.146	
Biotin	0.04666	
D-Ca pantothenate	2.546	
Choline chloride	5.792	
Folic acid	2.86	
i-Inositol	11.32	
Niacinamide	2.6	
Nicotinamide	0.0074	
para-Aminobenzoic acid	0.2	

	12
TABLE	1-continued

$\begin{tabular}{ c c c c c } \hline Concentrations mg/l \\ \hline Pyridoxal HCl & 2.4124 \\ Pyridoxin HCl & 0.2 \\ Riboflavin & 0.2876 \\ Thiamin HCl & 2.668 \\ Vitamin B_{12} & 0.2782 \\ L-Alanine & 11.78 \\ L-Asparagine H_2O & 14.362 \\ L-Arginine HCl & 92.6 \\ L-Arginine HCl & 92.6 \\ L-Arginine HCl & 92.6 \\ L-Arginine HCl & 62.04 \\ L-Cystine 2HCl & 62.04 \\ L-Cysteine HCl • H_2O & 7.024 \\ L-Glutamic acid & 36.94 \\ L-Glutamic acid & 36.94 \\ L-Glutamine & 730 \\ L-Hydroxypyroline & 4 \\ L-Isoleucine & 73.788 \\ L-Leucine & 75.62 \\ L-Lysine HCl & 100.29 \\ L-Methionine & 21.896 \\ L-Tyrosine + 2Na & 69.76 \\ L-Valine & 62.74 \\ Penicillin/streptomycin & 100 U/ml \\ Insulin (human) & 5 µg/ml \\ Bovine serum albumin & 5 µg/ml \\ Bovine serum albumin & 5 µg/ml \\ Phuronic F68 & 0.01\% \\ (Serva, Heidelberg, FRG) \\ Foetal calf serum & 0.3-3\% \\ \end{bmatrix}$	HL-60 me	tium
Pyridoxal HCl 2.4124 Pyridoxin HCl 0.2 Riboflavin 0.2876 Thiamin HCl 2.668 Vitamin B_{12} 0.2782 L-Alanine 11.78 L-Asparagine H2O 14.362 L-Arginine 40 L-Asparagine H2O 14.362 L-Arginine 40 L-Asparate 33.32 L-Cystine 2HCl 62.04 L-Cysteine HCl • H2O 7.024 L-Glutamic acid 36.94 L-Glutamine 730 L-Glutamine 730 L-Glutamine 73.788 L-Leucine 75.62 L-Lysine HCl 102.9 L-Hydroxypyroline 4 L-Isoleucine 75.62 L-Lysine HCl 102.9 L-Methionine 21.896 L-Proline 26.9 L-Serine 31.3 L-Threonine 53 L-Tyrosine • 2Na 69.76 L-Valine 62.74 Penicillin/streptomycin 100 U/m1 Insulin (human) 5 µg/	Components	Concentrations mg/l
Pyridoxin HCl 0.2 Riboflavin 0.2876 Thiamin HCl 2.668 Vitamin B_{12} 0.2782 L-Alanine 11.78 L-Asparagine H2O 14.362 L-Arginine 40 L-Arginine HCl 92.6 L-Asparate 33.32 L-Cystine 2HCl 62.04 L-Cysteine HCl • H2O 7.024 L-Glutamic acid 36.94 L-Glutamine 730 L-Glutamine 730 L-Glutamine 730 L-Glutamine 73.788 L-Leucine 75.62 L-Lysine HCl 102.9 L-Hydroxypyroline 4 L-Isoleucine 73.788 L-Leucine 75.62 L-Lysine HCl 102.9 L-Methionine 21.896 L-Proline 26.9 L-Serine 31.3 L-Threonine 53 L-Tryptophan 11.008 L-Typtonie • 2Na 69.76 L-Valine 62.74 Penicillin/streptomycin 100 <td>Pyridoxal HCl</td> <td>2.4124</td>	Pyridoxal HCl	2.4124
Riboflavin 0.2876 Thiamin HCl 2.668 Vitamin B ₁₂ 0.2782 L-Alanine 11.78 L-Aspartic acid 10 L-Arginine 40 L-Arginine HCl 92.6 L-Aspartate 33.32 L-Cysteine HCl • H ₂ O 7.024 L-Glutamic acid 36.94 L-Glycine 21.5 L-Histidine 3 L-Histidine 3 L-Soleucine 73.788 L-Leucine 75.62 L-Lysine HCl 102.9 L-Methionine 21.896 L-Phonylalanine 43.592 L-Proline 26.9 L-Serine 31.3 L-Threonine 53 L-Tryptophan 11.008 L-Tyrosine • 2Na 69.76 L-Valine 62.74 Penicillin/streptomycin 100 U/ml Insulin (human) 5 µg/ml	Pyridoxin HCl	0.2
Thiamin HCl 2.668 Vitamin B_{12} 0.2782 L-Alanine 11.78 L-Aspartic acid 10 L-Asparagine H ₂ O 14.362 L-Arginine 40 L-Arginine HCl 92.6 L-Asparate 33.32 L-Cystine 2HCl 62.04 L-Cysteine HCl • H ₂ O 7.024 L-Glutamic acid 36.94 L-Glycine 21.5 L-Histidine 3 L-Histidine 3 L-Hydroxypyroline 4 L-Isoleucine 73.62 L-Lysine HCl 102.9 L-Hydroxypyroline 4 L-Isoleucine 73.788 L-Leucine 75.62 L-Lysine HCl 102.9 L-Methionine 21.896 L-Phonylalanine 43.892 L-Phonylan 11.008 L-Tyrosine • 2Na 69.76 L-Valine 62.74 Penicillin/streptomycin 100 U/ml Insulin (human) 5 µg/ml Tranferrin (human) 5 µg/ml Tranferr	Riboflavin	0.2876
Vitamin B_{12} 0.2782 L-Alanine 11.78 L-Asparatic acid 10 L-Asparatic method 14.362 L-Arginine H ₂ O 14.362 L-Arginine HCI 92.6 L-Asparate 33.32 L-Cystine 2HCI 62.04 L-Cysteine HCI • H ₂ O 7.024 L-Glutamic acid 36.94 L-Glycine 21.5 L-Histidine 3 L-Histidine HCI • H ₂ O 27.392 L-Hydroxypyroline 4 L-Isoleucine 75.62 L-Lysine HCI 102.9 L-Methionine 21.896 L-Phenylalanine 43.592 L-Phenylalanine 26.9 L-Serine 31.3 L-Threonine 53 L-Tyrosine • 2Na 69.76 L-Tyrosine • 2Na 69.76 L-Valine 62.74 Penicillin/streptomycin 100 U/ml Insulin (human) 5 µg/ml Tranferrin (human) 5 µg/ml Primatone RL (Sheffield Products, 0.25% 0.25% Norwich NY, USA)	Thiamin HCl	2.668
L-Alanine 11.78 L-Aspartic acid 10 L-Asparagine H ₂ O 14.362 L-Arginine 40 L-Arginine HCl 92.6 L-Aspartate 33.32 L-Cystine 2HCl 62.04 L-Glutamic acid 36.94 L-Glutamic acid 36.94 L-Glutamic acid 3 L-Glycine 21.5 L-Histidine 3 L-Histidine HCl • H ₂ O 27.392 L-Hydroxpyproline 4 L-Isoleucine 73.788 L-Leucine 75.62 L-Phenylalanine 43.592 L-Phenylalanine 43.592 L-Proline 26.9 L-Serine 31.3 L-Threonine 53 L-Tryptophan 11.008 L-Tyrosine • 2Na 69.76 L-Valine 62.74 Penicillin/streptomycin 100 Insulin (human) 5 µg/ml Tranferrin (human) 5 µg/ml Primatone RL (Sheffield Products, 0.25% 0.25% Norwich NY, USA) 0.01%	Vitamin B ₁₂	0.2782
L-Aspartic acid 10 L-Asparagine H ₂ O 14.362 L-Arginine HCI 92.6 L-Aspartate 33.32 L-Cystine 2HCI 62.04 L-Cysteine HCI \bullet H ₂ O 7.024 L-Glutamic acid 36.94 L-Glutamine 730 L-Glutamine 730 L-Glutamine 7 L-Glutamine 7 L-Histidine 3 L-Histidine 3 L-Leucine 75.62 L-Lysine HCI 102.9 L-Hydroxypyroline 4 L-Isoleucine 73.788 L-Leucine 75.62 L-Lysine HCI 102.9 L-Methionine 21.896 L-Proline 26.9 L-Serine 31.3 L-Threonine 53 L-Tryptophan 11.008 L-Typtophan 11.008 L-Valine 62.74 Penicillin/streptomycin 100 U/ml Insulin (human) 5 µg/ml Tranferrin (human) 5 µg/ml Tranferrin (human) 67	L-Alanine	11.78
L-Asparagine H_2O 14.362 L-Arginine 40 L-Arginine HCI 92.6 L-Asparata 33.32 L-Cystine 2HCI 62.04 L-Cysteine HCI • H_2O 7.024 L-Glutamic acid 36.94 L-Glycine 21.5 L-Histidine 3 L-Histidine 3 L-Histidine 3 L-Lysine HCI • H_2O 27.392 L-Histidine 3 L-Histidine 3 L-Histidine 3 L-Lysine HCI 102.9 L-Lysine HCI 102.9 L-Methionine 26.9 L-Serine 31.3 L-Threonine 53 L-Trytosine • 2Na 69.76 L-Valine 62.74 Penicillin/streptomycin 100 U/ml Insulin (human) 5 µg/ml Tranferrin (human) 5 µg/ml Tranferrin (human) 67 µg/ml Phuronic F68 0.01% Norwich NY, USA) Phuronic F68 Norwich NY, USA) Phuronic F68 <td< td=""><td>L-Aspartic acid</td><td>10</td></td<>	L-Aspartic acid	10
L-Arginine 40 L-Arginine HCl 92.6 L-Arginine HCl 92.6 L-Aspartate 33.32 L-Cystine 2HCl 62.04 L-Cysteine HCl • H ₂ O 7.024 L-Glutamic acid 36.94 L-Glutamine 730 L-Glutamine 730 L-Histidine 3 L-Histidine HCl • H ₂ O 27.392 L-Hydroxypyroline 4 L-Isoleucine 73.788 L-Leucine 75.62 L-Lysine HCl 102.9 L-Methionine 21.896 L-Phenylalanine 43.592 L-Provoline 26.9 L-Serine 31.3 L-Threonine 53 L-Trytosine • 2Na 69.76 L-Valine 62.74 Penicillin/streptomycin 100 U/ml Insulin (human) 5 µg/ml Tranferrin (human) 5 µg/ml Primatone RL (Sheffield Products, 0.25% 0.25% Norwich NY, USA) Pluronic F68 Phuronic F68 0.01% (Serva, Heidelberg, FRG) <td< td=""><td>L-Asparagine H₂O</td><td>14.362</td></td<>	L-Asparagine H ₂ O	14.362
L-Arginine HCl 92.6 L-Aspartate 33.32 L-Cysteine 2HCl 62.04 L-Cysteine HCl • H ₂ O 7.024 L-Glutamic acid 36.94 L-Glutamic acid 36.94 L-Glutamine 730 L-Glycine 21.5 L-Histidine 3 L-Histidine 3 L-Histidine HCl • H ₂ O 27.392 L-Hydroxypyroline 4 L-Isoleucine 75.62 L-Lysine HCl 102.9 L-Methionine 21.896 L-Phonylalanine 43.592 L-Proline 26.9 L-Serine 31.3 L-Threonine 53 L-Tyrosine • 2Na 69.76 L-Valine 62.74 Penicillin/streptomycin 100 U/ml Insulin (human) 5 µg/ml Tranferrin (human) 5 µg/ml Primatone RL (Sheffield Products, 0.25% 0.25% Norwich NY, USA) Pluronic F68 Phuronic F68 0.01% (Serva, Heidelberg, FRG) Foetal calf serum	L-Arginine	40
L-Aspartate 33.32 L-Cystine 2HCl 62.04 L-Cysteine HCl • H ₂ O 7.024 L-Glutamic acid 36.94 L-Glutamice 730 L-Glutamice 21.5 L-Histidine 3 L-Histidine HCl • H ₂ O 27.392 L-Hydroxypyroline 4 L-Isoleucine 73.788 L-Leucine 75.62 L-Jysine HCl 102.9 L-Methionine 21.8966 L-Phonylalanine 43.592 L-Proline 26.9 L-Serine 31.3 L-Threonine 53 L-Tryptophan 11.008 L-Tyrosine • 2Na 69.76 L-Valine 62.74 Penicillin/streptomycin 100 U/ml Insulin (human) 5 µg/ml Tranferrin (human) 5 µg/ml Primatone RL (Sheffield Products, 0.25% 0.25% Norwich NY, USA) 7 Phuronic F68 0.01% (Serva, Heidelberg, FRG) 7 Foetal calf serum $0.3-3\%$ </td <td>L-Arginine HCl</td> <td>92.6</td>	L-Arginine HCl	92.6
L-Cystine 2HCl 62.04 L-Cysteine HCl • H ₂ O 7.024 L-Glutamicacid 36.94 L-Glycine 21.5 L-Histidine 3 L-Hydroxypyroline 4 L-Isoleucine 73.788 L-Leucine 75.62 L-Lysine HCl 102.9 L-Methionine 21.896 L-Phonylalanine 43.592 L-Proline 26.9 L-Serine 31.3 L-Threonine 53 L-Tryptophan 11.0008 L-Tyrosine • 2Na 69.76 L-Valine 62.74 Penicillin/streptomycin 100 U/ml Insulin (human) 5 µg/ml Tranferrin (human) 5 µg/ml Tranferrin (human) 67 µg/ml Primatone RL (Sheffield Products, 0.25% 0.25% Norwich NY, USA)	L-Aspartate	33.32
L-Cysteine HCl • H ₂ O 7.024 L-Glutamic acid 36.94 L-Glutamine 730 L-Glycine 21.5 L-Histidine 3 L-Hydroxypyroline 4 L-Isoleucine 75.62 L-Lysine HCl 102.9 L-Methionine 21.896 L-Phenylalanine 43.592 L-Proline 26.9 L-Serine 31.3 L-Threonine 53 L-Tryptophan 11.0008 L-Tyrosine • 2Na 69.76 L-Valine 62.74 Penicillin/streptomycin 100 U/ml Insulin (human) 5 µg/ml Tranferrin (human) 5 µg/ml Primatone RL (Sheffield Products, 0.25% 0.25% Norwich NY, USA) 0.01% Pluronic F68 0.01% (Serva, Heidelberg, FRG)	L-Cystine 2HCl	62.04
L-Glutamic acid 36.94 L-Glutamine 730 L-Glycine 21.5 L-Histidine 3 L-Histidine HCl • H ₂ O 27.392 L-Hydroxypyroline 4 L-Isoleucine 73.788 L-Luccine 75.62 L-Lysine HCl 102.9 L-Methionine 21.896 L-Phenylalanine 43.592 L-Proline 26.9 L-Serine 31.3 L-Threonine 53 L-Tyrosine • 2Na 69.76 L-Valine 62.74 Penicillin/streptomycin 100 U/ml Insulin (human) 5 µg/ml Tranferrin (human) 5 µg/ml Primatone RL (Shefield Products, 0.25% 0.25% Norwich NY, USA) Pluronic F68 Pluronic F68 0.01% (Serva, Heidelberg, FRG) $Foetal calf serum$	L-Cysteine HCl • H ₂ O	7.024
L-Glutamine 730 L-Glycine 21.5 L-Histidine 3 L-Histidine HCl • H ₂ O 27.392 L-Hydroxypyroline 4 L-Isoleucine 73.788 L-Leucine 75.62 L-Physine HCl 102.9 L-Methionine 21.896 L-Phenylalanine 43.592 L-Proline 26.9 L-Serine 31.3 L-Threonine 53 L-Typosine • 2Na 69.76 L-Valine 62.74 Penicillin/streptomycin 100 U/ml Insulin (human) 5 $\mu g/ml$ Tranferrin (human) 5 $\mu g/ml$ Primatone RL (Sheffield Products, 0.25% 0.25% Norwich NY, USA) Pluronic F68 Phuronic F68 0.01% (Serva, Heidelberg, FRG) Foetal calf serum	L-Glutamic acid	36.94
L-Glycine 21.5 L-Histidine 3 L-Histidine 3 L-Histidine 27.392 L-Hydroxypyroline 4 L-Isoleucine 73.788 L-Leucine 75.62 L-Lysine HCl 102.9 L-Methionine 21.896 L-Phenylalanine 43.592 L-Proline 26.9 L-Serine 31.3 L-Threonine 53 L-Tryptophan 11.008 L-Tyrosine • 2Na 69.76 L-Valine 62.74 Penicillin/streptomycin 100 U/ml Insulin (human) 5 µg/ml Tranferrin (human) 5 µg/ml Bovine serum albumin 67 µg/ml Primatone RL (Sheffield Products, 0.25% 0.25% Norwich NY, USA) Pluronic F68 Phuronic F68 0.01% (Serva, Heidelberg, FRG) Foetal calf serum	L-Glutamine	730
L-Histidine 3 L-Histidine 27.392 L-Hydroxypyroline 4 L-Isoleucine 73.788 L-Luceine 73.788 L-Leucine 75.62 L-Lysine HCl 102.9 L-Methionine 21.896 L-Phenylalanine 43.592 L-Proline 26.9 L-Serine 31.3 L-Threonine 53 L-Trytosine • 2Na 69.76 L-Valine 62.74 Penicillin/streptomycin 100 U/ml Insulin (human) 5 µg/ml Tranferrin (human) 15 µg/ml Bovine serum alburnin 67 µg/ml Primatone RL (Sheffield Products, 0.25% 0.25% Norwich NY, USA) Pluronic F68 Pluronic F68 0.01% (Serva, Heidelberg, FRG) Foetal calf serum	L-Glycine	21.5
L-Histidine HCl • H_2O 27.392 L-Hydroxypyroline 4 L-Isoleucine 73.788 L-Leucine 75.62 L-Lysine HCl 102.9 L-Methionine 21.896 L-Phenylalanine 43.592 L-Proline 26.9 L-Serine 31.3 L-Threonine 53 L-Tryotophan 11.008 L-Tyrosine • 2Na 69.76 L-Valine 62.74 Penicillin/streptomycin 100 U/ml Insulin (human) 5 µg/ml Tranferrin (human) 15 µg/ml Bovine serum albumin 67 µg/ml Primatone RL (Sheffield Products, 0.25% 0.25% Norwich NY, USA) Pluronic F68 Phuronic F68 0.01% (Serva, Heidelberg, FRG) Foetal calf serum	L-Histidine	3
L-Hydroxypyroline4L-Isoleucine73.788L-Leucine75.62L-Lysine HCl102.9L-Methionine21.896L-Phenylalanine43.592L-Proline26.9L-Serine31.3L-Threonine53L-Tyytophan11.008L-Tyrosine • 2Na69.76L-Valine62.74Penicillin/streptomycin100 U/mlInsulin (human)5 $\mu g/ml$ Trafferrin (human)15 $\mu g/ml$ Primatone RL (Sheffield Products, Norwich NY, USA)0.25%Norwic F680.01%(Serva, Heidelberg, FRG)0.3-3%	L-Histidine HCl • H ₂ O	27.392
L-Isoleucine 73.788 L-Leucine 75.62 L-Lysine HCl 102.9 L-Methionine 21.896 L-Phenylalanine 43.592 L-Proline 26.9 L-Serine 31.3 L-Threonine 53 L-Tryptophan 11.008 L-Yaline 69.76 L-Valine 62.74 Penicillin/streptomycin 100 U/ml Insulin (human) 5 µg/ml Tranferrin (human) 5 µg/ml Primatone RL (Sheffield Products, 0.25% 0.25% Norwich NY, USA) Pluronic F68 Phuronic F68 0.01% (Serva, Heidelberg, FRG) Foetal calf serum	L-Hydroxypyroline	4
L-Leucine 75.62 L-Lysine HCl 102.9 L-Methionine 21.896 L-Phenylalanine 43.592 L-Proline 26.9 L-Serine 31.3 L-Threonine 53 L-Tryptophan 11.008 L-Tyrosine • 2Na 69.76 L-Valine 62.74 Penicillin/streptomycin 100 U/ml Insulin (human) 5 µg/ml Tranferrin (human) 15 µg/ml Primatone RL (Sheffield Products, 0.25% 0.25% Norwich NY, USA) Pluronic F68 Phuronic F68 0.01% (Serva, Heidelberg, FRG) Foetal calf serum	L-Isoleucine	73.788
L-Lysine HCl 102.9 L-Methionine 21.896 L-Phenylalanine 43.592 L-Proline 26.9 L-Serine 31.3 L-Threonine 53 L-Tryptophan 11.0008 L-Tyrosine • 2Na 69.76 L-Valine 62.74 Penicillin/streptomycin 100 U/ml Insulin (human) 5 µg/ml Tranferrin (human) 15 µg/ml Bovine serum albumin 67 µg/ml Primatone RL (Sheffield Products, 0.25% 0.25% Norwich NY, USA) 9 Pluronic F68 0.01% (Serva, Heidelberg, FRG) 5 Foetal calf serum 0.3-3%	L-Leucine	75.62
L-Methionine21.896L-Phenylalanine 43.592 L-Proline 26.9 L-Serine 31.3 L-Threonine 53 L-Tryptophan 11.008 L-Tyrosine • 2Na 69.76 L-Valine 62.74 Penicillin/streptomycin 100 U/mlInsulin (human) 5 µg/mlTranferrin (human) 67 µg/mlPrimatone RL (Sheffield Products, Norwich NY, USA) 0.25% Norwich NY, USA) 0.01% Foetal calf serum $0.3-3\%$	L-Lysine HCl	102.9
L-Phenylalanine 43.592 L-Proline 26.9 L-Serine 31.3 L-Threonine 53 L-Threonine 69.76 L-Tyrosine • 2Na 69.76 L-Valine 62.74 Penicillin/streptomycin 100 U/mlInsulin (human) 5 µg/mlTrafferrin (human) 15 µg/mlPrimatone RL (Sheffield Products, 0.25% Norwich NY, USA)Pluronic F68Pluronic F68 0.01% (Serva, Heidelberg, FRG) $0.3-3\%$	L-Methionine	21.896
$\begin{array}{llllllllllllllllllllllllllllllllllll$	L-Phenylalanine	43.592
L-Serine 31.3 L-Threonine 53 L-Tryptophan 11.008 L-Tryptosine • 2Na 69.76 L-Valine 62.74 Penicillin/streptomycin 100 U/mlInsulin (human) 5 µg/mlTranferrin (human) 15 µg/mlBovine serum albumin 67 µg/mlPrimatone RL (Sheffield Products, Norwich NY, USA) 0.25% Norwich F68 0.01% (Serva, Heidelberg, FRG)Foetal calf serumFoetal calf serum $0.3-3\%$	L-Proline	26.9
L-Threonine53L-Tryptophan11.008L-Tyrosine • 2Na69.76L-Valine62.74Penicillin/streptomycin100 U/mlInsulin (human)5 µg/mlTranferrin (human)15 µg/mlBovine serum albumin67 µg/mlPrimatone RL (Sheffield Products,0.25%Norwich NY, USA)25%Pluronic F680.01%(Serva, Heidelberg, FRG)0.3-3%	L-Serine	31.3
L-Tryptophan11.008L-Tyrosine • 2Na69.76L-Valine62.74Penicillin/streptomycin100Insulin (human)5 µg/mlTranferrin (human)15 µg/mlBovine serum albumin67 µg/mlPrimatone RL (Sheffield Products,0.25%Norwich NY, USA)100Pluronic F680.01%(Serva, Heidelberg, FRG)0.3-3%	L-Threonine	53
L-Tyrosine • 2Na69.76L-Valine62.74Penicillin/streptomycin100 U/mlInsulin (human)5 µg/mlTranferrin (human)15 µg/mlBovine serum albumin67 µg/mlPrimatone RL (Sheffield Products,0.25%Norwich NY, USA)Pluronic F68Pluronic F680.01%(Serva, Heidelberg, FRG)0.3-3%	L-Tryptophan	11.008
L-Valine62.74Penicillin/streptomycin100 U/mlInsulin (human)5 µg/mlTranferrin (human)15 µg/mlBovine serum albumin67 µg/mlPrimatone RL (Sheffield Products, Norwich NY, USA)0.25%Norwich NY, USA)0.01%Pluronic F680.01%(Serva, Heidelberg, FRG)0.3-3%	L-Tyrosine • 2Na	69.76
Penicillin/streptomycin 100 U/ml Insulin (human) 5 µg/ml Tranferrin (human) 15 µg/ml Bovine serum albumin 67 µg/ml Primatone RL (Sheffield Products, 0.25% Norwich NY, USA) 9 Pluronic F68 0.01% (Serva, Heidelberg, FRG) 0.3-3%	L-Valine	62.74
Insulin (human)5 µg/mlTranferrin (human)15 µg/mlBovine serum albumin67 µg/mlPrimatone RL (Sheffield Products,0.25%Norwich NY, USA)Pluronic F68Pluronic F680.01%(Serva, Heidelberg, FRG)0.3-3%	Penicillin/streptomycin	100 U/ml
Tranferrin (human)15 µg/mlBovine serum albumin67 µg/mlPrimatone RL (Sheffield Products,0.25%Norwich NY, USA)Pluronic F68Pluronic F680.01%(Serva, Heidelberg, FRG)0.3-3%	Insulin (human)	5 μg/ml
Bovine serum albumin67 µg/mlPrimatone RL (Sheffield Products, Norwich NY, USA)0.25%Pluronic F680.01%(Serva, Heidelberg, FRG)0.3-3%	Tranferrin (human)	15 μg/ml
Primatone RL (Sheffield Products, Norwich NY, USA)0.25%Pluronic F680.01%(Serva, Heidelberg, FRG)0.3-3%	Bovine serum albumin	67 μg/ml
Norwich NY, USA) Pluronic F68 0.01% (Serva, Heidelberg, FRG) Foetal calf serum 0.3-3%	Primatone RL (Sheffield Products,	0.25%
Pluronic F680.01%(Serva, Heidelberg, FRG)0.3-3%Foetal calf serum0.3-3%	Norwich NY, USA)	
(Serva, Heidelberg, FRG) Foetal calf serum 0.3-3%	Pluronic F68	0.01%
Foetal calf serum 0.3-3%	(Serva, Heidelberg, FRG)	
	Foetal calf serum	0.3-3%

The centrifugate was washed with isotonic phosphate buffer (PBS; 0.2 g/1 KC1, 0.2 g/1 KH₂PO₄, 8.0 g/1 NaC1, 2.16 g/l Na₂HPO₄.7H₂0), which had been treated with 5% dimethylformamide, 10 mM benzamidine, 100 U/ml aprotinin, 10 µM leupeptin, 1 µM pepstatin, 1 mM o-phenanthroline, 5 45 mM iodoacetamide, 1 mM phenyl-methylsulphonyl fluoride (referred to hereinafter as PBS-M). The washed cells were extracted at a density of 2.5.108 cells/ml in PBS-M with Triton X-100 (final concentration 1.0%). The cell extract was clarified by centrifugation (15,000×g, 1 hour; 100,000×g, 1 50 hour).

Example 3

Production of Monoclonal (TNF-BP) Antibodies

A centrifugation supernatant from the cultivation of HL60 cells on an experimental laboratory scale, obtained according to Example 2, was diluted with PBS in the ratio 1:10. The diluted supernatant was applied at 4° C. (flow rate: 0.2 60 ml/min.) to a column which contained 2 ml of Affigel 10 (Bio Rad Catalogue No. 153-6099) to which had been coupled 20 mg of recombinant human TNF- α [Pennica, D. et al. (1984) Nature 312, 724; Shirai, T. et al. (1985) Nature 313, 803; Wang, A. M. et al. (1985) Science 228, 149] according to the 65 recommendations of the manufacturer. The column was washed at 4° C. and a throughflow rate of 1 ml/min firstly with 20 ml of PBS which contained 0.1% Triton X 114 and thereafter with 20 ml of PBS. Thus-1-enriched TNF-BP was eluted at 22° C. and a flow rate of 2 ml/min with 4 ml of 100 mM glycine, pH 2.8, 0.1% decyl-maltoside. The eluate was concentrated to 10 μ l in a Centricon 30 unit [Amicon].

10 μ l of this eluate were mixed with 20 μ l of complete 5 Freund's adjuvant to give an emulsion. 10 μ l of the emulsion were injected according to the procedure described by Holmdahl, R. et al. [(1985), J. Immunol. Methods 32, 379] on each of days 0, 7 and 12 into a hind paw of a narcotized Balb/c mouse.

The immunized mice were sacrificed on day 14, the popliteal lymph nodes were removed, minced and suspended by repeated pipetting in Iscove's medium (IMEM, GIBCO Catalogue No. 074-2200) which contained 2 g/l NaHCO3. According to a modified procedure of De St. Groth and 15 Scheidegger [J. Immunol. Methods (1980), 35, 1] 5×10^7 cells of the lymph nodes were fused with 5×10^7 PAI mouse myeloma cells (J. W. Stocker et al., Research Disclosure, 217, May 1982, 155-157) which were in logarithmic growth. The cells were mixed, collected by centrifugation and resus- 20 pended in 2 ml of 50% (v/v) polyethylene glycol in IMEM at room temperature by slight shaking and diluted by the slow addition of 10 ml of IMEM during careful shaking for 10 minutes. The cells were collected by centrifugation and resuspended in 200 ml of complete medium [IMEM+20% foetal 25 calf serum, glutamine (2.0 mM), 2-mercaptoethanol $(100 \,\mu\text{l})$, 100 µM hypoxanthine, 0.4 µM aminopterine and 16 µM thymidine (HAT)]. The suspension was distributed on 10 tissue 10 culture dishes each containing 96 wells and incubated at 37° C. for 11 days without changing the medium in an atmo- 30 sphere of 5% CO₂ and a relative humidity of 98%.

The antibodies are distinguished by their inhibitory action on the binding of TNF to HL60 cells or by their binding to antigens in the filter test according to Example 1. The following procedure was used to detect the biological activity of 35 anti(TNF-BP) antibodies: 5×10^6 HL60 or U937 cells were incubated in complete RPMI 1640 medium together with affinity-purified monoclonal anti-(TNF-BP) antibodies or control antibodies (i.e. those which are not directed against TNF-BP) in a concentration range of 1 ng/ml to 10 μ g/ml. ⁴⁰ After incubation at 37° C. for one hour the cells were collected by centrifugition and washed with 4.5 ml of PBS at 0° C. They were resuspended in 1 ml of complete RPMI 1640 medium (Example 2) which additionally contained 0.1% sodium azide and 125 I-TNF α (10⁶ cpm/ml) with or without 45 the addition of unlabelled TNF α (see above). The specific radioactivity of the $^{125}\mbox{I-TNF}\alpha$ amounted to 700 Ci/mmol. The cells were incubated at 4° C. for 2 hours, collected and washed 4 times at 0° C. with 4.5 ml of PBS which contained 1% BSA and 0.001% Triton X 100 (Fluka). The radioactivity 50 bound to the cells was measured in a y-scintillation counter. The cell-bound radioactivity of cells which had not been treated with anti-(TNF-BP) antibodies was determined in a comparative experiment (approximately 10 000 cpm/5×10⁶ 55 cells).

Example 4

Affinity Chromatography

60

For the further purification, a monoclonal anti-(55 kD TNF-BP) antibody (2.8 mg/ml gel), obtained according to Example 3, TNF α (3.9 mg/ml gel) and bovine serum albumin (BSA, 8.5 mg/ml gel) were each covalently coupled to CNBractivated Sepharose 4B (Pharmacia, Uppsala, Sweden) 65 according to the directions of the manufacturer. The cell extract obtained according to Example 2 was passed through

the thus-prepared columns which were connected in series in the following sequence: BSA-Sepharose pre-column, immune affinity column (anti-(55 kD-TNF-BP) antibody), TNF α -ligand affinity column. After completion of the application the two last-mentioned columns were separated and washed individually with in each case 100 ml of the following buffer solutions: (1) PBS, 1.0% Triton X-100, 10 mM benzamidine, 100 U/ml aprotinin; (2) PBS, 0.1% Triton X-100, 0.5M NaCl, 10 mM ATP, 10 mM benzamidine, 100 U/ml aprotinin; and (3) PBS, 0.1% Triton X-100, 10 mM benzamidine, 100 U/ml aprotinin. Not only the immune affinity column, but also the TNF α -ligand affinity column were then each eluted with 100 mM glycine pH 2.5, 100 mM NaCl, 0.2% decylmaltoside, 10 mM benzamidine, 100 U/ml aprotinin. The fractions of each column which were active in the filter test according to Example 1 were thereafter combined and neutralized with 1M Tris pH 8.0.

The thus-combined TNF-BP active fractions of the immune affinity chromatography on the one hand and of the $TNF\alpha$ -ligand affinity chromatography on the other hand were, for further purification, again applied to in each case one small TNFa-ligand affinity column. Thereafter, these two columns were washed with in each case 40 ml of (1) PBS, 1.0% Triton X-100, 10 mM benzamidine, 100 U/ml aprotinin, (2) PBS, 0.1% Triton X-100, 0.5M NaCl, 10 mM ATP, 10 mM benzamidine, 100 U/ml aprotinin, (3) PBS, 0.1% Triton X-100, (4) 50 mM Tris pH 7.5, 150 mM NaCl, 1.0% NP-40, 1.0% desoxycholate, 0.1% SDS, (5) PBS, 0.2% decyl-maltoside. Subsequently, the columns were eluted with 100 mM glycine pH 2.5, 100 mM NaCl, 0.2% decylmaltoside. Fractions of 0.5 ml from each column were collected and the fractions from each column which were active according to the filter test (Example 1) were combined and concentrated in a Centricon unit (Amicon, molecular weight exclusion 10,000).

Example 5

Separation by Means of HPLC

The active fractions obtained according to Example 4 were each applied according to their different source (immune or ligand affinity chromatography) to C1/C8 reversed phase HPLC columns (ProRPC, Pharmacia, 5×20 mm) which had been equilibrated with 0.1% trifluoroacetic acid, 0.1% octylglucoside. The columns were then eluted with a linear acetonitrile gradient (0-80%) in the same buffer at a flow of 0.5 ml/min. Fractions of 1.0 ml were collected from each column and the active fractions from each column were combined (detection according to Example 1).

Example 6

Separation by Means of SDS-PAGE

The fractions which were obtained according to Example 5 and which were active according to the filter test (Example 1) were further separated by SDS-PAGE according to [34]. For this purpose, the samples were heated to 95° C. for 3 minutes in SDS sample buffer and subsequently separated electrophoretically on a 12% acrylamide separation gel with a 5% collection gel. The following standard proteins were used as a reference for the determination of the apparent molecular weights on the SDS-PAGE gel: phosphorylase B (97.4 kD), BSA (66.2 kD), ovalbumin (42.7 kD), carboanhydrase (31.0 kD), soya trypsin inhibitor (21.5 kD) and lysozyme (14.4 kD).

Under the mentioned conditions there were obtained for samples which has been obtained according to Example 4 by TNF- α -ligand affinity chromatography of immune affinity chromatography eluates and which had been further separated by HPLC according to Example 5 two bands of 55 kD 5 and 51 kD as well as three weaker bands of 38 kD, 36 kD and 34 kD. These bands were transferred electro-phoretically during 1 hour at 100 V in 25 mM Tris, 192 mM glycine, 20% methanol on to a PVDF membrane (Immobilon, Millipore, Bedford, Mass. USA) in a Mini Trans Blot System (BioRad, Richmond, Calif., USA). Thereafter, the PVDF membrane was either protein-stained with 0.15% Serva-Blue (Serva, Heidelberg, FRG) in methanol/water/glacial acetic acid (50/ 40/10 parts by volume) or blocked with skimmed milk powder and subsequently incubated with 125 I-TNF α according to 15 the filter test conditions described in Example 1 in order to detect bands having TNF-BP activity. This showed that all bands produced in the protein staining bonded TNF α specifically. In the Western blot according to Towbin et al. [38] all of these bands also bonded the monoclonal anti-55 kD-TNF-BP 20 antibody produced according to Example 3. In this case, a procedure according to that described in Example 1 with Na¹²⁵I radioactively-labelled, affinity-purified (mouse immuno-globulin-Sepharose-4B affinity column) rabbitanti-mouse-immunoglobulin antibody was used for the auto-25 radiographic detection of this antibody.

Samples which had been obtained according to Example 4 by two-fold TNF- α -ligand affinity chromatography of the throughput of the immune affinity chromatography and which had been further separated by HPLC according to Example 5 showed under the above-specified SDS-PAGE and blot transfer conditions two additional bands of 75 kD and 65 kD, both of which bonded TNF specifically in the filter test (Example 1). In the Western blot according to Towbin et al. (see above) the proteins of these two bands did not react with anti-(55 kD TNF-BP) antibody produced according to Example 3. They reacted, however, with a monoclonal antibody which had been produced starting from the 75 kD band (anti-75 kD TNF-BP antibody) according to Example 3.

Example 7

Amino Acid Sequence Analysis

For the amino acid sequence analysis, the fractions which 45 had been obtained according to Example 5 and which were active according to the filter test (Example 1) were separated using the SDS-PAGE conditions described in Example 6, but now reducing (SDS sample buffer with 125 mM dithiothreitol). The same bands as in Example 6 were found, but because 50 of the reducing conditions of the SDS-PAGE in comparison to Example 6 all showed an about 1-2 kD higher molecular weight. These bands were then transferred according to Example 6 on to PVDF membranes and stained with 0.15% Serva-Blue in methanol/water/glacial acetic acid (50/400/10 55 parts by volume) for 1 minute, decolorized with methanol/ water/glacial acetic acid (45/48/7 parts by volume), rinsed with water, dried in air and thereafter cut out. The conditions given by Hunkapiller [34] were adhered to in all steps in order to avoid N-terminal blocking. Initially, the purified TNF-BP 60 were used unaltered for the amino acid sequencing. In order to obtain additional sequence information, the TNF-BP after reduction and S-carboxymethylation [Jones, B. N. (1986) in "Methods of Protein Micro-characterisation", J. E. Shively, ed., Humana Press, Clifton N.J., 124-125] were cleaved with 65 cyanogen bromide (Tarr, G. E. in "Methods of Protein Microcharacterisation", 165-166, loc. cit.), trypsin and/or protein-

ase K and the peptides were separated by HPLC according to known methods of protein chemistry. Thus-prepared samples were then sequenced in an automatic gas phase microsequencing apparatus (Applied Biosystems Model 470A, ABI, Foster City, Calif., USA) with an on-line automatic HPLC PTH amino acid analyzer (Applied Biosystems Model 120, ABI see above) connected to the outlet, whereby the following amino acid sequences were determined:

1. For the 55 kD band (according to non-reducing SDS-10 PAGE):

Leu-Val-Pro-His-Leu-Gly-Asp-Arg-Glu-Lys-Arg-Asp-Ser-Val-Cys-Pro-Gln-Gly-Lys-Tyr-Ile-His-Pro-Gln-X-Asn-Ser-Ile (SEQ ID NO: 5),

and

Ser-Thr-Pro-Glu-Lys-Glu-Gly-Glu-Leu-Glu-Gly-Thr-Thr-Thr-Lys (SEQ ID NO: 6) in which X stands for an amino acid residue which could not be determined,

- 2. for the 51 kD and 38 kD bands (according to non-reducing SDS-PAGE):
- Leu-Val-Pro-His-Leu-Gly-Asp-Arg-Glu (SEQ ID NO: 15)
- 3. for the 6510 band (according to non-reducing SDS-PAGE) In the N-terminal sequencing of the 6510 band two parallel sequences were determined up to the 15th residue without interruption. Since one of the two sequences corresponded to a partial sequence of ubiquitin [36,37], the following sequence was derived for the 65 kD band:
 - Leu-Pro-Ala-Gln-Val-Ala-Phe-X-Pro-Tyr-Ala-Pro-Glu-Pro-Gly-Ser-Thr-Cys. (SEQ ID NO: 10)
 - in which X stands for an amino acid residue which could not be determined.
- Additional peptide sequences for 75(65)kDa-TNF-BP were determined:
 - Ile-X-Pro-Gly-Phe-Gly-Val-Ala-Tyr-Pro-Ala-Leu-Glu (SEQ ID NO: 11)

and

Ser-Gln-Leu-Glu-Thr-Pro-Glu-Thr-Leu-Leu-Gly-Ser-Thr-Glu-Glu-Lys-Pro-Leu (SEQ ID NO: 7) and Val-Phe-Cys-Thr (SEQ ID NO: 8)

and

Asn-Gln-Pro-Gln-Ala-Pro-Gly-Val-Glu-Ala-Ser-Gly-Ala-Gly-Glu-Ala (SEQ ID NO: 9) and Leu-Cys-Ala-Pro (SEQ ID NO: 12)

and

Val-Pro-His-Leu-Pro-Ala-Asp (SEQ ID NO: 13)

and

Gly-Ser-Gln-Gly-Pro-Glu-Gln-Gln-X-X-Leu-Ile-X-Ala-Pro (SEQ ID NO: 14), in which X stands for an amino acid residue which could not be determined.

Example 8

Determination of Base Sequences of Complementary DNA (cDNA)

Starting from the amino acid sequence according to formula IA there were synthesized having regard to the genetic code for the amino acid residues 2-7 and 17-23 corresponding completely degenerated oligonucleotides in suitable complementarity ("sense" and "antisense" oligonucleotides). Total cellular RNA was isolated from HL60 cells [42,43] and the first cDNA strand was synthesized by oligo-dT priming or by priming with the "antisense" oligonucleotide using a cDNA synthesis kit (RPN 1256, Amersham, Amersham; England) according to the instructions of the manufacturer. This cDNA strand and the two synthesized degenerate "sense" and "antisense" oligonucleotides were used in a polymerase chain reaction (PCR, Perkin Elmer Cetus, Norwalk, Conn., USA

according to the instructions of the manufacturer) to synthesize as a cDNA fragment the base sequence coding for the amino acid residues 8-16 (formula IA). The base sequence of this cDNA fragment accorded to: 5'-AGGGAGAA-GAGAGATAGTGTGTGTCCC-3' (SEQ ID NO: 16). This 5 cDNA fragment was used as a probe in order to identify according to a known procedure a cDNA clone coding for the 55 kD TNF-BP in a Xgt11-cDNA gene bank from human placenta (42, 43). This clone was then cut according to usual methods from the X-vector and cloned in the plasmids pUC18 10(Pharmacia, Uppsala, Sweden) and pUC19 (Pharmacia, Uppsala, Sweden) and in the M13 mp 18/M13 mp 19 bacteriophage (Pharmacia, Uppsala, Sweden) (42, 43). The nucleotide sequence of this cDNA clone was determined using a Sequenase kit (U.S. Biochemical, Cleveland, Ohio, USA) accord- 15 ing to the details of the manufacturer. The nucleotide sequence and the amino acid sequence derived therefrom for the 55 kD TNF-BP and its signal peptide (amino acid "-28" to amino acid "0") is given in FIG. 1 using the abbreviations for bases such as amino acids usual in the state of the art. From 20 sequence comparisons with other already known receptor protein sequences there can be determined a N-terminal domain containing approximately 180 amino acids and a C-terminal domain containing 220 amino acids which are separated from one another by a transmembrane region of 19 25 amino acids (underlined in FIG. 1) which is typical according to the sequence comparisons. Hypothetical glycosylation sites are characterized in FIG. 1 by asterisks above the corresponding amino acid.

Essentially analogous techniques were used to identify 30 75/65 kD TNF-BP-coding partial cDNA sequences, whereby however, in this case genomic human DNA and completely degenerated 14-mer and 15-mer "sense" and "antisense" oligonucleotides derived from peptide IIA were used in order to produce a primary 26 by cDNA probe in a polymerase chain 35 duced two stop codons of the translation after amino acid 182. reaction. This cDNA probe was then used in a HL-60 cDNA library to identify cDNA clones of different lengths. This cDNA library was produced using isolated HL60 RNA and a cDNA cloning kit (Amersham) according to the details of the manufacturer. The sequence of such a cDNA clone is given in 40 FIG. 4, whereby repeated sequencing lead to the following correction. A threonine coded by "ACC" not "TCC", has to be at position 3 instead of the serine.

Example 9

Expression in COS 1 Cells

Vectors starting from the plasmid "pN11" were constructed for the expression in COS cells. The plasmid "pN11" 50 contains the efficient promoter and enhancer of the "major gene of human cytomegalovirus immediate-early" ("HCMV"; Boshart et al., Cell 41, 521-530, 1985). After the promoter there is situated a short DNA sequence which contains several restriction cleavage sites, which are present only 55 once in the plasmid ("polylinker"), inter alia the cleavage sites for HindIII, Ball, BamHI and PvuII (see sequence). PvuII

5'-AAGCTTGGCCAGGATCCAGCTGACT-GACTGATCGCGAGATC-3' (SEQ ID NO: 17) 3'-TTCGAACCGGTCCTAGGTCGACTGACT-GACTAGCGCTCTAG-5' (SEQ ID NO: 18)

After these cleavage sites there are situated three translation stop codons in all three reading frames. After the polylinker sequence there is situated the 2nd intron and the 65 polyadenylation signal of the preproinsulin gene of the rat (Lomedico et al., Cell 18, 545-558, 1979). The plasmid also

contains the replication origin of the SV40 virus and a fragment from pBR322 which confers E. coli-bacteria ampicillin resistance and permits the replication of the plasmid in E. coli.

For the construction of the expression vector "pN123", this plasmid "pN11" was cleaved the restriction endo-nuclease PvuII and subsequently treated with alkaline phosphatase. The dephosphorylated vector was thereafter isolated from an agarose gel (V1). The 5'-projecting nucleotides of the EcoRIcleaved 1.3 kb fragment of the 55 kD TNF-BP-cDNA (see Example 8) were filled in using Klenow enzyme. Subsequently, this fragment was isolated from an agarose gel (F1). Thereafter, V1 and F1 were joined together using T4-ligase. E. coli HB101 cells were then transformed with this ligation batch according to known methods [42]. By means of restriction analyses and DNA sequencing according to known methods [42] there were identified transformants which had been transformed with a plasmid and which contained the 1.3 kb EcoRI fragment of the 55 kD TNF-BP-cDNA in the correct orientation for expression via the HCMV-promoter. This vector received the designation "pN123".

The following procedure was used for the construction of the vector "pK19". A DNA fragment which contained only the cDNA coding for the extracellular part of the 55 kD TNF-BP (amino acids -28 to 182 according to FIG. 1) was obtained by PCR technology (Saiki et al., Science 230, 1350-1354, 1985, see also Example 8). The following oligonucleotides were used in order to amplify the cDNA from "pN123" coding for the extracellular part of the 55 kD TNF-BP:

BAMHI 5'-CACAGGGATCCATAGCTGTCTG-GCATGGGCCTCTCCAC-3' (SEQ ID NO: 19) ASP718

3'-CGTGACTCCTGAGTCCGTGGTGTAT-

TATCTCTAGACCA TGGCCC-5' (SEQ ID NO: 20) By means of these oligonucleotides there were also intro-The thus-amplified DNA fragment was cleaved with BamHI and Asp718, the thereby resulting projecting ends were filled in using Klenow enzyme and this fragment was subsequently isolated from an agarose gel (F2). F2 was then ligated with V1 and the entire batch was used for the transformation of E. coli HB101, as already described. Transformants which had been transformed with a plasmid containing the DNA fragment in the correct orientation for the expression via the HCMVpromoter were identified by DNA sequencing (see above). The plasmid isolated therefrom received the designation "pK19".

Transfection of the COS cells with the plasmids "pN123" or "pK19" was carried out according to the lipofection method published by Feigner et al. (Proc. Natl. Acad. Sci. USA 84, 7413-7417, 1987). 72 hours after the transfection had been effected the cells transfected with "pN123" were analyzed for binding with ¹²⁵I-TNF α according to known methods. The results of the Scatchard analysis [Scatchard, G., Ann. N.Y. Acad. Sci. 51, 660, 1949] of the thus-obtained binding data (FIG. 2A) is given in FIG. 2B. The culture supernatants of the cells transfected with "pK19" were investigated in a "sandwich" test. For this purpose, PVC microtitre plates (Dynatech, Arlington, Va., USA) were sensitized with 100 µl/well of a rabbit-anti-mouse immunoglobulin (10 60 µg/ml PBS). Subsequently, the plates were washed and incubated (3 hours, 20° C.) with an anti-55 kD TNF-BP antibody which had been detected by its antigen binding and isolated according to Example 3, but which did not inhibit the TNFbinding to cells. The plates were then again washed and incubated overnight at 4° C. with 100 µl/well of the culture supernatant (diluted 1:4 with buffer A containing 1% skimmed milk powder: 50 mM Tris/HCl pH 7.4, 140 mM

NaCl, 5 mM EDTA, 0.02% Na azide). The plates were emptied and incubated at 4° C. for 2 hours with buffer A containing ¹²⁵I-TNF α (10⁶ cpm/ml, 100 µl/well) with or without the addition of 2 µg/ml of unlabelled TNF. Thereafter, the plates were washed 4 times with PBS, the individual wells were cut ⁵ out and measured in a λ -counter. The results of 5 parallel transfections (columns # 2, 3, 4, 6 and 7), of two control transfections

with the pN11 vector (columns # 1, 5) and of a control with HL60 cell lysate (column # 8) are given in FIG. 3.

Example 10

Expression in Insect Cells

The plasmid "pVL941" (Luckow and Summers, 1989, "High Level Expression of Nonfused Foreign Genes with *Autographa california* Nuclear Polyhedrosis virus Expression Vectors", Virology 170, 31-39) was used for the expression in a baculovirus expression system and was modified as 20 follows. The single EcoRI restriction cleavage site in "pVL941" was removed by cleaving the plasmid with EcoRI and the projecting 5'-end was filled in with Klenow enzyme. The plasmid pVL941/E obtained therefrom was digested with BamHI and Asp718 and the vector trunk was subse-25 quently isolated from an agarose gel. This fragment was ligated with a synthetic oligonucleotide of the following sequence:

BamHI EcoRI Asp718

5'-GATCCAGAATTCATAATAG-3' (SEQ ID NO: 21) 3'-GTCTTAAGTATTATCCATG-5' (SEQ ID NO: 22)

E. coli HB101 was transformed with the ligation batch and transformants containing a plasmid in which the oligonucleotide had been incorporated correctly were identified by restriction analysis and DNA sequencing according to known 35. methods (see above); this plasmid was named "pNR704". For the construction of the transfer vector "pN113", this plasmid "pNR704" was cleaved with EcoRI, treated with alkaline phosphatase and the thus-1-produced vector trunk (V2) was subsequently isolated from an agarose gel. The 1.3 kb frag- 40 ment of the 55 kD TNF-BP-cDNA cleaved with EcoRI as above was ligated with fragment V2. Transformants obtained with this ligation batch, which contained a plasmid containing the cDNA insert in the correct orientation for the expression via the polyhedron promoter, were identified (see 45 above). The vector isolated therefrom received the designation "pN113".

The following procedure was used for the construction of the transfer vector "pN119". The 1.3 kb EcoRI/EcoRI fragment of the 55 kD TNF-BP cDNA in the "pUC19" plasmid 50 (see Example 8) was digested with BanI and ligated with the following synthetic oligonucleotide:

BanI Asp718

5'-GCACCACATAATAGAGATCTGGTACCGGGAA-3' (SEQ ID NO: 23)

3'-GTGTATTATCTCTAGACCATGGCCC-5' (SEQ ID NO: 24)

Two stop codons of the translation after amino acid 182 and a cleavage site for the restriction endo-nuclease Asp718 are incorporated with the above adaptor. After carrying out ligation the batch was digested with EcoRI and Asp718 and the partial 55 kD TNF-BP fragment (F3) was isolated. Furthermore, the plasmid "pNR704", likewise cleaved with Asp718 and EcoRI, was ligated with F3 and the ligation batch was transformed into *E. coli HB*101. The identification of the 65 transformants which contained a plasmid in which the partial 55 kD TNF-BP cDNA had been correctly integrated for the 20

expression was effected as already described. The plasmid isolated from these transformants received the name "pN119".

The following procedure was used for the construction of the transfer vector "pN124". The cDNA fragment coding for the extracellular part of the 55 kD TNF-BP, described in Example 9, was amplified with the specified oligo-nucleotides with the aid of PCR technology as described in Example 9. This fragment was cleaved with BamHI and Asp718 and isolated from an agarose gel (F4). The plasmid "pNR704" was also cleaved with BamHI and Asp718 and the vector trunk (V4) was isolated (see above). The fragments V4 and F4 were ligated, *E. coli* HB101 was transformed therewith and the recombinant transfer vector "pN124" was identified and isolated as described.

The following procedure was used for the transfection 10 of the insect cells. 3 µg of the transfer vector "pN113" were transfected with 1 µg of DNA of the Autographa californica nuclear polyhedrosisvirus (AcMNPV) (EP 127839) in Sf9 cells (ATCC CRL 1711). Polyhedron-negative viruses were identified and purified from "plaques" [52]. Sf9 cells were again infected with these recombinant viruses as described in [52]. After 3 days in the culture the infected cells were investigated for TNF-binding using 125 I-TNF α . For this purpose, the transfected cells were washed from the cell culture dish with a Pasteur pipette and resuspended at a cell density of 5×10^6 cells/ml of culture medium [52] which contained 10 ng/ml of ¹²⁵I-TNF- α , not only in the presence of, but also in the absence of 5 μ g/ml of non-labelled TNF- α and incubated on ice for 2 hours. Thereafter, the cells were washed with pure culture medium and the cell-bound radio-activity was counted in a γ -counter (see Table 2).

TABLE 2

Cells	Cell-bound radioactivity per 10 ⁶ cells
Non-infected cells	60 cpm
Infected cells	$1600 \pm 330 \text{ cpm}^{-1}$

1) Average and standard deviation from 4 experiments

Example 11

Analogously to the procedure described in Example 9, the cDNA fragment coding for the extracellular region of the 55 kDa TNF-BP was amplified in a polymerase chain reaction, but now using the following oligonucleotides as the primer: Oligonucleotide 1:

Sst I 5'-TAC GAG CTC GGC CAT AGC TGT CTG GCA TG-3' (SEQ ID NO: 25)

Oligonucleotide 2:

Sst I 5'-ATA GAG CTC TGT GGT GCC TGA GTC CTC 55 AG-3' (SEQ ID NO: 26)

This cDNA fragment was ligated in the pCD4-Hy3 vector [DSM 5523; European Patent Application No. 90107393.2; Japanese Patent Application No. 108967/90; U.S. Pat. No. 51,077,390] from which the CD4-cDNA had been removed via the SstI restriction cleavage sites. SstI cleavage sites are situated in vector pCD4-H γ 3 not only in front of, but also behind the CD4-partial sequence fragment. The construction was transfixed in J558 myeloma cells (ATCC No. TIB6) by means of protoplast fusion according to Oi et al. (Procd. Natl. Acad. Sci. USA 80, 825-829, 1983). Transfectants were selected by adding 5 µg/ml of mycophenolic acid and 250 g/ml of xanthin (Traunecker et al., Eur. J. Immunol. 16, 851-

55

65

854 [1986]) in basic medium (Dulbecco's modified Eagle's Medium, 10% foetal calf serum, 5×10⁻⁵M 2-mercaptoethanol). The expression product secreted by the transfixed cells could be purified using usual methods of protein chemistry, e.g. TNF-BP-antibody affinity chromatography. Unless not 5 already specifically indicated, standard procedures as described e.g. by Freshney, R. I. in "Culture of Animal Cells", Alan R. Liss, Inc., New York (1983) were used for the cultivation of the cell lines employed, for the cloning, for the selection or for the expansion of the cloned cells.

REFERENCES

- 1. G. E. Nedwin, S. L. Naylor, A. Y. Sakaguchi, D. Smith, Jarrett-Nedwin, D. Pennica, D. V. Goeddel and P. W. Gray: 15 Nucl. Acids Res. 12, 6361, 1985
- 2. B. Beutler and A. Cerami: New England J. Med. 316, 379, 1987
- 3. L. J. Old: Science 230, 630, 1985
- 4. G. Trinchieri, M. Kobavashi, M. Rosen, R. Loudon, M. 20 Murphy and B. Perussia: J. exp. Med. 164, 1206, 1986
- 5. J. Vilcek, V. J. Palombella, D. Henriksen-de Stefano, C. Swenson, R. Feinman, M. Hirai and M. Tsujimoto: J. exp. Med. 163, 632, 1986
- 6. B. J. Sugarman, B. B. Aggarwal, P. E. Hass, I. S. Figari, M. 25 A. Palladino and H. M. Shepard: Science 230, 943, 1985
- 7. J. R. Gamble, J. M. Harlan, S. J. Klebanoff and M. A. Vadas: Proc. Natl. Acad. Sci. USA 82, 8667, 1985
- 8. N. Sato, T. Goto, K. Haranaka, N. Satomi, H. Nariuchi, Y. Mano and Y. Sawasaki: J. Natl. Cancer Inst. 76, 1113, 1986 30
- 9. A. H. Stolpen, E. C. Guinan, W. Fiers and J. S. Pober: Am. J. Pathol. 123, 16, 1986
- 10. J. S. Pober, L. A. Lapierre, A. H. Stolpen, T. A. Brock, T. A. Springer, W. Fiers, M. P. Bevilacqua, D. L. Mendrick and M. A Gimbrone: J. Immunol. 138, 3319, 1987
- 11. M. Kawakami, P. Pekala, M. Lane and A. Cerami: Proc. Natl. Acad. Sci. USA 22, 912, 1982
- 12. T. Collins, L. A. Lapierre, W. Fiers, J. L. Strominger and J. S Pober: Proc. Natl. Acad. Sci. USA 83, 446, 1986
- 13. G. H. W. Wong and D. V. Goeddel: Nature 323, 819, 1986 40
- 14. J. W. Lowenthal, D. W. Ballard, E. B•hnlein and W. C. Greene: Proc. Natl. Acad. Sci. USA ja, 2331, 1989
- 15. M. J. Lenardo, C. M. Fan, T. Maniatis and D. Baltimore: Cell 51, 287, 1989
- 16. A. E. Goldfeld and T. Maniatis: Proc. Natl. Acad. Sci. 45 44. E. Harlow and D. Lane: Antibodies, A Laboratory USA 86, 1490, 1989
- 17. A. Waage, A. Halsteuren and T. Espevik: Lancet, Febr. 14, 1987, 355,
- 18. C. O. Jacob and H. O. McDevitt: Nature 331, 356, 1988
- 19. G. E. Grau, L. F. Fajardo, P. Piguet, B. Allet, P. Lambert 50 46. D. Pennica and D. V. Goeddel, in Lymphokines, Vol. 13, and P. Vassalli: Science 237, 1210, 1987
- 20. B. Beutler, I. W. Milsark and A. C. Cerami: Science 229, 869, 1985
- 21. B. B. Aggarwal, T. E. Eessalu and P. E. Hass: Nature 318, 665, 1985
- 22. M. Tsujimoto, Y. K. Yip and J. Vilcek: Proc. Natl. Acad. Sci. USA 82, 7626, 1985
- 23. C. Baglioni, S. McCandless, J. Tavernier and W. Fiers: J. Biol. Chem. 260, 13395, 1985
- 24. P. Hohmann, R. Remy, M. Brockhaus and A. P. G. M. van 60 Loon: J. Biol. Chem., im Druck
- 25. F. C. Kull, S. Jacobs and P. Cuatrecasas: Proc. Natl. Acad. Sci. USA 82, 5756, 1985
- 26. A. A. Creasy, R. Yamamoto and Ch. R. Vitt: Proc. Natl. Acad. Sci. USA 84, 3293, 1987
- 27. G. B. Stauber, R. A. Aiyer and B. B. Aggarwal: J. Biol. Chem. 263, 19098, 1988

- 28. K. Hirano, K. Yamamoto, Y. Kobayashi and T. Osawa: J. Biochem. 105, 120, 1989
- 29. Y. Niitsu, N. Watanabe, H. Sone, H. Neda, N. Yamauchi, M. Maeda and I. Urushizaki: J. Biol. Resp. Modifiers 7, 276, 1988
- 30. I. Olsson, A. Grubb, U. Gullberg, M. Lantz, E. Nilsson, C. Peetre and H. Thysell: Abstract, 2nd Intern. Conference on Tumor Necrosis Factor and Related Cytokines, Napa, Calif., 15.-20. January 1989
- 10 31. H. R. Loetscher and M. Brockhaus: Abstract, 2nd Intern. Conference on Tumor Necrosis Factor and Related Cytokines, Napa, Calif., 15.-20. Januar 1989
 - 32. M. Brockhaus, H. Loetscher, H.-P. Hohmann and W. Hunziker: Abstract, 2nd Intern. Conference on Tumor Necrosis Factor and Related Cytokines, Napa, Calif., 15.-20. January 1989
 - 33. C. R. Cantor and P. R. Schimmel, in Biophysical Chemistry, W.H. Freeman, ed., San Francisco, 1980, p. 850
 - 34. M. W. Hunkapiller, E. Lujan, F. Ostrander, L. E. Hood: Methods Enzymol. 91, 227, 1983
 - 35. U. K. Lammli: Nature 227, 680, 1970
 - 36. T. St. John, W. M. Gallatin, M. Siegelman. H. T. Smith, V. A. Fried and I. L. Weissman: Science 231, 845, 1986
 - 37. M. Siegelman, M. W. Bond, W. M. Gallatin, T. St. John, H. T. Smith, V.A. Fried and I.L. Weissman: Science 231, 823, 1986
 - 38. H. Towbin, T. Staehelin and J. Gordon: Proc. Natl. Acad. Sci. USA 76, 4350, 1979
 - 39. Dinarello, Ch. A., in Lymphokines, Vol. 14, E. Pick, ed., p. 1, Academic Press, London, 1987
 - 40. D. J. Merchant, R. H. Kahn and W. H. Murphy: Handbook of Cell and Organ Culture, Burgess Publ. Co., Minneapolis, 1969
 - 41. G. E. Grau, T. E. Taylor, M. E. Molyneux, J. J. Wirima, P. Vassalli, M. Hommel and P. Lambert: New Engl. J. Med. 320. 1586. 1989
 - 42. J. Sambrook, E. F. Fritsch and T. Maniatis: Molecular Cloning, A Laboratory Manual, 2nd ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, Cold Spring Harbor Laboratory Press, 1989
 - 43. F. M. Ausubel, R. Brent, R. E. Kingston, D. D. Moore, J. A. Smith, J. G. Seidman and K. Struhl: Current Protocols in Molecular Biology 1987-1988, S. Wiley and Sons, New York, 1987
 - Manual, Cold Spring Harbor Laboratory Publications, New York, 1988
 - 45. S. Fazekas de St. Groth and D. Scheidegger: J. Immunol. Methods 35, 1, 1980
 - D. R. Webb and D. V. Goeddel, eds. p. 163, Academic Press, London, 1987
 - 47. J. Tavernier, L. Franzen, A. Marmenout, J. van der Heyden, R. Muller, M. Ruysschaert, A. van Vliet, R. Banden and W. Fiers, in Lymphokines, Vol. 13, D. R. Webb and D. V. Goeddel, eds., p. 181, Academic Press, London
 - 48. P. J. Fraker and J. C. Speck: Biochem. Biophys. Res. Commun. 80, 849, 1987
 - 49. D. H. Erlich, D. H. Gelfand, R. K. Saiki: Nature 331, 61, 1988
 - 50. Bosserhoff, J. Wallach and R. W. Frank: J. Chromatogr. 473, 71, 1989
 - 51. R. Lathe: J. Mol. Biol. 183, 1, 1985
 - 52. Luckow and Summers, "A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures", Texas Agricultural Experimental Station, Texas A & M University, Bulletin No. 1555, 2nd edition, 1988

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 26 <210> SEQ ID NO 1 <211> LENGTH: 2111 <212> TYPE: DNA <213> ORGANISM: Homo sapiens <400> SEQUENCE: 1

gaattcgggg	gggttcaaga	tcactgggac	caggccgtga	tctctatgcc	cgagtctcaa	60
ccctcaactg	tcaccccaag	gcacttggga	cgtcctggac	agaccgagtc	ccgggaagcc	120
ccagcactgc	cgctgccaca	ctgccctgag	cccaaatggg	ggagtgagag	gccatagctg	180
tctggcatgg	gcctctccac	cgtgcctgac	ctgctgctgc	cgctggtgct	cctggagctg	240
ttggtgggaa	tatacccctc	aggggttatt	ggactggtcc	ctcacctagg	ggacagggag	300
aagagagata	gtgtgtgtcc	ccaaggaaaa	tatatccacc	ctcaaaataa	ttcgatttgc	360
tgtaccaagt	gccacaaagg	aacctacttg	tacaatgact	gtccaggccc	ggggcaggat	420
acggactgca	gggagtgtga	gageggetee	ttcaccgctt	cagaaaacca	cctcagacac	480
tgcctcagct	gctccaaatg	ccgaaaggaa	atgggtcagg	tggagatctc	ttcttgcaca	540
gtggaccggg	acaccgtgtg	tggctgcagg	aagaaccagt	accggcatta	ttggagtgaa	600
aaccttttcc	agtgcttcaa	ttgcagcctc	tgcctcaatg	ggaccgtgca	cctctcctgc	660
caggagaaac	agaacaccgt	gtgcacctgc	catgcaggtt	tctttctaag	agaaaacgag	720
tgtgtctcct	gtagtaactg	taagaaaagc	ctggagtgca	cgaagttgtg	cctaccccag	780
attgagaatg	ttaagggcac	tgaggactca	ggcaccacag	tgctgttgcc	cctggtcatt	840
ttctttggtc	tttgcctttt	atccctcctc	ttcattggtt	taatgtatcg	ctaccaacgg	900
tggaagtcca	agctctactc	cattgtttgt	gggaaatcga	cacctgaaaa	agaggggggag	960
cttgaaggaa	ctactactaa	gcccctggcc	ccaaacccaa	gcttcagtcc	cactccaggc	1020
ttcaccccca	ccctgggctt	cagtcccgtg	cccagttcca	ccttcacctc	cagetecace	1080
tatacccccg	gtgactgtcc	caactttgcg	gctccccgca	gagaggtggc	accaccctat	1140
caggggggctg	accccatcct	tgcgacagcc	ctcgcctccg	accccatccc	caaccccctt	1200
cagaagtggg	aggacagcgc	ccacaagcca	cagagcctag	acactgatga	ccccgcgacg	1260
ctgtacgccg	tggtggagaa	cgtgcccccg	ttgcgctgga	aggaattcgt	gcggcgccta	1320
gggctgagcg	accacgagat	cgatcggctg	gagctgcaga	acgggcgctg	cctgcgcgag	1380
gcgcaataca	gcatgctggc	gacctggagg	cggcgcacgc	cgcggcgcga	ggccacgctg	1440
gagctgctgg	gacgcgtgct	ccgcgacatg	gacctgctgg	gctgcctgga	ggacatcgag	1500
gaggcgcttt	gcggccccgc	cgccctcccg	cccgcgccca	gtcttctcag	atgaggctgc	1560
gcccctgcgg	gcagctctaa	ggaccgtcct	gcgagatcgc	cttccaaccc	cacttttttc	1620
tggaaaggag	gggtcctgca	ggggcaagca	ggagctagca	gccgcctact	tggtgctaac	1680
ccctcgatgt	acatagcttt	tctcagctgc	ctgcgcgccg	ccgacagtca	gcgctgtgcg	1740
cgcggagaga	ggtgcgccgt	gggctcaaga	gcctgagtgg	gtggtttgcg	aggatgaggg	1800
acgctatgcc	tcatgcccgt	tttgggtgtc	ctcaccagca	aggetgeteg	ggggcccctg	1860
gttcgtccct	gagcettttt	cacagtgcat	aagcagtttt	ttttgttttt	gttttgtttt	1920
gttttgtttt	taaatcaatc	atgttacact	aatagaaact	tggcactcct	gtgccctctg	1980
cctggacaag	cacatagcaa	gctgaactgt	cctaaggcag	gggcgagcac	ggaacaatgg	2040

-continued

ggc	cttca	agc 1	tgga	gctg	tg ga	actt	ttgta	a ca	taca	ctaa	aati	cctga	aag t	taaa	aaaaaa	2100
aaco	ccgaa	att (C													211:
<21) <21; <21; <21;	0> SI L> LI 2> T 3> OF	EQ II ENGTI (PE : RGAN]	D NO H: 4 PRT ISM:	2 55 Home	o saj	pien	5									
<40	D> SI	EQUEI	NCE:	2												
Met 1	Gly	Leu	Ser	Thr 5	Val	Pro	Asp	Leu	Leu 10	Leu	Pro	Leu	Val	Leu 15	Leu	
Glu	Leu	Leu	Val 20	Gly	Ile	Tyr	Pro	Ser 25	Gly	Val	Ile	Gly	Leu 30	Val	Pro	
His	Leu	Gly 35	Asp	Arg	Glu	Lys	Arg 40	Asp	Ser	Val	Суз	Pro 45	Gln	Gly	Lys	
Tyr	Ile 50	His	Pro	Gln	Asn	Asn 55	Ser	Ile	Суз	Суз	Thr 60	ГЛа	Суз	His	Гла	
Gly 65	Thr	Tyr	Leu	Tyr	Asn 70	Asp	Суз	Pro	Gly	Pro 75	Gly	Gln	Asp	Thr	Aap 80	
Сув	Arg	Glu	Сув	Glu 85	Ser	Gly	Ser	Phe	Thr 90	Ala	Ser	Glu	Asn	His 95	Leu	
Arg	His	Сув	Leu 100	Ser	Сүз	Ser	Lys	Cys 105	Arg	Lys	Glu	Met	Gly 110	Gln	Val	
Glu	Ile	Ser 115	Ser	Суз	Thr	Val	Asp 120	Arg	Asp	Thr	Val	Cys 125	Gly	Суз	Arg	
ГЛа	Asn 130	Gln	Tyr	Arg	His	Tyr 135	Trp	Ser	Glu	Asn	Leu 140	Phe	Gln	Сув	Phe	
Asn 145	Сув	Ser	Leu	Суа	Leu 150	Asn	Gly	Thr	Val	His 155	Leu	Ser	Сув	Gln	Glu 160	
ГЛа	Gln	Asn	Thr	Val 165	Суз	Thr	Суз	His	Ala 170	Gly	Phe	Phe	Leu	Arg 175	Glu	
Asn	Glu	Суз	Val 180	Ser	Суз	Ser	Asn	Cys 185	Lys	Lys	Ser	Leu	Glu 190	Суз	Thr	
ГЛа	Leu	Cys 195	Leu	Pro	Gln	Ile	Glu 200	Asn	Val	Lys	Gly	Thr 205	Glu	Asp	Ser	
Gly	Thr 210	Thr	Val	Leu	Leu	Pro 215	Leu	Val	Ile	Phe	Phe 220	Gly	Leu	Суз	Leu	
Leu 225	Ser	Leu	Leu	Phe	Ile 230	Gly	Leu	Met	Tyr	Arg 235	Tyr	Gln	Arg	Trp	Lys 240	
Ser	Lys	Leu	Tyr	Ser 245	Ile	Val	Сүз	Gly	Lys 250	Ser	Thr	Pro	Glu	Lys 255	Glu	
Gly	Glu	Leu	Glu 260	Gly	Thr	Thr	Thr	Lys 265	Pro	Leu	Ala	Pro	Asn 270	Pro	Ser	
Phe	Ser	Pro 275	Thr	Pro	Gly	Phe	Thr 280	Pro	Thr	Leu	Gly	Phe 285	Ser	Pro	Val	
Pro	Ser 290	Ser	Thr	Phe	Thr	Ser 295	Ser	Ser	Thr	Tyr	Thr 300	Pro	Gly	Asp	Суа	
Pro 305	Asn	Phe	Ala	Ala	Pro 310	Arg	Arg	Glu	Val	Ala 315	Pro	Pro	Tyr	Gln	Gly 320	
Ala	Asp	Pro	Ile	Leu 325	Ala	Thr	Ala	Leu	Ala 330	Ser	Aap	Pro	Ile	Pro 335	Asn	
Pro	Leu	Gln	Lys 340	Trp	Glu	Asp	Ser	Ala 345	His	Lys	Pro	Gln	Ser 350	Leu	Asp	
Thr	Asp	Asp 355	Pro	Ala	Thr	Leu	Tyr 360	Ala	Val	Val	Glu	Asn 365	Val	Pro	Pro	

Leu Arg Trj 370	p Lys Glu Pl	he Val Arg 375	Arg Leu Gly	Leu Ser Asp 380) His Glu	
Ile Asp Are 385	g Leu Glu Lo 31	eu Gln Asn 90	Gly Arg Cys 395	Leu Arg Glu	ı Ala Gln 400	
Tyr Ser Met	t Leu Ala Ti 405	hr Trp Arg	Arg Arg Thr 410	Pro Arg Arg	g Glu Ala 415	
Thr Leu Glu	ı Leu Leu G 420	ly Arg Val	Leu Arg Asp 425	Met Asp Leu 430	ı Leu Gly	
Cys Leu Glu 43!	ı Asp Ile G 5	lu Glu Ala 440	Leu Cys Gly	Pro Ala Ala 445	a Leu Pro	
Pro Ala Pro 450	o Ser Leu L	eu Arg 455				
<210> SEQ : <211> LENG' <212> TYPE <213> ORGAI	ID NO 3 TH: 2339 : DNA NISM: Homo ;	sapiens				
<400> SEQUI	ENCE: 3					
tcggacaccg	tgtgtgactc	ctgtgaggac	agcacataca	cccagctctg	gaactgggtt	60
cccgagtgct	tgagctgtgg	ctcccgctgt	agctctgacc	aggtggaaac	tcaagcctgc	120
actcgggaac	agaaccgcat	ctgcacctgc	aggeeegget	ggtactgcgc	gctgagcaag	180
caggaggggt	gccggctgtg	cgcgccgctg	ccgaagtgcc	gcccgggctt	cggcgtggcc	240
agaccaggaa	ctgaaacatc	agacgtggtg	tgcaagccct	gtgccccggg	gacgttctcc	300
aacacgactt	catccacgga	tatttgcage	ccccaccaga	tctgtaacgt	ggtggccatc	360
cctgggaatg	caagcaggga	tgcagtctgc	acgtccacgt	cccccacccg	gagtatggcc	420
ccaggggcag	tacacttacc	ccagccagto	tccacacgat	cccaacacac	gcagccaagt	480
ccagaaccca	gcactgctcc	aagcacctco	tteetgetee	caatgggccc	cagcccccca	540
gctgaaggga	gcactggcga	cttcgctctt	ccagttggac	tgattgtggg	tgtgacagcc	600
ttgggtctac	taataatagg	agtggtgaad	tgtgtcatca	tgacccaggt	gaaaaagaag	660
cccttgtgcc	tgcagagaga	agccaaggto	cctcacttgc	ctgccgataa	ggcccggggt	720
acacagggcc	ccgagcagca	gcacctgctg	atcacagcgc	cgagctccag	cagcagctcc	780
ctggagagct	cggccagtgc	gttggacaga	agggcgccca	ctcggaacca	gccacaggca	840
ccaggcgtgg	aggccagtgg	ggccggggag	gcccgggcca	gcaccgggag	ctcagcagat	900
tetteecetg	gtggccatgg	gacccaggto	aatgtcacct	gcatcgtgaa	cgtctgtagc	960
agetetgace	acageteaca	gtgeteetee	caagccagct	ccacaatggg	agacacagat	1020
tccagcccct	cggagtcccc	gaaggacgac	caggtcccct	tctccaagga	ggaatgtgcc	1080
tttcqqtcac	aqctqqaqac	qccaqaqaco	ctqctqqqqa	qcaccqaaqa	qaaqcccctq	1140
ccccttqqaq	tqcctqatqc	tqqqatqaaq	cccaqttaac	caqqccqqtq	tqqqctqtqt	1200
cqtaqccaaq	gtggctgagc	cctggcagga	tqaccctqcq	aaqqqqccct	ggtccttcca	1260
qqcccccacc	actaggactc	tgaggetett	tctqqqccaa	gttcctctag	taccetecae	1320
agccgcagcc	tecetetgae	ctgcaggcca	agagcagagg	cagcgagttg	tggaaagcet	1380
ctqctqccat	qqcatataca	teteggaage	l ctgactaaaa	atggacgtto	gggggatget	1440
aaaaaaata	cctgagtete	tataadatad	eccaccasaa	tacacetace	agectgactt	1500
atagagaget	tagatttt	atttatta	ttatttatt	atttattat	agaactagaa	1560
tataa	tataatt		anget	5000900000	anatht -t	1.000
LCTGCCCAGC	LCTGGCTTCC	agaaaacccc	agcateett	LCCGCagagg	yycttctgg	16ZU

US 8,063,182 B1

-continued

agaggaggga tgctgcctga gtcacccatg aagacaggac agtgcttcag cctgaggctg agactgcggg atggtcctgg ggctctgtgc agggaggagg tggcagccct gtagggaacg gggtccttca agttagctca ggaggcttgg aaagcatcac ctcaggccag gtgcagtggc tcacgcctat gatcccagca ctttgggagg ctgaggcggg tggatcacct gaggttagga qttcqaqacc aqcctqqcca acatqqtaaa accccatctc tactaaaaat acaqaaatta gccqqqcqtq qtqqcqqqca cctataqtcc caqctactca qaaqcctqaq qctqqqaaat cgtttgaacc cgggaagcgg aggttgcagg gagccgagat cacgccactg cactccagcc tgggcgacag agcgagagtc tgtctcaaaa gaaaaaaaaa aagcaccgcc tccaaatgct aacttqtcct tttqtaccat qqtqtqaaaq tcaqatqccc aqaqqqccca qqcaqqccac catattcaqt qctqtqqcct qqqcaaqata acqcacttct aactaqaaat ctqccaattt tttaaaaaag taagtaccac tcaggccaac aagccaacga caaagccaaa ctctgccagc cacatecaae cececaeetq ceattigeae cetecgeett caeteegqtg tgeetgeag <210> SEQ ID NO 4 <211> LENGTH: 392 <212> TYPE: PRT <213> ORGANISM: Homo sapiens <400> SEQUENCE: 4 Ser Asp Thr Val Cys Asp Ser Cys Glu Asp Ser Thr Tyr Thr Gln Leu Trp Asn Trp Val Pro Glu Cys Leu Ser Cys Gly Ser Arg Cys Ser Ser Asp Gln Val Glu Thr Gln Ala Cys Thr Arg Glu Gln Asn Arg Ile Cys Thr Cys Arg Pro Gly Trp Tyr Cys Ala Leu Ser Lys Gln Glu Gly Cys Arg Leu Cys Ala Pro Leu Pro Lys Cys Arg Pro Gly Phe Gly Val Ala Arg Pro Gly Thr Glu Thr Ser Asp Val Val Cys Lys Pro Cys Ala Pro Gly Thr Phe Ser Asn Thr Thr Ser Ser Thr Asp Ile Cys Arg Pro His Gln Ile Cys Asn Val Val Ala Ile Pro Gly Asn Ala Ser Arg Asp Ala Val Cys Thr Ser Thr Ser Pro Thr Arg Ser Met Ala Pro Gly Ala Val His Leu Pro Gln Pro Val Ser Thr Arg Ser Gln His Thr Gln Pro Ser Pro Glu Pro Ser Thr Ala Pro Ser Thr Ser Phe Leu Leu Pro Met Gly Pro Ser Pro Pro Ala Glu Gly Ser Thr Gly Asp Phe Ala Leu Pro Val Gly Leu Ile Val Gly Val Thr Ala Leu Gly Leu Leu Ile Ile Gly Val Val Asn Cys Val Ile Met Thr Gln Val Lys Lys Pro Leu Cys Leu Gln Arg Glu Ala Lys Val Pro His Leu Pro Ala Asp Lys Ala Arg Gly Thr Gln Gly Pro Glu Gln Gln His Leu Leu Ile Thr Ala Pro Ser Ser

^{245 250 250 250}

Ser Ser Ser Ser Leu Glu Ser Ser Ala Ser Ala Leu Asp Arg Arg Ala 265 260 270 Pro Thr Arg Asn Gln Pro Gln Ala Pro Gly Val Glu Ala Ser Gly Ala 275 280 285 Gly Glu Ala Arg Ala Ser Thr Gly Ser Ser Ala Asp Ser Ser Pro Gly 290 295 300 Gly His Gly Thr Gln Val Asn Val Thr Cys Ile Val Asn Val Cys Ser 320 305 310 315 Ser Ser Asp His Ser Ser Gln Cys Ser Ser Gln Ala Ser Ser Thr Met 325 330 335 Gly Asp Thr Asp Ser Ser Pro Ser Glu Ser Pro Lys Asp Glu Gln Val 345 340 350 Pro Phe Ser Lys Glu Glu Cys Ala Phe Arg Ser Gln Leu Glu Thr Pro 355 360 365 Glu Thr Leu Leu Gly Ser Thr Glu Glu Lys Pro Leu Pro Leu Gly Val 370 375 380 Pro Asp Ala Gly Met Lys Pro Ser 385 390 <210> SEQ ID NO 5 <211> LENGTH: 28 <212> TYPE: PRT <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic peptide <220> FEATURE: <221> NAME/KEY: misc_feature <222> LOCATION: (25)..(25) <223> OTHER INFORMATION: Xaa = unknown amino acid <400> SEQUENCE: 5 Leu Val Pro His Leu Gly Asp Arg Glu Lys Arg Asp Ser Val Cys Pro 5 1 10 15 Gln Gly Lys Tyr Ile His Pro Gln Xaa Asn Ser Ile 20 25 <210> SEQ ID NO 6 <211> LENGTH: 15 <212> TYPE: PRT <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic peptide <400> SEQUENCE: 6 Ser Thr Pro Glu Lys Glu Gly Glu Leu Glu Gly Thr Thr Thr Lys 5 10 1 15 <210> SEQ ID NO 7 <211> LENGTH: 18 <212> TYPE: PRT <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic peptide <400> SEQUENCE: 7 Ser Gln Leu Glu Thr Pro Glu Thr Leu Leu Gly Ser Thr Glu Glu Lys 1 5 10 15 Pro Leu <210> SEQ ID NO 8 <211> LENGTH: 4 <212> TYPE: PRT

<213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic peptide <400> SEQUENCE: 8 Val Phe Cys Thr 1 <210> SEQ ID NO 9 <211> LENGTH: 16 <212> TYPE: PRT <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic peptide <400> SEQUENCE: 9 Asn Gln Pro Gln Ala Pro Gly Val Glu Ala Ser Gly Ala Gly Glu Ala 5 10 1 15 <210> SEQ ID NO 10 <211> LENGTH: 18 <212> TYPE: PRT <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic peptide <220> FEATURE: <221> NAME/KEY: misc_feature <222> LOCATION: (8)..(8) <223> OTHER INFORMATION: Xaa = unknown amino acid <400> SEQUENCE: 10 Leu Pro Ala Gln Val Ala Phe Xaa Pro Tyr Ala Pro Glu Pro Gly Ser 5 10 1 15 Thr Cys <210> SEQ ID NO 11 <211> LENGTH: 13 <212> TYPE: PRT <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic peptide <220> FEATURE: <221> NAME/KEY: misc_feature <222> LOCATION: (2)..(2) <223> OTHER INFORMATION: Xaa = unknown amino acid <400> SEQUENCE: 11 Ile Xaa Pro Gly Phe Gly Val Ala Tyr Pro Ala Leu Glu 10 1 5 <210> SEQ ID NO 12 <211> LENGTH: 4 <212> TYPE: PRT <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic peptide <400> SEQUENCE: 12 Leu Cys Ala Pro 1 <210> SEQ ID NO 13 <211> LENGTH: 7 <212> TYPE: PRT <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic peptide

```
<400> SEQUENCE: 13
```

Val Pro His Leu Pro Ala Asp 1 5

<210> SEQ ID NO 14 <211> LENGTH: 15 <212> TYPE: PRT <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic peptide <220> FEATURE: <221> NAME/KEY: misc_feature <222> LOCATION: (9)..(10) <223> OTHER INFORMATION: Xaa = unknown amino acid <220> FEATURE: <221> NAME/KEY: misc_feature <222> LOCATION: (13)..(13) <223> OTHER INFORMATION: Xaa = unknown amino acid <400> SEQUENCE: 14 Gly Ser Gln Gly Pro Glu Gln Gln Xaa Xaa Leu Ile Xaa Ala Pro 1 5 10 15 <210> SEQ ID NO 15 <211> LENGTH: 9 <212> TYPE: PRT
<213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic peptide <400> SEQUENCE: 15 Leu Val Pro His Leu Gly Asp Arg Glu 1 5 <210> SEQ ID NO 16 <211> LENGTH: 27 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic primer <400> SEQUENCE: 16 agggagaaga gagatagtgt gtgtccc <210> SEQ ID NO 17 <211> LENGTH: 41 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic primer <400> SEQUENCE: 17 aagettggee aggatecage tgactgaetg ategegagat e <210> SEO ID NO 18 <211> LENGTH: 41 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Antisense primer <400> SEQUENCE: 18 gatetegega teagteagte agetggatee tggeeaaget t <210> SEQ ID NO 19 <211> LENGTH: 38 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE:

27

41

41

-continued	
<223> OTHER INFORMATION: Synthetic primer	
<400> SEQUENCE: 19	
cacagggatc catagctgtc tggcatgggc ctctccac	38
<210> SEQ ID NO 20 <211> LENGTH: 44 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Antisense primer	
<400> SEQUENCE: 20	
cccggtacca gatetetatt atgtggtgee tgagteetea gtge	44
<210> SEQ ID NO 21 <211> LENGTH: 19 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic primer	
<400> SEQUENCE: 21	
gatccagaat tcataatag	19
<210> SEQ ID NO 22 <211> LENGTH: 19 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Antisense primer	
<400> SEQUENCE: 22	
gtacctatta tgaattetg	19
<210> SEQ ID NO 23 <211> LENGTH: 31 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic primer	
<400> SEQUENCE: 23	
gcaccacata atagagatet ggtaceggga a	31
<210> SEQ ID NO 24 <211> LENGTH: 25 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Antisense primer	
<400> SEQUENCE: 24	
cccggtacca gatctctatt atgtg	25
<210> SEQ ID NO 25 <211> LENGTH: 29 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic primer	
<400> SEQUENCE: 25	
tacgageteg gecatagetg tetggeatg	29

		-
cont	- 1 m	11ed
COILC		ucu

	a	
<211> LENGTH: 29		
<212> TYPE: DNA		
<213> ORGANISM: Artificial sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic primer		
<400> SEQUENCE: 26		
atagagetet gtggtgeetg agteeteag	29	

The invention claimed is:

1. A protein comprising

 (a) a human tumor necrosis factor (TNF)-binding soluble fragment of an insoluble human TNF receptor, wherein
 ¹⁵ the insoluble human TNF receptor (i) specifically binds human TNF, (ii) has an apparent molecular weight of about 75 kilodaltons on a non-reducing SDS-polyacrylamide gel, and (iii) comprises the amino acid sequence LPAQVAFXPYAPEPGSTC (SEQ ID NO: 10); and

(b) all of the domains of the constant region of a human immunoglobulin IgG heavy chain other than the first domain of said constant region;

wherein said protein specifically binds human TNF. 25

2. The protein of claim **1**, wherein the soluble fragment comprises the peptides LCAP (SEQ ID NO: 12) and VFCT (SEQ ID NO: 8).

3. The protein of claim **2**, wherein the soluble fragment further comprises the peptide LPAQVAFXPYAPEPGSTC ³⁰ (SEQ ID NO: 10).

4. The protein of claim **1**, wherein said human immunoglobulin IgG heavy chain is IgG_1 .

5. The protein of claim 4, wherein said domains of the constant region of the human immunoglobulin heavy chain ³⁵ consist essentially of the immunoglobulin amino acid sequence encoded by pCD4-H γ 1 vector (deposited at Deutschen Sammlung von Mikroorganismen und Zellkulturen GmbH (DSM) in Braunschweig, FRG under No. DSM 5314). ₄₀

6. A pharmaceutical composition comprising the protein of claim **4** and a pharmaceutically acceptable carrier material.

7. The protein of claim 1, wherein the protein is purified.

8. The protein of claim **1**, wherein the protein is produced by CHO cells.

9. The protein of claim **1**, wherein the protein consists of (a) the soluble fragment of the receptor and (b) all of the domains of the constant region of the human immunoglobulin IgG heavy chain other than the first domain of the constant region.

10. The protein of claim 1, wherein said domains of the 50 constant region of the human immunoglobulin heavy chain consist essentially of the immunoglobulin amino acid sequence encoded by pCD4-H γ 1 vector (deposited at Deutschen Sammlung von Mikroorganismen and Zellkulturen GmbH (DSM) in Braunschweig, FRG under No. DSM 5314) ⁵⁵ or by pCD4-H γ 3 vector (deposited at Deutschen Sammlung von Mikroorganismen und Zellkulturen GmbH (DSM) in Braunschweig, FRG under No. DSM 5523).

11. The protein of claim 1, wherein the protein consists essentially of the extracellular region of the insoluble human TNF receptor and all the domains of the constant region of a human IgG_1 immunoglobulin heavy chain other than the first domain of the constant region.

12. A pharmaceutical composition comprising the protein ₆₅ of claim **11** and a pharmaceutically acceptable carrier material.

13. A protein comprising

(a) a human tumor necrosis factor (TNF)-binding soluble fragment of an insoluble human TNF receptor, wherein the insoluble human TNF receptor (i) specifically binds human TNF, (ii) has an apparent molecular weight of about 75 kilodaltons on a non-reducing SDS-polyacrylamide gel, and (iii) comprises the amino acid sequences LPAQVAFXPYAPEPGSTC (SEQ ID NO: 10), LCAP (SEQ ID NO: 12), VFCT (SEQ ID NO: 8), NQPQAPGVEASGAGEA (SEQ ID NO: 9) and VPHL-PAD (SEQ ID NO: 13),

wherein the soluble fragment comprises the peptides LCAP (SEQ ID NO: 12) and VFCT (SEQ ID NO: 8); and

(b) all of the domains of the constant region of a human IgG₁ heavy chain other than the first domain of the constant region;

wherein said protein specifically binds human TNF.

14. The protein of claim 13, wherein the protein is purified. 15. The protein of claim 13, wherein the protein is produced by CHO cells.

16. The protein of claim 13, wherein the protein consists of (a) the soluble fragment of the receptor and (b) all of the domains of the constant region of the human IgG_1 heavy chain other than the first domain of the constant region.

17. The protein of claim 13, wherein the soluble fragment further comprises the peptide LPAQVAFXPYAPEPGSTC (SEQ ID NO: 10).

18. A protein encoded by a polynucleotide which comprises two nucleic acid subsequences,

- (a) one of said subsequences encoding a human TNFbinding soluble fragment of an insoluble human TNF receptor protein having an apparent molecular weight of about 75 kilodaltons on a non-reducing SDS-polyacrylamide gel, said soluble fragment comprising the amino acid sequence LPAQVAFXPYAPEPGSTC (SEQ ID NO: 10), and
- (b) the other of said subsequences encoding all of the domains of the constant region of the heavy chain of a human IgG immunoglobulin other than the first domain of said constant region,

wherein said protein specifically binds human TNF.

19. The protein of claim **18**, wherein the soluble fragment comprises the peptides LCAP (SEQ ID NO: 12) and VFCT (SEQ ID NO: 8).

20. The protein of any one of claim **18** or **19**, wherein said human immunoglobulin heavy chain is IgG_1 .

21. The protein of claim 20, wherein said domains of the constant region of the human immunoglobulin heavy chain consist essentially of the immunoglobulin amino acid sequence encoded by the DNA insert of pCD4-H γ 1 vector (deposited at Deutschen Sammlung von Mikroorganismen und Zellkulturen GmbH (DSM) in Braunschweig, FRG under No. DSM 5314).

22. The protein of claim 18, wherein the protein is purified.23. The protein of claim 18, wherein the protein is produced by CHO cells.

24. The protein of claim **18**, wherein the protein consists of (a) the soluble fragment of the receptor and (b) all of the ⁵ domains of the constant region of the human IgG_1 heavy chain other than the first domain of the constant region.

25. The protein of claim 18, wherein said domains of the constant region of the human immunoglobulin heavy chain consist essentially of the immunoglobulin amino acid sequence encoded by pCD4-H γ 1 vector (deposited at Deutschen Sammlung von Mikroorganismen und Zellkulturen GmbH (DSM) in Braunschweig, FRG under No. DSM 5314) or by pCD4-H γ 3 vector (deposited at Deutschen Sammlung von Mikroorganismen und Zellkulturen GmbH (DSM) in Braunschweig, FRG under No. DSM 5514) in Braunschweig, FRG under No. DSM 5523).

26. A protein consisting of

- (a) a human tumor necrosis factor (TNF)-binding soluble fragment of an insoluble human TNF receptor, wherein the insoluble human TNF receptor (i) specifically binds human TNF, and (ii) has an apparent molecular weight of about 75 kilodaltons on a non-reducing SDS-polyacrylamide gel and (iii) comprises the amino acid sequence LPAQVAFXPYAPEPGSTC (SEQ ID NO: 10),
- wherein the soluble fragment comprises the peptides LCAP (SEQ ID NO:12) and VFCT (SEQ ID NO:8), and
- (b) all of the domains of the constant region of a human IgG_1 heavy chain other than the first domain of the constant region, 30
- wherein the protein specifically binds human TNF, and wherein the protein is produced by CHO cells.

27. The protein of claim **26**, wherein the soluble fragment comprises the peptide LPAQVAFXPYAPEPGSTC (SEQ ID NO: 10).

28. The protein of claim 26, wherein the protein is purified.29. A pharmaceutical composition comprising the protein of any of claim 1, 18, 26, or 27 and a pharmaceutically acceptable carrier material.

30. A protein comprising

- (a) human tumor necrosis factor (TNF) binding soluble fragment of the amino acid sequence encoded by the cDNA insert of the plasmid deposited with the ATCC on Oct. 17, 2006 under accession number PTA 7942,
- (b) all of the domains of the constant region of a human immunoglobulin IgG heavy chain other than the first domain of said constant region;

wherein said protein specifically binds human TNF.

31. The protein of claim **30**, consisting of the soluble fragment and all the domains of the constant region of the human immunoglobulin IgG heavy chain other than the first domain of said constant region.

32. The protein of claim **30**, wherein the protein is expressed by a mammalian host cell.

33. The protein of claim **32**, wherein the mammalian host cell is a CHO cell.

34. The protein of claim **32**, consisting of the soluble fragment and all the domains of the constant region of the human immunoglobulin IgG heavy chain other than the first domain of said constant region.

35. The protein of claim **30**, wherein the protein consists essentially of the extracellular region of the human tumor necrosis factor (TNF) receptor amino acid sequence encoded by the cDNA insert, and all the domains of the constant region of a human IgG_1 immunoglobulin heavy chain other than the first domain of the constant region.

36. A pharmaceutical composition comprising the protein of claim **35** and a pharmaceutically acceptable carrier material.

* * * * *