



US010857231B2

(12) **United States Patent**
Dix et al.

(10) **Patent No.:** **US 10,857,231 B2**
(45) **Date of Patent:** **Dec. 8, 2020**

(54) **FORMULATIONS OF VEG ANTAGONIST FUSION PROTEINS AND METHOD OF MANUFACTURING THEM**

(71) Applicant: **REGENERON PHARMACEUTICALS, INC.**,
Tarrytown, NY (US)

(72) Inventors: **Daniel B. Dix**, LaGrangeville, NY (US); **Kelly Frye**, Mendham, NJ (US); **Susan Kautz**, Albany, NY (US)

(73) Assignee: **REGENERON PHARMACEUTICALS, INC.**,
Tarrytown, NY (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **16/535,610**

(22) Filed: **Aug. 8, 2019**

(65) **Prior Publication Data**

US 2019/0388539 A1 Dec. 26, 2019

Related U.S. Application Data

(60) Continuation of application No. 15/692,893, filed on Aug. 31, 2017, now Pat. No. 10,406,226, which is a continuation of application No. 15/342,989, filed on Nov. 3, 2016, now abandoned, which is a continuation of application No. 15/064,343, filed on Mar. 8, 2016, now Pat. No. 9,511,140, which is a continuation of application No. 14/550,385, filed on Nov. 21, 2014, now Pat. No. 9,416,167, which is a continuation of application No. 13/909,745, filed on Jun. 4, 2013, now Pat. No. 8,921,316, which is a continuation of application No. 13/428,510, filed on Mar. 23, 2012, now Pat. No. 8,710,004, which is a continuation of application No. 13/343,214, filed on Jan. 4, 2012, now Pat. No. 8,404,638, which is a division of application No. 12/835,065, filed on Jul. 13, 2010, now Pat. No. 8,110,546, which is a continuation of application No. 11/387,256, filed on Mar. 22, 2006, now abandoned.

(60) Provisional application No. 60/665,125, filed on Mar. 25, 2005.

(51) **Int. Cl.**

A61K 39/395 (2006.01)
A61K 38/17 (2006.01)
A61K 9/00 (2006.01)
A61K 9/08 (2006.01)
A61K 31/4172 (2006.01)
A61K 31/7012 (2006.01)
A61K 47/02 (2006.01)
A61K 47/26 (2006.01)
A61K 38/16 (2006.01)
A61K 47/10 (2017.01)
A61K 47/18 (2017.01)
A61K 47/22 (2006.01)
C07K 14/71 (2006.01)
C07K 16/22 (2006.01)
A61K 47/12 (2006.01)
A61K 9/19 (2006.01)

(52) **U.S. Cl.**

CPC **A61K 39/39591** (2013.01); **A61K 9/0019** (2013.01); **A61K 9/08** (2013.01); **A61K 9/19** (2013.01); **A61K 31/4172** (2013.01); **A61K 31/7012** (2013.01); **A61K 38/16** (2013.01); **A61K 38/179** (2013.01); **A61K 47/02** (2013.01); **A61K 47/10** (2013.01); **A61K 47/12** (2013.01); **A61K 47/183** (2013.01); **A61K 47/22** (2013.01); **A61K 47/26** (2013.01); **C07K 14/71** (2013.01); **C07K 16/22** (2013.01); **C07K 2318/20** (2013.01); **C07K 2319/30** (2013.01)

(58) **Field of Classification Search**

None

See application file for complete search history.

(56)

References Cited

U.S. PATENT DOCUMENTS

5,399,670 A	3/1995	Bhattacharya et al.
5,763,401 A	6/1998	Nayar
5,851,999 A	12/1998	Ulrich et al.
6,011,003 A	1/2000	Charmock-jonee et al.
6,100,071 A	8/2000	Davis-Smyth et al.
6,267,958 B1	7/2001	Andya et al.
6,270,993 B1	8/2001	Shibuya et al.
6,875,432 B2	4/2005	Liu et al.
6,897,294 B2	5/2005	Davis -Smyth et al.
7,001,892 B1	2/2006	Chmielweski et al.
7,052,691 B2	5/2006	Sleeman et al.
7,060,268 B2	6/2006	Andya et al.
7,070,959 B1	7/2006	Papadopoulos et al.
7,087,411 B2	8/2006	Daly et al.
7,374,757 B2	5/2008	Papadopoulos et al.
8,084,234 B2	12/2011	Papadopoulos et al.
8,343,737 B2	1/2013	Papadopoulos et al.
8,647,842 B2 †	2/2014	Papadopoulos

(Continued)

FOREIGN PATENT DOCUMENTS

WO	WO1993000807 A1	1/1993
WO	WO1999013909 A1	3/1999

(Continued)

OTHER PUBLICATIONS

Amand et al., Controllability analysis of protein glycosylation in CHO cells, PLOS One, 9, e87943, 2014. (Year: 2014).

(Continued)

Primary Examiner — Elly-Gerald Stoica

(74) *Attorney, Agent, or Firm* — Thomas Triolo; Karl Bozicevic

(57)

ABSTRACT

Formulations of a vascular endothelial growth factor (VEGF)-specific fusion protein antagonist are provided including a pre-lyophilized formulation, a reconstituted lyophilized formulation, and a stable liquid formulation. Preferably, the fusion protein has the sequence of SEQ ID NO:4.

114 Claims, No Drawings

Specification includes a Sequence Listing.

(56)

References Cited**U.S. PATENT DOCUMENTS**

2002/0004478	A1	1/2002	Danko et al.
2003/0092606	A1	5/2003	L'Italien et al.
2003/0113316	A1	6/2003	Kaisheva et al.
2003/0202972	A1	10/2003	Andya et al.
2004/0197324	A1	10/2004	Liu et al.
2005/0032699	A1	2/2005	Holash et al.
2005/0276808	A1	12/2005	Cedarbaum
2005/0281831	A1	12/2005	Davis-Smyth et al.

FOREIGN PATENT DOCUMENTS

WO	WO199962536	A2	12/1999
WO	WO2000075319	A1	12/2000
WO	WO2002060489	A1	8/2002
WO	WO2004103159	A2	12/2004
WO	WO2004106378	A2	12/2004
WO	WO2005000895	A2	1/2005
WO	WO200502972	A1	3/2005
WO	WO2005072772	A1	8/2005
WO	WO2006047325	A1	5/2006
WO	WO 2006/088650		8/2006
WO	WO2007149334	A2	12/2007

OTHER PUBLICATIONS

Borys et al., Culture pH affects expression rates and glycosylation of recombinant mouse placental lactogen proteins by Chinese hamster ovary (CHO) cells, *Biotechnology*, 11, 720-724, 1993. (year: 1993).

Carpenter, J.F. (1997) Rational Design of Stable Lyophilized . . . , *Pharm. Res.* 14(8):969-975.

Daugherty et al., (2006) Formulation and delivery issues for monoclonal . . . , *Adv. Drug Delivery Rev.* 58:668-706.

Fraser et al. (2004) Single injection of Vascular Trap . . . , *J. Clin. Endocrin. & Metabol.* 90(2): 1114-1122.

Glade-Blender, J. et al., VEGF Blocking Therapy in the Treatment of Cancer, Expert Opinion on Biological Therapy. Ashley London GB vol. 3, No. 2 Apr. 2003, pp. 263-276.

Hanks Solution <http://www.lifetechnologies.com/us/en/home/technical-resources/media-formulation.152.html>.

Holash, Jocelyn et al., (2002) VEGF-Trap: AVEGF blocker with potent antitumor effects, *Proceedings of the National Academy of Sciences of the United States of America*, vol. 99 No. 17 Aug. 20, 2002, 11393-11398.

Katayama, et al., (2004) Retrospective statistical analysis of lyophilized . . . , *J. Pharm. Sci.* 93(10): 2609-2623.

Kim, Eugene S. et al., *Proceedings of the National Academy of Sciences of the United States of America*, vol. 99, No. 17, Aug. 20, 2002.

Lopez et al., "Comparative enhancer effects of Span 20 with Tween 20 and Azone of the in vitro percutaneous penetration of compounds with different lipophilicities" *International Journal of Pharmaceutics* (Jul. 1, 2000) 202(1-2):133-140.

Mi et al., (2002) Effects of polyethylene glycol molecular weight and concentration . . . , *PDS J. Pharm. Sci. Technol.*, 56:115-123.

Stewart, M.W. (2012) Clinical and Differential utility of VEGF Inhibitors . . . , *Clinical Ophthalmology*, 2012:6; 175-1186.

Wang et al., Glycoengineering of CHO cells to improve product quality. Paula Meleady (ed.), *Heterologous Protein Production in CHO Cells: Methods and Protocols*, *Methods in Molecular Biology*, vol. 1603, DOI 10.1007/978-1-4939-6972-2_2, © Springer Science+Business Media LLC 2017. (Year: 2017).

Wang, W. (1999) Instability, stabilization and formulation of liquid protein . . . , *Int'l J. Pharmaceutics* 185(2): 129-188.

Webb et al. (2002) A new mechanism for decreasing aggregation . . . , *J. Pharm. Sci.* 93(10): 2609-2623.

Randolph and Jones, *Surfactant-Protein Interactions in Rational Design of Stable Protein Formulations*, pp. 159-175, published 2002, Kluwer Academic/Plenum publishers, New York, NY.†

Kendrick et al., *Physical Stability of Proteins in Aqueous Solution in Rational Design of Stable Protein Formulations*, pp. 61-84 (relabeled as pp. 1-19 in provided version), published 2002, Kluwer Academic/Plenum publishers, New York, NY.†

† cited by third party

FORMULATIONS OF VEGF ANTAGONIST FUSION PROTEINS AND METHOD OF MANUFACTURING THEM

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a continuation of U.S. patent application Ser. No. 15/342,989, filed on Nov. 3, 2016, which is a continuation of U.S. patent application Ser. No. 15/064,343, filed on Mar. 8, 2016 and granted on Dec. 6, 2016 as U.S. Pat. No. 9,511,140, which is a continuation of U.S. patent application Ser. No. 14/550,385, filed on Nov. 21, 2014 and granted on Aug. 16, 2016 as U.S. Pat. No. 9,416,167, which is a continuation of U.S. patent application Ser. No. 13/909,745, filed on Jun. 4, 2013 and granted on Dec. 30, 2014 as U.S. Pat. No. 8,921,316, which is a continuation of U.S. patent application Ser. No. 13/428,510, filed on Mar. 23, 2012 and granted on Apr. 29, 2014 as U.S. Pat. No. 8,710,004, which is a continuation of U.S. patent application Ser. No. 13/343,214, filed on Jan. 4, 2012 and granted on Mar. 26, 2013 as U.S. Pat. No. 8,404,638, which is a division of U.S. patent application Ser. No. 12/835,065, filed on Jul. 13, 2010 and granted on Feb. 7, 2012 as U.S. Pat. No. 8,110,546, which is a continuation of U.S. patent application Ser. No. 11/387,256, filed on Mar. 22, 2006, which claims the benefit of priority under 35 USC § 119(e) of U.S. Provisional Application No. 60/665,125, filed on Mar. 25, 2005, all of which are herein specifically incorporated by reference in their entirety.

BACKGROUND OF THE INVENTION

Field of the Invention

The present invention is directed to pharmaceutical formulations comprising agents capable of inhibiting vascular endothelial growth factor (VEGF), and to methods for making and using such formulations. The invention includes pharmaceutical formulations having increased stability.

Statement of Related Art

Vascular endothelial growth factor (VEGF) expression is nearly ubiquitous in human cancer, consistent with its role as a key mediator of tumor neoangiogenesis. Blockade of VEGF function, by binding to the molecule or its VEGFR-2 receptor, inhibits growth of implanted tumor cells in multiple different xenograft models (see, for example, Gerber et al. (2000) *Cancer Res.* 60:6253-6258). A soluble VEGF-specific fusion protein antagonist, termed a "VEGF trap" has been described (Kim et al. (2002) *Proc. Natl. Acad. Sci. USA* 99:11399-404; Holash et al. (2002) *Proc. Natl. Acad. Sci. USA* 99:11393-8), which references are specifically incorporated by reference in their entirety.

Lyophilization (freeze drying under controlled conditions) is commonly used for long term storage of proteins. The lyophilized protein is substantially resistant to degradation, aggregation, oxidation, and other degenerative processes while in the freeze-dried state (see, for example, U.S. Pat. No. 6,436,897).

BRIEF SUMMARY OF THE INVENTION

Stable formulations of a VEGF-specific fusion protein antagonist are herein provided. The pharmaceutically acceptable formulations of the invention comprise the

VEGF "trap" antagonist with a pharmaceutically acceptable carrier. In specific embodiments, liquid and freeze-dried, or lyophilized formulations are provided.

In a first aspect, the invention features a stable liquid formulation of a VEGF-specific fusion protein antagonist, comprising a fusion protein comprising a receptor component consisting essentially of an immunoglobulin-like (Ig) domain 2 of a first VEGF receptor and Ig domain 3 of a second VEGF receptor, and a multimerizing component, one or more buffers, and one or more thermal stabilizers. In a specific embodiment of the VEGF-specific fusion protein antagonist, the first VEGF receptor is Flt1 and the second VEGF receptor is Flk1 or Flt4. In a more specific embodiment the fusion protein has the amino acid sequence of SEQ ID NO:2 or SEQ ID NO:4. In one embodiment, the buffer is a phosphate buffer and/or citrate. More preferably, the buffers are phosphate and citrate. In one embodiment, the thermal stabilizers are NaCl and/or sucrose. More preferably, the thermal stabilizers are both NaCl and sucrose.

In a specific embodiment, the stable liquid formulation of a VEGF-specific fusion protein antagonist comprises 1-10 mM phosphate buffer, 1-10 mM citrate, 25-150 mM NaCl, 5-30% sucrose, 10-50 mg/ml of the fusion protein, at a pH of about 6-6.5. In a more specific embodiment, the stable liquid formulation comprises 5 mM phosphate buffer, 5 mM citrate buffer, 100 mM NaCl, 20% sucrose, 25 mg/ml of the fusion protein, at a pH of about 6.0. Additionally, polysorbate may be present, for example 0.05-0.15% polysorbate 20. The stable liquid formulation of the VEGF-specific fusion protein antagonist of the invention exhibits little or no precipitation after storage of a 25 mg/ml VEGF formulation for about 6 months at -80° C. and little or no precipitation after storage for 6 months at 5° C.

In a second aspect, the invention features a high concentration stable liquid formulation of a VEGF antagonist comprising 1-50 mM histidine, 25-150 mM NaCl, 5-30% sucrose, 50-100 mg/ml of the fusion protein, at a pH of about 6-6.5, and either 0.1-0.5% polysorbate or 1-5% PEG. In a more specific embodiment, the high concentration stable liquid formulation comprises 10 mM histidine, 50 mM NaCl, 5-20% sucrose, 50-100 mg/ml of the fusion protein, at a pH of about 6.0-6.5, with either 0.1% polysorbate (e.g., polysorbate 20) or 3% PEG (e.g., PEG 3350). The high concentration stable liquid formulation of the VEGF-specific fusion protein antagonist of the invention exhibits less than about 3% degradation after 15 months of storage at 5° C. (75 or 100 mg/ml VEGF trap protein) or less than about 1.5% degradation after 24 months (50 mg/ml).

In a third aspect, the invention features a pre-lyophilized formulation of a vascular endothelial growth factor (VEGF)-specific fusion protein antagonist, comprising (i) fusion protein comprising a receptor component consisting essentially of an immunoglobulin-like (Ig) domain 2 of a first VEGF receptor and Ig domain 3 of a second VEGF receptor, and a multimerizing component, (ii) a buffer, (iii) an organic co-solvent or bulking agent, and (iv) one or more lyoprotectants. In various embodiments, the buffer is histidine, the organic co-solvent or bulking agent is PEG, and the lyoprotectant(s) is at least one of glycine and sucrose. In one embodiment, the pre-lyophilized formulation of the invention does not contain a preservative.

In one embodiment of the pre-lyophilized formulation of the invention, the formulation comprises 5-50 mM histidine, 0.1-3.0% PEG, 0.25-3.0% glycine, 0.5-6.0% sucrose, and 5-75 mg/ml of the fusion protein, at a pH of about 6.0-6.5. In any embodiment, the pre-lyophilized formulation may

3

further comprise up to 0.05 mM citrate and/or 0.003-0.005% polysorbate. The polysorbate present may be, for example, polysorbate 20.

In a more specific embodiment, the pre-lyophilized formulation comprises about 10 mM histidine, about 1.5% PEG 3350, about 0.75% glycine, about 2.5% sucrose, and about 12.5 to 75 mg/ml VEGF-specific fusion protein, at a pH of about 6.25. In specific embodiments, the fusion protein comprises the protein sequence of SEQ ID NO:4, present as a multimer, e.g., a dimer. In separate embodiments, the reconstituted formulation is 2 times the concentration of the pre-lyophilized formulation, e.g., a 20 mg fusion protein/ml pre-lyophilized formulation is reconstituted to a final formulation of 60 mg fusion protein/ml. Generally, the lyophilized formulation is reconstituted with sterile water suitable for injection. In one embodiment, the reconstitution liquid may be bacteriostatic water.

In a preferred embodiment, the pre-lyophilized formulation consists essentially of about 10 mM histidine, about 1.5% PEG 3350, about 0.75% glycine, about 2.5% sucrose, and about 50 mg/ml of the fusion protein having the sequence of SEQ ID NO:4 as a dimer, at a pH of about 6.25. Citrate (less than or equal to about 0.02 mM) and/or polysorbate (less than or equal to about 0.0005%) may be present. Optionally, the pre-lyophilized formulation does not contain a preservative, a phosphate buffer, and/or more than trace amounts of NaCl. In one embodiment, the pre-lyophilized formulation consists of about 10 mM histidine, about 1.5% PEG 3350, about 0.75% glycine, about 2.5% sucrose, and about 50 mg/ml of the VEGF trap protein (SEQ ID NO:4), pH 6.3, and upon reconstitution contains 20 mM histidine, 3% PEG, 1.5% glycine, about 5% sucrose, and about 100 mg/ml VEGF trap protein.

In a fourth aspect, the invention features a method of producing a lyophilized formulation of a VEGF-specific fusion protein antagonist, comprising subjecting the pre-lyophilized formulation of the invention to lyophilization to generate a lyophilized formulation. The lyophilized formulation may be lyophilized by any method known in the art for lyophilizing a liquid.

In a fifth related aspect, the invention features a method of producing a reconstituted lyophilized formulation of a VEGF-specific fusion protein antagonist, comprising reconstituting the lyophilized formulation of the invention to a reconstituted formulation. In one embodiment, the reconstituted formulation is twice the concentration of the pre-lyophilized formulation, e.g., the method of the invention comprises: (a) producing a pre-lyophilized formulation of a VEGF-specific fusion protein antagonist, (b) subjecting the pre-lyophilized formulation of step (a) to lyophilization; and (c) reconstituting the lyophilized formulation of step (b).

In specific embodiments of the method of producing a reconstituted lyophilized formulation, a pre-lyophilized solution is present in a vial as a 25 mg VEGF-specific fusion protein antagonist per ml solution of pre-lyophilized formulation, which is lyophilized and reconstituted to an 50 mg/ml solution. In another embodiment, a 30 mg/ml pre-lyophilized solution is lyophilized and reconstituted to a 60 mg/ml solution. In another embodiment, a 40 mg/ml pre-lyophilized solution is lyophilized and reconstituted to a 80 mg/ml solution. In another embodiment, a 12.5 mg/ml pre-lyophilized solution is lyophilized and reconstituted to a 25 mg/ml solution. In another embodiment, a 50 mg/ml pre-lyophilized solution is lyophilized and reconstituted to a 100 mg/ml solution. In another embodiment, a 75 mg/ml pre-lyophilized solution is lyophilized and reconstituted to a

4

150 mg/ml solution. Preferably, the reconstituted lyophilized formulation does not contain a preservative.

Other objects and advantages will become apparent from a review of the ensuing detailed description.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is not limited to particular methods, and experimental conditions described, as such methods and conditions may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting unless indicated, since the scope of the present invention will be limited only by the appended claims.

As used in this specification and the appended claims, the singular forms "a", "an", and "the" include plural references unless the context clearly dictates otherwise. Thus for example, references to "a method" include one or more methods, and/or steps of the type described herein and/or which will become apparent to those persons skilled in the art upon reading this disclosure.

Unless stated otherwise, all technical and scientific terms and phrases used herein have the same meaning as commonly understood by one of ordinary skill in the art to which the invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials are now described. All publications mentioned herein are incorporated herein by reference.

General Description

Safe handling and administration of formulations comprising proteins represent significant challenges to pharmaceutical formulators. Proteins possess unique chemical and physical properties that present stability problems: a variety of degradation pathways exist for proteins, implicating both chemical and physical instability. Chemical instability includes deamination, aggregation, clipping of the peptide backbone, and oxidation of methionine residues. Physical instability encompasses many phenomena, including, for example, aggregation.

Chemical and physical stability can be promoted by removing water from the protein. Lyophilization (freeze-drying under controlled conditions) is commonly used for long-term storage of proteins. The lyophilized protein is substantially resistant to degradation, aggregation, oxidation, and other degenerative processes while in the freeze-dried state. The lyophilized protein is normally reconstituted with water optionally containing a bacteriostatic preservative (e.g., benzyl alcohol) prior to administration.

Definitions

The term "carrier" includes a diluent, adjuvant, excipient, or vehicle with which a composition is administered. Carriers can include sterile liquids, such as, for example, water and oils, including oils of petroleum, animal, vegetable or synthetic origin, such as, for example, peanut oil, soybean oil, mineral oil, sesame oil and the like.

The term "excipient" includes a non-therapeutic agent added to a pharmaceutical composition to provide a desired consistency or stabilizing effect. Suitable pharmaceutical excipients include, for example, starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium

stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like.

The term "lyophilized" or "freeze-dried" includes a state of a substance that has been subjected to a drying procedure such as lyophilization, where at least 50% of moisture has been removed.

The phrase "bulking agent" includes a compound that is pharmaceutically acceptable and that adds bulk to a lyo cake. Generally, acceptable bulking agents known to the art include, for example, carbohydrates, including simple sugars such as dextrose, ribose, fructose and the like, alcohol sugars such as mannitol, inositol and sorbitol, disaccharides including trehalose, sucrose and lactose, naturally occurring polymers such as starch, dextrans, chitosan, hyaluronate, proteins (e.g., gelatin and serum albumin), glycogen, and synthetic monomers and polymers. In the formulations of the invention, PEG 3350 is an organic co-solvent which is used to stabilize the fusion protein when agitated, mixed, or handled, and as a bulking agent to help produce an acceptable bulk.

The term "lyoprotectant" includes a substance that may be added to a freeze-dried or lyophilized formulation to help maintain protein structure when freeze-dried or lyophilized.

A "preservative" includes a bacteriostatic, bacteriocidal, fungistatic or fungicidal compound that is generally added to formulations to retard or eliminate growth of bacteria or other contaminating microorganisms in the formulations. Preservatives include, for example, benzyl alcohol, phenol, benzalkonium chloride, m-cresol, thimerosal, chlorbutanol, methylparaben, propylparaben and the like. Other examples of pharmaceutically acceptable preservatives can be found in the USP.

VEGF Antagonists

An VEGF antagonist is a compound capable of blocking or inhibiting the biological action of vascular endothelial growth factor (VEGF), and includes fusion proteins capable of trapping VEGF. In a preferred embodiment, the VEGF antagonist is the fusion protein of SEQ ID NO:2 or 4; more preferably, SEQ ID NO:4. In specific embodiments, the VEGF antagonist is expressed in a mammalian cell line such as a CHO cell and may be modified posttranslationally. In a specific embodiment, the fusion protein comprises amino acids 27-457 of SEQ ID NO:4 and is glycosylated at Asn residues 62, 94, 149, 222 and 308.

The VEGF antagonist of the methods and formulations of the invention can be prepared by any suitable method known in the art, or that comes to be known. The VEGF antagonist is preferably substantially free of protein contaminants at the time it is used to prepare the pharmaceutically acceptable formulation. By "substantially free of protein contaminants" is meant, preferably, that at least 90% of the weight of protein of the VEGF-specific fusion protein antagonist preparation used for making a formulation is VEGF fusion protein antagonist protein, more preferably at least 95%, most preferably at least 99%. The fusion protein is preferably substantially free of aggregates. "Substantially free of aggregates" means that at least 90% of the weight of fusion protein is not present in an aggregate at the time the fusion protein is used to prepare the pharmaceutically effective formulation. The fusion protein of the methods and formulations of the invention may contain low or trace amounts of compounds as a results of the purification process, for example, low or trace amounts of citrate and/or polysorbate. In one embodiment of the pre-lyophilized formulation of the invention containing about 50 mg of fusion protein/ml, citrate may be present at a concentration of about 0.02 mM

and/or polysorbate may be present at a concentration of about 0.0005%. If the pre-lyophilized formulation is reconstituted after lyophilization to half of the original volume (e.g., 100 mg/ml of fusion protein), the resulting concentrations may be 0.04 mM citrate and/or 0.001% polysorbate.

Lyophilization and Lyophilized Formulations

In one aspect of the invention, a pharmaceutically acceptable formulation comprising a VEGF-specific fusion protein antagonist is provided, wherein the formulation is a freeze-dried or lyophilized formulation. Lyophilized formulations can be reconstituted into solutions, suspensions, emulsions, or any other suitable form for administration or use. Lyophilized formulations are typically first prepared as liquids, then frozen and lyophilized. The total liquid volume before lyophilization can be less, equal to, or more than, the final reconstituted volume of the lyophilized formulation. The lyophilization process is well known to those of ordinary skill in the art, and typically includes sublimation of water from a frozen formulation under controlled conditions.

Lyophilized formulations can be stored at a wide range of temperatures. Lyophilized formulations may be stored below 25° C., for example, refrigerated at 4° C., or at room temperature (e.g., approximately 25° C.). Preferably, lyophilized formulations are stored below about 25° C., more preferably, at about 4-20° C.; below about 4° C.; below about -20° C.; about -40° C.; about -70° C., or about -80° C.

Lyophilized formulations are typically reconstituted for use by addition of an aqueous solution to dissolve the lyophilized formulation. A wide variety of aqueous solutions can be used to reconstitute a lyophilized formulation. Preferably, lyophilized formulations are reconstituted using water. Lyophilized formulations are preferably reconstituted with a solution consisting essentially of water (e.g., USP WFI, or water for injection) or bacteriostatic water (e.g., USP WFI with 0.9% benzyl alcohol). However, solutions comprising buffers and/or excipients and/or one or more pharmaceutically acceptable carries can also be used.

Freeze-dried or lyophilized formulations are typically prepared from liquids, that is, from solutions, suspensions, emulsions, and the like. Thus, the liquid that is to undergo freeze-drying or lyophilization preferably comprises all components desired in a final reconstituted liquid formulation. As a result, when reconstituted, the freeze-dried or lyophilized formulation will render a desired liquid formulation upon reconstitution. A preferred liquid formulation used to generate a freeze-dried or lyophilized formulation comprises a VEGF-specific fusion protein antagonist in a pharmaceutically effective amount, a buffer, a stabilizer, and a bulking agent. Freeze-dried or lyophilized formulations preferably comprise histidine, since histidine, in comparison to phosphate, is more effective at stabilizing the fusion protein when the fusion protein is lyophilized. Organic co-solvents, such as PEG 3350, are used to stabilize the fusion protein when agitated, mixed, or handled. A lyoprotectant is preferably used in freeze-dried or lyophilized formulations. Lyoprotectants help to maintain the secondary structure of proteins when freeze-dried or lyophilized. Two preferred example lyoprotectants are glycine and sucrose, which are preferably used together.

Stable Liquid Formulations

In one aspect, the invention provides a stable pharmaceutically acceptable formulation comprising a VEGF-specific fusion protein antagonist, wherein the formulation is a liquid formulation. Preferably, the liquid formulation comprises a pharmaceutically effective amount of the fusion protein. The

7

formulation can also comprise one or more pharmaceutically acceptable carriers, buffers, bulking agents, stabilizers, preservatives, and/or excipients. An example of a pharmaceutically acceptable liquid formulation comprises a VEGF-specific fusion protein antagonist in a pharmaceutically effective amount, a buffer, a co-solvent, and one or more stabilizers.

A preferred liquid formulation comprises phosphate buffer, an organic co-solvent, and one or more thermal stabilizers to minimize formation of aggregates and low molecular weight products when stored, and about 10 mg/ml to about 50 mg/ml fusion protein, wherein the formulation is from about pH 6.0-6.5. A preferred liquid formulation comprises about 5 mM phosphate buffer, about 5 mM citrate, about 100 mM NaCl, about 25% sucrose, and about 1050 mg/ml fusion protein, wherein the formulation is at a pH of about 6.0; optionally polysorbate may be present (e.g., 0.1% polysorbate 20). Although either NaCl or sucrose can be used as a stabilizer, a combination of NaCl and sucrose has been established to stabilize the fusion protein more effectively than either individual stabilizer alone.

Stability is determined in a number of ways at specified time points, including determination of pH, visual inspection of color and appearance, determination of total protein content by methods known in the art, e.g., UV spectroscopy, SDS-PAGE, size-exclusion HPLC, bioassay determination of activity, isoelectric focusing, and isoaspartate quantification. In one example of a bioassay useful for determining VEGF antagonist activity, a BAF/3 VEGFR1/EPOR cell line is used to determine VEGF165 binding by the VEGF-specific fusion protein antagonist of the invention.

Formulations, whether liquid or freeze-dried and lyophilized, can be stored in an oxygen-deprived environment. Oxygen-deprived environments can be generated by storing the formulations under an inert gas such as, for example, argon, nitrogen, or helium.

EXAMPLES

Before the present methods are described, it is to be understood that this invention is not limited to particular methods, and experimental conditions described, as such methods and conditions may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting, since the scope of the present invention will be limited only to the appended claims.

As used in this specification and the appended claims, the singular forms "a", "an", and "the" include plural references unless the context clearly dictates otherwise. Thus for example, a reference to "a method" includes one or more methods, and/or steps of the type described herein and/or which will become apparent to those persons skilled in the art upon reading this disclosure and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials are now described.

Example 1. Stability of a 50 mg/ml Liquid Formulation of VEGF Trap

A liquid formulation containing 10 mM phosphate, 50 mM NaCl, 0.1% polysorbate 20, 20% sucrose, and 50 mg/ml

8

VEGF trap (SEQ ID NO:4), pH 6.25, was stored at 5° C. and samples tested at 3, 6, 9, 12, 18 and 24 months. Stability was determined by SE-HPLC. The results, shown in Table 1, show that 98.6% and 98.3% of VEGF trap protein remained intact (non-degraded) at 12 and 24 months, respectively. Turbidity was measured at OD405 nm; and percent recovered protein by size exclusion HPLC.

TABLE 1

Stability of 50 mg/ml VEGF Trap Protein When Stored at 5° C. (VGFT-SS065)					
Months	Visual Appearance	Turbidity	pH	% VEGF Trap Recovered	% VEGF Trap Native Configuration
0	Pass	0.00	6.2	100	99.0
3	Pass	0.00	6.2	102	98.8
6	Pass	0.01	6.2	103	98.7
9	Pass	0.01	6.3	102	98.2
12	Pass	0.01	6.3	106	98.6
18	Pass	0.00	6.3	103	98.4
24	Pass	0.00	6.2	93	98.3

A liquid formulation containing 10 mM phosphate, 50 mM NaCl, 3% PEG 3350, 20% sucrose, and 50 mg/ml VEGF trap (SEQ ID NO:4), pH 6.25, was stored at 5° C. and samples tested at 3, 6, 9, 12, 18 and 24 months. Stability results are shown in Table 2.

TABLE 2

Stability of 50 mg/ml VEGF Trap Protein When Stored at 5° C. (VGFT-SS065)					
Months	Visual Appearance	Turbidity	pH	% VEGF Trap Recovered	% VEGF Trap Native Configuration
0	Pass	0.00	6.2	100	99.0
3	Pass	0.00	6.2	100	98.8
6	Pass	0.01	6.3	103	98.5
9	Pass	0.00	6.3	103	98.3
12	Pass	0.01	6.3	110	98.3
18	Pass	0.00	6.3	113	98.0
24	Pass	0.01	6.2	90	97.8

Example 2. Stability of a 75 mg/ml Liquid Formulation of VEGF Trap

A liquid formulation containing 10 mM phosphate, 50 mM NaCl, 0.1% polysorbate 20, 20% sucrose, and 75 mg/ml VEGF trap (SEQ ID NO:4), pH 6.25, was stored at 5° C. and samples tested at 0, 1, 2.3, 3, 9, 12 and 15 months. Stability results are shown in Table 3.

TABLE 3

Stability of 75 mg/ml VEGF Trap Protein When Stored at 5° C. (VGFT-SS101)					
Months	Visual Appearance	Turbidity	pH	% VEGF Trap Recovered	% VEGF Trap Native Configuration
0	Pass	0.00	6.2	100	97.1
1	Pass	0.00	6.2	96	97.0
2.3	Pass	0.00	6.2	98	96.7
3	Pass	0.00	6.2	97	96.1
9	Pass	-0.01	6.0	101	96.0
12	Pass	0.00	6.3	110	94.5
15	Pass	0.00	6.3	92	95.6

A liquid formulation containing 10 mM phosphate, 50 mM NaCl, 3% PEG 3350, 20% sucrose, and 75 mg/ml VEGF trap (SEQ ID NO:4), pH 6.25, was stored at 5° C. and samples tested at 0, 1, 2.3, 3, 9, 12 and 15 months. Stability results are shown in Table 4.

TABLE 4

Stability of 75 mg/ml VEGF Trap Protein When Stored at 5° C. (VGFT-SS101)					
Months	Visual Appearance	Turbidity	pH	% VEGF Trap Recovered	% VEGF Trap Native Configuration
0	Pass	0.00	6.2	100	96.8
1	Pass	0.00	6.2	99	96.7
2.3	Pass	0.00	6.2	97	96.3
3	Pass	0.00	6.2	89	95.6
9	Pass	-0.01	6.2	98	95.4
12	Pass	-0.01	6.3	112	94.1
15	Pass	0.00	6.3	98	94.8

Example 3. Stability of a 100 mg/ml Liquid Formulation of VEGF Trap

A liquid formulation containing 10 mM phosphate, 50 mM NaCl, 0.1% polysorbate 20, 20% sucrose, and 100 mg/ml VEGF trap (SEQ ID NO:4), pH 6.25, was stored at 5° C. and samples tested at 0, 1, 2.3, 3, 9, 12 and 15 months. Stability results are shown in Table 5.

TABLE 5

Stability of 100 mg/ml VEGF Trap Protein Stored at 5° C. (VGFT-SS101)					
Months	Visual Appearance	Turbidity	pH	% VEGF Trap Recovered	% VEGF Trap Native Configuration
0	Pass	0.00	6.3	100	96.7
1	Pass	0.00	6.2	92	96.6
2.3	Pass	0.00	6.2	92	96.2
6	Pass	0.00	6.2	99	95.5
9	Pass	-0.01	6.2	92	95.5
12	Pass	-0.01	6.2	110	93.9
15	Pass	0.00	6.3	108	94.8

A liquid formulation containing 10 mM phosphate, 50 mM NaCl, 3% PEG 3350, 20% sucrose, and 100 mg/ml VEGF trap (SEQ ID NO:4), pH 6.25, was stored at 5° C. and samples tested at 0, 1, 2.3, 3, 9, 12 and 15 months. Stability results are shown in Table 6.

TABLE 6

Stability of 100 mg/ml VEGF Trap Protein Stored at 5° C. (VGFT-SS101)					
Months	Visual Appearance	Turbidity	pH	% VEGF Trap Recovered	% VEGF Trap Native Configuration
0	Pass	0.00	6.3	100	96.5
1	Pass	0.01	6.2	94	96.2
2.3	Pass	0.01	6.2	93	95.7
6	Pass	0.01	6.2	102	94.6
9	Pass	0.00	6.2	95	94.6
12	Pass	0.00	6.3	96	92.8
15	Pass	0.01	6.3	102	93.9

Example 4. Further Embodiments of Stable VEGF Trap Formulations

In one embodiment, the invention provides a stable liquid VEGF-binding fusion protein (VEGF trap) formulations comprising 5 mM phosphate, 5 mM citrate, 100 mM NaCl, 0.1% Polysorbate 20, 20% sucrose, 25 mg/ml VEGF trap protein, pH 6.0. This formulation can either be delivered subcutaneously or diluted and delivered by intravenous infusion. Due to the high osmolality of this formulation, it is diluted 3-fold to achieve an iso-osmolar solution for intravenous administration. Stability studies showed less than about 1% degradation was detected after 3 years of storage at 2-8° C.

In one embodiment, the invention features a lyophilized formulation which is preferably concentrated two-fold from the pre-lyophilized to the post-lyophilized formulation, e.g., 50 to 100 mg/ml; 75 to 150 mg/ml, or 100 to 200 mg/ml VEGF trap protein. In one specific embodiment, the pre-lyophilized formulation comprises 10 mM histidine, 1.5% PEG 3350, 0.75% glycine, 2.5% sucrose, 50 mg/ml VEGF trap protein, pH 6.3, and is reconstituted to a formulation comprising 20 mM histidine, 3% PEG 3350, 1.5% glycine, 5% sucrose, 100 mg/ml VEGF trap protein, pH 6.3. Stability studies showed no degradation of the VEGF trap was detected after 6 months of storage at 2-8° C.

In one embodiment of a liquid formulation, the formulation comprises 10 mM histidine, 50 mM NaCl, 5-20% sucrose, 50-100 mg/ml VEGF trap, and one of 0.1% polysorbate 20 or TY° PEG 3350. One advantage of this liquid formulation is that it provides a higher concentration of VEGF trap without requiring the manufacture of a lyophilized product. Thus, this formulation provides ease for subcutaneous delivery, for example, by allowing provision of a liquid pre-filled syringe at a concentration higher than that delivered by IV infusion. Also, this formulation could advantageously be used to provide lower infusion volumes and shorter infusion times. The amount of degradation determined by SE-HPLC following incubation at 5° C. for up to 15 or 24 months is summarized in Table 7.

TABLE 7

Stability of Liquid Formulation with 50-100 mg/ml VEGF Trap (VGFT-SS101)				
Incubation (months)	VEGF Trap (mg/ml)	% Polysorbate 20	% PEG 3350	% Degradation
24	50	0.1	—	0.7
24	50	—	3	1.3
15	75	0.1	—	1.5
15	75	—	3	2.0
15	100	0.1	—	1.9
15	100	—	3	2.6

Example 5. Stability and Activity of Lyophilized and Liquid

The stability of a reconstituted lyophilized formulation was determined over a 6 month period. The pre-lyophilized formulation contained 10 mM histidine, 1.5% PEG 3350, 2.5% sucrose, 0.75% glycine and 50 mg/ml VEGF trap protein. After lyophilization, the reconstituted formulation contained 20 mM histidine, 3% PEG 3350, 5% sucrose, 1.5% glycine, and 100 mg/ml VEGF trap protein (SEQ ID NO:4). The results are shown in Table 8. Activity was determined in a cell based bioassay which directly measures

11

the ability of the VEGF trap to inhibit the biological effects of human VEGF on a mouse Baf/3 VEGFR1/EpoR cell line. Therefore, this bioassay directly measures the biological activity of the protein. The results are expressed as percent relative potency (test sample IC₅₀/reference VEGF IC₅₀ standard×100). The binding affinity of VEGF to the VEGF trap is measured using a sensitive ELISA that specifically measures free VEGF in equilibrated mixtures containing VEGF and various concentrations of the VEGF trap. Results are expressed as percent relative binding (IC₅₀ of test sample/ICH, of reference×100). Measured pH ranged between 6.3-6.5. All solutions were visually clear. The concentration of VEGF trap recovered was determined with a UV spectrophotometer as mg/ml at A280 nm. The percent VEGF trap recovered in the native configuration (main peak purity) was determined with SE-HPLC.

TABLE 8

Stability of VEGF Trap Lyophilized Formulation Stored at 5° C. (VGT-RS475)				
Months	Bioassay	Binding Assay	% Recovered	% Native Configuration
0	120	126	97.9	98.7
1	117	74	97.9	98.6
1 + 24 hr	126	72	99.0	98.5
1 + 4 hr	94	81	101.5	98.2
3	101	98	98.1	98.6
3 + 24 hr	65	94	98.1	98.2
6 + 4 hr			96.9	98.7
6 + 24 hr			98.8	98.6

12

A formulation containing about 5 mM phosphate, 5 mM citrate, 100 mM NaCl, 0.1% polysorbate 20, 20% sucrose, and 25 mg/ml VEGF trap protein was tested for stability and activity over 36 months when stored at 5° C. The results are shown in Table 9. All samples were clear and colorless as determined by visual inspection. pH ranged from 6.0-6.1. *Binding assay results for two measurements (1 and 2 months) are expressed directly and not as a percent of the standard.

TABLE 9

Stability and Activity of Liquid Formulation (VGT-FS405)					
Months	% Native Configuration	Bioassay	Binding Assay	Protein Content mg/ml	
0	99.7	106	72	25.0	
1	99.9	119	4.4 μ M*	25.2	
2	99.6	102	5.4 μ M*	25.1	
3	99.6	97	88	25.1	
6	99.6	101	106	25.0	
9	99.4	89	126	25.4	
12	99.5	85	95	25.2	
18	99.4	99	81	25.5	
24	99.3	75	95	25.6	
36	98.8	109	79	25.6	

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 4

<210> SEQ ID NO 1

<211> LENGTH: 1453

<212> TYPE: DNA

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 1

```

aagcttgggc tgcaggctga tcgactctag aggatcgatc cccgggagag ctgcaattcg      60
caaccacccat ggtcagctac tgggacaccg gggctcctgt gtgcgcgctg ctcagctgtc      120
tgctttctcac aggatctagt tccggaggta gacctttcgt agagatgtac agtgaaatcc      180
ccgaaattat acacatgact gaaggaaggg agctcgtcat tccctgccgg gttacgtcac      240
ctaacatcac tgttacttta aaaaagtttc cacttgacac ttgtatccct gatggaaaac      300
gcataatctg ggacagtaga aaggggttca tcatatcaaa tgcaacgtac aaagaaatag      360
ggcttctgac ctgtgaagca acagtcaatg ggcatattgta taagacaaac tatctcacac      420
atcgacaaac caatacaatc atagatgtgg ttctgagtcc gtctcatgga attgaactat      480
ctgttgaggaga aaagcttgtc ttaaattgta cagcaagaac tgaactaaat gtggggattg      540
acttcaactg ggaataccct tcttcgaagc atcagcataa gaaacttgta aaccgagacc      600
taaaaaccca gtctgggagt gagatgaaga aatttttgag caccttaact atagatgggtg      660
taaccgggag tgaccaagga ttgtacacct gtgcagcatc cagtgggctg atgaccaaga      720
agaacagcac atttgtcagg gtccatgaaa agggcccggg cgacaaaact cacacatgcc      780

```


-continued

```

caccgtgccc agcacctgaa ctccctggggg gaccgtcagt cttcctcttc cccccaaaac 840
ccaaggacac cctcatgatc tcccggaacc ctgagggtcac atgcgtgggtg gtggacgtga 900
gccacgaaga ccctgaggtc aagttcaact ggtacgtgga cggcgtggag gtgcataatg 960
ccaagacaaa gccgcggggag gagcagtaca acagcacgta ccgtgtgggtc agcgtcctca 1020
ccgtcctgca ccaggactgg ctgaatggca aggagtacaa gtgcaagggtc tccaacaaag 1080
ccctcccagc ccccatcgag aaaaccatct ccaaagccaa agggcagccc cgagaaccac 1140
aggtgtacac cctgccccca tcccgggatg agctgaccaa gaaccagggtc agcctgacct 1200
gcctgggtcaa aggtcttctat cccagcgaca tcgccgtgga gtgggagagc aatgggcagc 1260
cggagaacaa ctacaagacc acgcctcccg tgctggactc cgacggctcc ttcttctct 1320
atagcaagct caccgtggac aagagcaggt ggcagcaggg gaacgtcttc tcatgtctcg 1380
tgatgcatga ggctctgcac aaccactaca cgcagaagag cctctccctg tctccgggta 1440
aatgagcggc cgc 1453

```

```

<210> SEQ ID NO 2
<211> LENGTH: 458
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic

```

```

<400> SEQUENCE: 2

```

```

Met Val Ser Tyr Trp Asp Thr Gly Val Leu Leu Cys Ala Leu Leu Ser
1           5           10          15
Cys Leu Leu Leu Thr Gly Ser Ser Ser Gly Gly Arg Pro Phe Val Glu
20          25          30
Met Tyr Ser Glu Ile Pro Glu Ile Ile His Met Thr Glu Gly Arg Glu
35          40          45
Leu Val Ile Pro Cys Arg Val Thr Ser Pro Asn Ile Thr Val Thr Leu
50          55          60
Lys Lys Phe Pro Leu Asp Thr Leu Ile Pro Asp Gly Lys Arg Ile Ile
65          70          75          80
Trp Asp Ser Arg Lys Gly Phe Ile Ile Ser Asn Ala Thr Tyr Lys Glu
85          90          95
Ile Gly Leu Leu Thr Cys Glu Ala Thr Val Asn Gly His Leu Tyr Lys
100         105         110
Thr Asn Tyr Leu Thr His Arg Gln Thr Asn Thr Ile Ile Asp Val Val
115         120         125
Leu Ser Pro Ser His Gly Ile Glu Leu Ser Val Gly Glu Lys Leu Val
130         135         140
Leu Asn Cys Thr Ala Arg Thr Glu Leu Asn Val Gly Ile Asp Phe Asn
145         150         155         160
Trp Glu Tyr Pro Ser Ser Lys His Gln His Lys Lys Leu Val Asn Arg
165         170         175
Asp Leu Lys Thr Gln Ser Gly Ser Glu Met Lys Lys Phe Leu Ser Thr
180         185         190
Leu Thr Ile Asp Gly Val Thr Arg Ser Asp Gln Gly Leu Tyr Thr Cys
195         200         205
Ala Ala Ser Ser Gly Leu Met Thr Lys Lys Asn Ser Thr Phe Val Arg
210         215         220
Val His Glu Lys Gly Pro Gly Asp Lys Thr His Thr Cys Pro Pro Cys
225         230         235         240

```


-continued

```

gacctgagg tcaagttaa ctggtacgtg gacggcgtgg aggtgcataa tgccaagaca    900
aagccgcggg aggagcagta caacagcacg taccgtgtgg tcagcgtcct caccgtcctg    960
caccaggact ggctgaatgg caaggagtac aagtgcaagg tctccaacaa agccctccca    1020
gcccccatcg agaaaacat ctccaaagcc aaagggcagc cccgagaacc acaggtgtac    1080
accctgcccc catcccggga tgagctgacc aagaaccagg tcagcctgac ctgcctggtc    1140
aaaggcttct atcccagcga catgcgcgtg gagtgggaga gcaatgggca gccggagaa    1200
aactacaaga ccacgcctcc cgtgctggac tccgacggct ccttcttct ctacagcaag    1260
ctcaccgtgg acaagagcag gtggcagcag gggaacgtct tctcatgctc cgtgatgcat    1320
gaggctctgc acaaccacta cagcagaag agcctctccc tgtctccggg taaatga      1377

```

```

<210> SEQ ID NO 4
<211> LENGTH: 458
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic

```

```

<400> SEQUENCE: 4

```

```

Met Val Ser Tyr Trp Asp Thr Gly Val Leu Leu Cys Ala Leu Leu Ser
 1             5             10             15
Cys Leu Leu Leu Thr Gly Ser Ser Ser Gly Ser Asp Thr Gly Arg Pro
      20             25             30
Phe Val Glu Met Tyr Ser Glu Ile Pro Glu Ile Ile His Met Thr Glu
      35             40             45
Gly Arg Glu Leu Val Ile Pro Cys Arg Val Thr Ser Pro Asn Ile Thr
      50             55             60
Val Thr Leu Lys Lys Phe Pro Leu Asp Thr Leu Ile Pro Asp Gly Lys
      65             70             75             80
Arg Ile Ile Trp Asp Ser Arg Lys Gly Phe Ile Ile Ser Asn Ala Thr
      85             90             95
Tyr Lys Glu Ile Gly Leu Leu Thr Cys Glu Ala Thr Val Asn Gly His
      100            105            110
Leu Tyr Lys Thr Asn Tyr Leu Thr His Arg Gln Thr Asn Thr Ile Ile
      115            120            125
Asp Val Val Leu Ser Pro Ser His Gly Ile Glu Leu Ser Val Gly Glu
      130            135            140
Lys Leu Val Leu Asn Cys Thr Ala Arg Thr Glu Leu Asn Val Gly Ile
      145            150            155            160
Asp Phe Asn Trp Glu Tyr Pro Ser Ser Lys His Gln His Lys Lys Leu
      165            170            175
Val Asn Arg Asp Leu Lys Thr Gln Ser Gly Ser Glu Met Lys Lys Phe
      180            185            190
Leu Ser Thr Leu Thr Ile Asp Gly Val Thr Arg Ser Asp Gln Gly Leu
      195            200            205
Tyr Thr Cys Ala Ala Ser Ser Gly Leu Met Thr Lys Lys Asn Ser Thr
      210            215            220
Phe Val Arg Val His Glu Lys Asp Lys Thr His Thr Cys Pro Pro Cys
      225            230            235            240
Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro
      245            250            255
Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
      260            265            270

```

-continued

Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp
		275					280					285			
Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu
	290					295					300				
Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu
305					310					315					320
His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn
			325						330					335	
Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly
		340						345					350		
Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Asp	Glu
		355					360					365			
Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr
	370					375					380				
Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn
385					390					395					400
Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe
			405					410						415	
Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn
		420						425				430			
Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr
		435					440					445			
Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly	Lys						
	450					455									

We claim:

1. A formulation comprising:
 10-50 mg/ml of a vascular endothelial growth factor (VEGF) antagonist fusion protein comprising amino acids 27-457 of SEQ ID NO: 4,
 a buffer comprising histidine,
 an organic co-solvent comprising polysorbate, and
 a stabilizing agent comprising a sugar, an amino acid or both,
 wherein said VEGF antagonist fusion protein exhibits less than about 3% degradation after 15 months of storage at 5° C.
2. The formulation of claim 1, wherein said buffer comprises 5-50 mM histidine.
3. The formulation of claim 1, wherein said buffer comprises about 10 mM histidine.
4. The formulation of claim 1, wherein said buffer is at a pH of about 6.0-6.5.
5. The formulation of claim 2, wherein said formulation comprises 40 mg/ml of said VEGF antagonist fusion protein.
6. The formulation of claim 2, wherein said formulation further comprises a tonicity agent.
7. The formulation of claim 6, wherein said tonicity agent is sodium chloride.
8. The formulation of claim 2, wherein said organic co-solvent comprises 0.1-0.5% polysorbate.
9. The formulation of claim 2, wherein said organic co-solvent comprises polysorbate 20.
10. The formulation of claim 2, wherein said stabilizing agent comprises a sugar.
11. The formulation of claim 10, wherein said sugar is selected from group consisting of dextrose, ribose, fructose, mannitol, inositol, sorbitol, trehalose, sucrose, and lactose.
12. The formulation of claim 10, wherein said sugar is 5-20% sucrose.
13. The formulation of claim 10, wherein said sugar is 0.5-6.0% sucrose.
14. The formulation of claim 2, wherein said stabilizing agent comprises an amino acid.
15. The formulation of claim 14, wherein said amino acid is glycine.
16. The formulation of claim 14, wherein said amino acid is 0.25-3.0% glycine.
17. The formulation of claim 9, wherein said formulation comprises 40 mg/ml of said VEGF antagonist fusion protein.
18. The formulation of claim 9, wherein said formulation further comprises a tonicity agent.
19. The formulation of claim 18, wherein said tonicity agent is sodium chloride.
20. The formulation of claim 9, wherein said stabilizing agent comprises a sugar.
21. The formulation of claim 20, wherein said sugar is selected from group consisting of dextrose, ribose, fructose, mannitol, inositol, sorbitol, trehalose, sucrose, and lactose.
22. The formulation of claim 20, wherein said sugar is 5-20% sucrose.
23. The formulation of claim 20, wherein said sugar is 0.5-6.0% sucrose.
24. The formulation of claim 9, wherein said stabilizing agent comprises an amino acid.
25. The formulation of claim 24, wherein said amino acid is glycine.
26. The formulation of claim 24, wherein said amino acid is 0.25-3.0% glycine.
27. A formulation comprising:
 10-50 mg/ml of a vascular endothelial growth factor (VEGF) antagonist fusion protein comprising amino acids 27-457 of SEQ ID NO: 4,

21

a buffer comprising histidine,
an organic co-solvent comprising polysorbate, and
a stabilizing agent comprising a sugar, an amino acid or
both,

wherein said VEGF antagonist fusion protein is capable
of inhibiting biological activity of human VEGF as
measured by a mouse Baf/3 VEGFR1/EpoR cell line
and achieving a percent relative potency of at least 75
relative to a reference VEGF IC₅₀ standard.

28. The formulation of claim 27, wherein said buffer
comprises 5-50 mM histidine.

29. The formulation of claim 27, wherein said percent
relative potency of at least 75 is achieved after 1 month of
storage at 5° C.

30. The formulation of claim 27, wherein said percent
relative potency of at least 75 is achieved after 24 months of
storage at 5° C.

31. A formulation comprising:

10-50 mg/ml of a vascular endothelial growth factor
(VEGF) antagonist fusion protein comprising amino
acids 27-457 of SEQ ID NO: 4,

a buffer comprising histidine,
an organic co-solvent comprising polysorbate, and
a stabilizing agent comprising a sugar, an amino acid or
both,

wherein said VEGF antagonist fusion protein is capable
of binding VEGF at a percent relative binding of at
least 88 relative to a reference VEGF IC₅₀ standard,
after storage at 5° C. for 3 months.

32. The formulation of claim 31, wherein said buffer
comprises 5-50 mM histidine.

33. A method of manufacturing a liquid formulation, said
method comprising combining a vascular endothelial
growth factor (VEGF) antagonist fusion protein that com-
prises amino acids 27-457 of SEQ ID NO: 4 with a buffer
comprising histidine, an organic co-solvent comprising
polysorbate, and a stabilizing agent comprising a sugar, an
amino acid or both;

wherein the concentration of said VEGF antagonist fusion
protein is 10-50 mg/ml in said liquid formulation; and
wherein said VEGF antagonist fusion protein exhibits less
than about 3% degradation after 15 months of storage
at 5° C.

34. The method of claim 33, wherein said buffer com-
prises 5-50 mM histidine.

35. A method of manufacturing a liquid formulation, said
method comprising combining a vascular endothelial
growth factor (VEGF) antagonist fusion protein that com-
prises amino acids 27-457 of SEQ ID NO: 4 with a buffer
comprising histidine, an organic co-solvent comprising
polysorbate, and a stabilizing agent comprising a sugar, an
amino acid or both;

wherein the concentration of said VEGF antagonist fusion
protein is 10-50 mg/ml in said liquid formulation; and
wherein said VEGF antagonist fusion protein is capable
of inhibiting biological activity of human VEGF as
measured by a mouse Baf/3 VEGFR1/EpoR cell line
and achieving a percent relative potency of at least 75
relative to a reference VEGF IC₅₀ standard.

36. The method of claim 35, wherein said buffer com-
prises 5-50 mM histidine.

37. The method of claim 35, wherein said percent relative
potency of at least 75 is achieved after 1 month of storage
at 5° C.

38. The formulation of claim 35, wherein said percent
relative potency of at least 75 is achieved after 24 months of
storage at 5° C.

22

39. A method of manufacturing a liquid formulation, said
method comprising combining a vascular endothelial
growth factor (VEGF) antagonist fusion protein that com-
prises amino acids 27-457 of SEQ ID NO: 4 with a buffer
comprising histidine, an organic co-solvent comprising
polysorbate, and a stabilizing agent comprising a sugar, an
amino acid or both;

wherein the concentration of said VEGF antagonist fusion
protein is 10-50 mg/ml in said liquid formulation; and
wherein said VEGF antagonist fusion protein is capable
of binding VEGF at a percent relative binding of at
least 88 relative to a reference VEGF IC₅₀ standard,
after storage at 5° C. for 3 months.

40. The method of claim 39, wherein said buffer com-
prises 5-50 mM histidine.

41. The formulation of claim 2, wherein said organic
co-solvent comprises 0.05-0.15% polysorbate 20.

42. The formulation of claim 2, wherein said organic
co-solvent comprises 0.003-0.15% polysorbate.

43. The formulation of claim 29, wherein said buffer
comprises 5-50 mM histidine.

44. The formulation of claim 29, wherein said buffer
comprises about 10 mM histidine.

45. The formulation of claim 43, wherein said buffer is at
a pH of about 6.0-6.5.

46. The formulation of claim 43, wherein said formulation
comprises 40 mg/ml of said VEGF antagonist fusion protein.

47. The formulation of claim 46, wherein said formulation
further comprises a tonicity agent.

48. The formulation of claim 47, wherein said tonicity
agent is sodium chloride.

49. The formulation of claim 46, wherein said organic
co-solvent comprises 0.1-0.5% polysorbate.

50. The formulation of claim 46, wherein said organic
co-solvent comprises 0.05-0.15% polysorbate 20.

51. The formulation of claim 46, wherein said organic
co-solvent comprises 0.003-0.15% polysorbate.

52. The formulation of claim 46, wherein said stabilizing
agent comprises a sugar.

53. The formulation of claim 52, wherein said sugar is
selected from group consisting of dextrose, ribose, fructose,
mannitol, inositol, sorbitol, trehalose, sucrose, and lactose.

54. The formulation of claim 52, wherein said sugar is
5-20% sucrose.

55. The formulation of claim 52, wherein said sugar is
0.5-6.0% sucrose.

56. The formulation of claim 46, wherein said stabilizing
agent comprises an amino acid.

57. The formulation of claim 56, wherein said amino acid
is glycine.

58. The formulation of claim 31, wherein said buffer
comprises about 10 mM histidine.

59. The formulation of claim 32, wherein said buffer is at
a pH of about 6.0-6.5.

60. The formulation of claim 32, wherein said formulation
comprises 40 mg/ml of said VEGF antagonist fusion protein.

61. The formulation of claim 60, wherein said formulation
further comprises a tonicity agent.

62. The formulation of claim 61, wherein said tonicity
agent is sodium chloride.

63. The formulation of claim 60, wherein said organic
co-solvent comprises 0.1-0.5% polysorbate.

64. The formulation of claim 60, wherein said organic
co-solvent comprises 0.05-0.15% polysorbate 20.

65. The formulation of claim 60, wherein said organic
co-solvent comprises 0.003-0.15% polysorbate.

23

66. The formulation of claim 60, wherein said stabilizing agent comprises a sugar.

67. The formulation of claim 66, wherein said sugar is selected from group consisting of dextrose, ribose, fructose, mannitol, inositol, sorbitol, trehalose, sucrose, and lactose.

68. The formulation of claim 66, wherein said sugar is 5-20% sucrose.

69. The formulation of claim 66, wherein said sugar is 0.5-6.0% sucrose.

70. The formulation of claim 60, wherein said stabilizing agent comprises an amino acid.

71. The formulation of claim 70, wherein said amino acid is glycine.

72. The method of claim 33, wherein said buffer comprises about 10 mM histidine.

73. The method of claim 34, wherein said buffer is at a pH of about 6.0-6.5.

74. The method of claim 34, wherein said formulation comprises 40 mg/ml of said VEGF antagonist fusion protein.

75. The method of claim 74, wherein said formulation further comprises a tonicity agent.

76. The method of claim 75, wherein said tonicity agent is sodium chloride.

77. The method of claim 74, wherein said organic co-solvent comprises 0.1-0.5% polysorbate.

78. The method of claim 74, wherein said organic co-solvent comprises 0.05-0.15% polysorbate 20.

79. The method of claim 74, wherein said organic co-solvent comprises 0.003-0.15% polysorbate.

80. The method of claim 74, wherein said stabilizing agent comprises a sugar.

81. The method of claim 80, wherein said sugar is selected from group consisting of dextrose, ribose, fructose, mannitol, inositol, sorbitol, trehalose, sucrose, and lactose.

82. The method of claim 80, wherein said sugar is 5-20% sucrose.

83. The method of claim 80, wherein said sugar is 0.5-6.0% sucrose.

84. The method of claim 74, wherein said stabilizing agent comprises an amino acid.

85. The method of claim 84, wherein said amino acid is glycine.

86. The method of claim 37, wherein said buffer comprises 5-50 mM histidine.

87. The method of claim 37, wherein said buffer comprises about 10 mM histidine.

88. The method of claim 86, wherein said buffer is at a pH of about 6.0-6.5.

89. The method of claim 86, wherein said formulation comprises 40 mg/ml of said VEGF antagonist fusion protein.

90. The method of claim 89, wherein said formulation further comprises a tonicity agent.

24

91. The method of claim 90, wherein said tonicity agent is sodium chloride.

92. The method of claim 89, wherein said organic co-solvent comprises 0.1-0.5% polysorbate.

93. The method of claim 89, wherein said organic co-solvent comprises 0.05-0.15% polysorbate 20.

94. The method of claim 89, wherein said organic co-solvent comprises 0.003-0.15% polysorbate.

95. The method of claim 89, wherein said stabilizing agent comprises a sugar.

96. The method of claim 95, wherein said sugar is selected from group consisting of dextrose, ribose, fructose, mannitol, inositol, sorbitol, trehalose, sucrose, and lactose.

97. The method of claim 95, wherein said sugar is 5-20% sucrose.

98. The method of claim 95, wherein said sugar is 0.5-6.0% sucrose.

99. The method of claim 89, wherein said stabilizing agent comprises an amino acid.

100. The method of claim 99, wherein said amino acid is glycine.

101. The method of claim 39, wherein said buffer comprises about 10 mM histidine.

102. The method of claim 40, wherein said buffer is at a pH of about 6.0-6.5.

103. The method of claim 40, wherein said formulation comprises 40 mg/ml of said VEGF antagonist fusion protein.

104. The method of claim 103, wherein said formulation further comprises a tonicity agent.

105. The method of claim 104, wherein said tonicity agent is sodium chloride.

106. The method of claim 103, wherein said organic co-solvent comprises 0.1-0.5% polysorbate.

107. The method of claim 103, wherein said organic co-solvent comprises 0.05-0.15% polysorbate 20.

108. The method of claim 103, wherein said organic co-solvent comprises 0.003-0.15% polysorbate.

109. The method of claim 103, wherein said stabilizing agent comprises a sugar.

110. The method of claim 109, wherein said sugar is selected from group consisting of dextrose, ribose, fructose, mannitol, inositol, sorbitol, trehalose, sucrose, and lactose.

111. The method of claim 109, wherein said sugar is 5-20% sucrose.

112. The method of claim 109, wherein said sugar is 0.5-6.0% sucrose.

113. The method of claim 103, wherein said stabilizing agent comprises an amino acid.

114. The method of claim 113, wherein said amino acid is glycine.

* * * * *