



US011253572B2

**(12) United States Patent**  
**Yancopoulos****(10) Patent No.: US 11,253,572 B2****(45) Date of Patent: \*Feb. 22, 2022****(54) USE OF A VEGF ANTAGONIST TO TREAT**  
**ANGIOGENIC EYE DISORDERS**(2013.01); *A61K 2039/505* (2013.01); *C07K*  
*2319/30* (2013.01); *C07K 2319/32* (2013.01)**(71) Applicant: REGENERON**  
**PHARMACEUTICALS, INC.,**  
Tarrytown, NY (US)**(58) Field of Classification Search**

None

See application file for complete search history.

**(72) Inventor: George Yancopoulos,** Yorktown  
Heights, NY (US)**(56) References Cited**

## U.S. PATENT DOCUMENTS

**(73) Assignee: REGENERON**  
**PHARMACEUTICALS, INC.,**  
Tarrytown, NY (US)

6,171,586	B1	1/2001	Lam et al.
6,833,349	B2	12/2004	Xia et al.
6,879,294	B2	5/2005	Davis-Smyth et al.
6,897,294	B2	5/2005	Davis-Smyth et al.
7,070,959	B1	7/2006	Papadopoulos
7,303,746	B2	12/2007	Wiegand
7,303,747	B2	12/2007	Wiegand et al.
7,303,748	B2	12/2007	Wiegand
7,306,799	B2	12/2007	Wiegand
7,374,757	B2	5/2008	Papadopoulos et al.
7,374,758	B2	5/2008	Papadopoulos et al.
7,378,095	B2	5/2008	Cao et al.
7,396,664	B2	7/2008	Daly et al.
7,482,002	B2	1/2009	Cedarbaum
7,521,049	B2	4/2009	Wiegand et al.
7,531,173	B2	5/2009	Wiegand et al.
7,750,138	B2	7/2010	Fang et al.
7,951,585	B2	5/2011	Ke
8,092,803	B2	1/2012	Furfine et al.

(Continued)

**(\*) Notice:** Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 0 days.This patent is subject to a terminal dis-  
claimer.**(21) Appl. No.: 17/352,892****(22) Filed: Jun. 21, 2021****(65) Prior Publication Data**

US 2021/0308217 A1 Oct. 7, 2021

**Related U.S. Application Data****(63)** Continuation of application No. 17/350,958, filed on  
Jun. 17, 2021, which is a continuation of application  
No. 17/112,404, filed on Dec. 4, 2020, which is a  
continuation of application No. 17/072,417, filed on  
Oct. 16, 2020, which is a continuation of application  
No. 16/055,847, filed on Aug. 6, 2018, now Pat. No.  
10,857,205, and a continuation of application No.  
16/397,267, filed on Apr. 29, 2019, now Pat. No.  
10,888,601, which is a continuation of application  
No. 16/159,282, filed on Oct. 12, 2018, now Pat. No.  
10,828,345, which is a continuation of application  
No. 15/471,506, filed on Mar. 28, 2017, now Pat. No.  
10,130,681, which is a continuation of application  
No. 14/972,560, filed on Dec. 17, 2015, now Pat. No.  
9,669,069, which is a continuation of application No.  
13/940,370, filed on Jul. 12, 2013, now Pat. No.  
9,254,338, which is a continuation-in-part of  
application No. PCT/US2012/020855, filed on Jan.  
11, 2012.**(60)** Provisional application No. 61/432,245, filed on Jan.  
13, 2011, provisional application No. 61/434,836,  
filed on Jan. 21, 2011, provisional application No.  
61/561,957, filed on Nov. 21, 2011.**(51) Int. Cl.***A61K 38/17* (2006.01)*C07K 14/71* (2006.01)*C07K 19/00* (2006.01)*C07K 16/22* (2006.01)*A61K 9/00* (2006.01)*A61K 39/00* (2006.01)**(52) U.S. Cl.**CPC ..... *A61K 38/179* (2013.01); *A61K 9/0048*  
(2013.01); *C07K 14/71* (2013.01); *C07K 16/22*

## FOREIGN PATENT DOCUMENTS

CN	1304427	C	3/2007
CN	100502945	C	6/2009

(Continued)

## OTHER PUBLICATIONS

U.S. Appl. No. 16/055,847—Third Party Submissions dated May 1,  
2019.

(Continued)

*Primary Examiner* — Christine J Saoud*Assistant Examiner* — Jon M Lockard**(74) Attorney, Agent, or Firm** — Thomas Triolo; Karl  
Bozicevic**(57)****ABSTRACT**

The present invention provides methods for treating angio-  
genic eye disorders by sequentially administering multiple  
doses of a VEGF antagonist to a patient. The methods of the  
present invention include the administration of multiple  
doses of a VEGF antagonist to a patient at a frequency of  
once every 8 or more weeks. The methods of the present  
invention are useful for the treatment of angiogenic eye  
disorders such as age related macular degeneration, diabetic  
retinopathy, diabetic macular edema, central retinal vein  
occlusion, branch retinal vein occlusion, and corneal neo-  
vascularization.

**30 Claims, 1 Drawing Sheet****Specification includes a Sequence Listing.**

(56)

**References Cited****U.S. PATENT DOCUMENTS**

8,216,575	B2	7/2012	Yu
9,254,338	B2	2/2016	Yancopoulos
9,657,084	B2	5/2017	Ke et al.
9,669,069	B2	6/2017	Yancopoulos
10,130,681	B2	11/2018	Yancopoulos
10,406,226	B2	9/2019	Dix et al.
10,464,992	B2	11/2019	Furfine et al.
10,828,345	B2	11/2020	Yancopoulos
2003/0113316	A1	6/2003	Kaisheva et al.
2003/0138417	A1	7/2003	Kaisheva et al.
2003/0171320	A1	9/2003	Guyer
2004/0197324	A1	10/2004	Liu et al.
2004/0213787	A1	10/2004	Sleeman et al.
2004/0266688	A1	12/2004	Nayak
2005/0032699	A1	2/2005	Holash et al.
2005/0163798	A1	7/2005	Papadopoulos et al.
2005/0260203	A1	11/2005	Wiegand et al.
2005/0281822	A1	12/2005	Cedarbaum et al.
2006/0030000	A1	2/2006	Alitalo et al.
2006/0058234	A1	3/2006	Daly et al.
2006/0172944	A1	8/2006	Wiegand et al.
2006/0217311	A1	9/2006	Dix et al.
2007/0190058	A1	8/2007	Shams
2008/0220004	A1	9/2008	Wiegand et al.
2009/0264358	A1	10/2009	Yu
2013/0295094	A1	11/2013	Yancopoulos
2016/0130337	A1	5/2016	Gekkieva et al.
2019/0290725	A1	9/2019	Vitti et al.
2019/0388539	A1	12/2019	Dix et al.
2020/0017572	A1	1/2020	Furfine et al.

**FOREIGN PATENT DOCUMENTS**

CN	100567325	C	12/2009
CN	102233132	B	10/2013
CN	102380096	B	4/2014
CN	103212075	B	6/2017
CN	107115294	A	9/2017
EP	2663325		11/2013
JP	2010-509369		3/2010
WO	97/04801		2/1997
WO	WO 97/04801		2/1997
WO	WO 2000/75319		12/2000
WO	WO 2004/106378	A2	12/2004
WO	WO 2005/000895	A2	1/2005
WO	WO 2006/047325		3/2006
WO	WO 2007/022101	A2	2/2007
WO	WO 2008/063932		5/2008
WO	WO 2012/097019		7/2012

**OTHER PUBLICATIONS**

U.S. Appl. No. 16/159,282—Third Party Submissions dated May 31, 2019.

Adsis R&D Profile “Aflibercept: AVE 0005, AVE 005, AVE0005, VEGF Trap—Regeneron, VEGF Trap (R1R2), VEGF Trap-Eye.” Drugs R D, 9(4):261-269 (2008).

Anonymous “Lucentis (ranibizumab injection) Intravitreal Injection” pp. 103 (Jun. 2006).

Anonymous “Anti-VEGF 2019: The State of the Art” Review of Ophthalmology (published Aug. 5, 2019).

Barbazetto, “Dosing Regimen And The Frequency Of Macular Hemorrhages In Neovascular Age-Related Macular Degeneration Treated With Ranibizumab.” Retina, 30(9):1376-85 (2010).

Bayer Investor News, “Bayer and Regeneron Start additional Phase 3 Study for VEGF Trap-Eye in Wet Age-related Macular Degeneration.” (May 8, 2008).

Bayer Investor News, “VEGF Trap-Eye: New Data Confirm Successes in the Treatment of Age-related Macular Degeneration” (Sep. 28, 2008).

Benz et al. “CLEAR-IT-2: Interim Results Of The Phase II, Randomized, Controlled Dose-and Interval-ranging Study Of Repeated

Intravitreal VEGF Trap Administration In Patients With Neovascular Age-related Macular Degeneration (AMD)” ARVO Annual Meeting Abstract (May 2007).

Boyer, “A Phase IIIb Study to Evaluate the Safety of Ranibizumab in Subjects with Neovascular Age-related Macular Degeneration.” Ophthalmology, 116(9):1731-39 (Sep. 2009).

Brown, “Ranibizumab versus Verteporfin for Neovascular Age-Related Macular Degeneration.” N Engl J Med, 355(14):1432-44 (Oct. 5, 2006).

Brown, “Primary Endpoint Results of a Phase II Study of Vascular Endothelial Growth Factor Trap-Eye in Wet Age-related Macular Degeneration.” Ophthalmology, 118(6): 1089-97 (Jun. 2011).

Brown, “Long-term Outcomes of Ranibizumab Therapy for Diabetic Macular Edema: The 36-Month Results from Two phase III Trials.” Ophthalmology, 120(10):2013-22 (Oct. 2013).

Browning et al. “Aflibercept for age-related macular degeneration: a game-changer or quiet addition?” American Journal of Ophthalmology, 154(2):222-226 (Aug. 2012).

Campochiaro et al. “Ranibizumab for Macular Edema Due to Retinal Vein Occlusions Implication of VEGF as a Critical Stimulator” Molecular Therapy, 16(4):791-799 (2008).

Campochiaro, “Ranibizumab for Macular Edema following Branch Retinal Vein Occlusion: six-month primary end point results of a phase III study.” Ophthalmology, 117(6):1102-1112 (Jun. 2010).

Campochiaro, “Sustained Benefits from Ranibizumab for Macular Edema following Central Retinal Vein Occlusion: Twelve-Month Outcomes of a phase III Study.” Ophthalmology, 188(10):2041-49 (Oct. 2011).

Cao, “A Subretinal Matrigel Rat Choroidal Neovascularization (CNV) Model and Inhibition of CNV and Associated Inflammation and Fibrosis by VEGF Trap” Investigative Ophthalmology & Visual Science, 51 (11):6009-6017 (Nov. 2010).

Center for Drug Evaluation and Research BLA Application No. 125156 Medical Review, (Jun. 2006) <URL:https://www.accessdata.fda.gov/drugsatfda\_docs/nda/2006/125156s000\_Lucentis\_MedR.pdf>.

Charles, Steve (Guest Lecturer) “VEGF Trap Has Positive DME Data” Tenth Annual Retina Fellows Forum Jan. 29 and 30, Chicago, (Article Date Mar. 1, 2010).

Chatziralli et al. “Intravitreal aflibercept for neovascular age-related macular degeneration in patients aged 90 years or older: 2-year visual acuity outcomes” Eye (2018) 32:1523-1529.

Chung et al. “Ziv-aflibercept: A novel angiogenesis inhibitor for the treatment of metastatic colorectal cancer” Am J Heath-Syst Pharm (Nov. 1, 2013) 70:1887-1896.

Cooper et al., “Increased Renal Expression of Vascular Endothelial Growth Factor (VEGF) and Its Receptor VEGFR-2 in Experimental Diabetes” Diabetes (1999) 48:2229-2239.

Croll et al., “VEGF-mediated inflammation precedes angiogenesis in adult brain” Experimental Neurology (2004) 187:388-402.

Csaky, “Safety Implications of Vascular Endothelial Growth Factor Blockade for Subjects Receiving Intravitreal Anti-Vascular Endothelial Growth Factor Therapies.” Am. J. Ophthalmology, 148(5):647-56, (Nov. 2009).

DeVries et al., “Antibodies against Vascular Endothelial Growth Factor Improve Early Renal Dysfunction in Experimental Diabetes” J. Am. Soc. Nephrol (2001) 12:993-1000.

Do et al., “An exploratory study of the safety, tolerability and bioactivity of a single intravitreal injection of vascular endothelial growth factor Trap-Eye in patients with diabetic macular oedema” Br J Ophthalmol. 93(2):144-1449 (Feb. 2009).

Do et al., “The DA VINCI Study: phase 2 primary results of VEGF Trap-Eye in patients with diabetic macular edema” Ophthalmology, 118(9): 1819-1826 (Sep. 2011).

Do, “One-Year Outcomes of the DA VINCI Study of VEGF Trap-Eye in Eyes with Diabetic Macular Edema.” Ophthalmology, 119(8):1658-65 (2012).

Do et al. “Results of a Phase 1 Study of Intravitreal VEGF Trap in Subjects with Diabetic Macular Edema: The CLEAR-IT DME Study” ARVO Annual Meeting Abstract (May 2007).

(56)

**References Cited****OTHER PUBLICATIONS**

- Do et al. "VEGF Trap-Eye Vision-specific Quality of Life through 52 Weeks in Patients with Neovascular AMD in CLEAR-IT 2: A Phase 2 Clinical Trial" ARVO Annual Meeting Abstract (Apr. 2009).
- Eichten, "Rapid decrease in tumor perfusion following VEGF blockade predicts long-term tumor growth inhibition in preclinical tumor models" *Angiogenesis*, 16:429-441 (2013).
- Engelbert, "Treat And Extend" Dosing Of Intravitreal Antivascular Endothelial Growth Factor Therapy For Type 3 Neovascularization/Retinal Angiomatous Proliferation." *Retina*, 29(10):1424-31 (2009).
- Engelbert, "Long-Term Follow-Up For Type 1 (Subretinal Pigment Epithelium) Neovascularization Using A Modified 'Treat And Extend' Dosing Regimen Of Intravitreal Antivascular Endothelial Growth Factor Therapy." *Retina*, 30(9):1368-75 (2010).
- Engelbert, "The 'Treat and Extend' Dosing Regimen of Intravitreal Anti-Vascular Endothelial Growth Factor Therapy for Neovascular Age-Related Macular Degeneration." *Ophthalmology Management*, Issue 42, (Jun. 2010) available at <http://www.visioncareprofessional.com/emails/amdupdate/index.asp?issue=42>.
- Eremina et al., "Glomerular-specific alterations of VEGF-A expression lead to distinct congenital and acquired renal diseases" *Journal of Clinical Investigation* (Mar. 2003) 111(5):707-716.
- Eriksson et al., "Structure, Expression and Receptor-Binding Properties of Novel Vascular Endothelial Growth Factors" *Vascular Growth Factors and Angiogenesis*, Springer (1999) pp. 41-57.
- The Eyetech Study Group, "Anti-Vascular Endothelial Growth Factor Therapy for Subfoveal Choroidal Neovascularization Secondary to Age-related Macular Degeneration" *American Academy of Ophthalmology*, 110(5):979-986 (May 2003).
- Eylea®, Highlights of Prescribing Information, Revised Aug. 2018.
- Ferrara, N. "Vascular Endothelial Growth Factor: Molecular and Biological Aspects" *Advances in Organ Biology* (1999) pp. 1-30.
- Ferrara et al., "Clinical applications of angiogenic growth factors and their inhibitors" *Nature Medicine* (Dec. 1999) 5(12):1359-1364.
- Flyvbjerg et al., "Amelioration of Long-Term Renal Changes in Obese Type 2 Diabetic Mice by a Neutralizing Vascular Endothelial Growth Factor Antibody" *Diabetes* (Oct. 2002) 51:3090-3094.
- Fung, "An Optical Coherence Tomography-Guided, Variable Dosing Regimen with Intravitreal Ranibizumab (Lucentis) for Neovascular Age-related Macular Degeneration." *Am J Ophthalmology*, 143(4):566-83 (Apr. 2007).
- Gale, "Complementary and Coordinated Roles of the VEGFs and Angiopoietins during Normal and Pathologic Vascular Formation." *Cold Spring Harbor Symposia on Quantitative Biology*, vol. LXVII, pp. 267-273 (2002).
- Garcia-Quintanilla, "Pharmacokinetics of Intravitreal Anti-VEGF Drugs in Age-Related Macular Degeneration." *Pharmaceutics*, 11:365 (2019).
- Gomez-Manzano, "VEGF Trap induces antiglioma effect at different stages of disease." *Neuro-Oncology*, 10:940-945 (Dec. 2008).
- Gragoudas, "Pegaptanib for Neovascular Age-Related Macular Degeneration." *N Engl J Med*, 351(27):2805-16, (Dec. 30, 2004).
- Gutierrez et al., "Intravitreal bevacizumab (Avastin) in the treatment of macular edema secondary to retinal vein occlusion" *Clin. Ophthalmol.*, 2(4):787,791 (2008).
- Haller et al., "VEGF Trap-Eye In CRVO: Primary Endpoint Results of the Phase 3 COPERNICUS Study" ARVO Annual Meeting Abstract (Apr. 2011).
- Heier et al., "CLEAR-IT 2: Phase 2, Randomized Controlled Dose and Interval-Ranging Study of Intravitreal VEGF Trap Eye in Patients with Neovascular Age-Related Macular Degeneration: Predictive Factors for Visual Acuity" ARVO Annual Meeting Abstract (Apr. 2009).
- Heier, J., "Intravitreal VEGF Trap for AMD: An Update, The CLEAR-IT 2 Extension Study" Presented at the annual meeting of the Association for Research in Vision and Ophthalmology, Retina Today (2009) pp. 44-45.
- Heier et al., "rhuFab V2 (anti-VEGF Antibody) for Treatment of Exudative AMD" Symposium 8:Experimental and Emerging Treatments for Choroidal Neovascularization, 10 pp (2002).
- Heier et al., "RhuFab V2 in Wet AMD—6 Month Continued Improvement Following Multiple Intravitreal Injections" *Invest Ophthalmol Vis Sci*, 44(E-Abstract):972 (2003).
- Heier et al., "Intravitreal Aflibercept (VEGF Trap-Eye) in Wet Age-related macular Degeneration," *Ophthalmology*, 119:2537-2548 (2012).
- Heier, "Intravitreal Aflibercept for Diabetic Macular Edema: 148-Week Results from the VISTA and VIVID Studies." *Ophthalmology*, 123(11):2376-2385 (Nov. 2016).
- Heier et al., "The 1-year Results of CLEAR-IT 2, a Phase 2 Study of Vascular Endothelial Growth Factor Trap-Eye Dosed As-needed After 12-week Fixed Dosing" *Ophthalmology* 2011;118:1098-1106 (Jun. 2011).
- Heier et al., "The 1-year Results of CLEAR-IT 2, a Phase 2 Study of Vascular Endothelial Growth Factor Trap-Eye Dosed As-needed After 12-week Fixed Dosing: Erratum" *Ophthalmology* 2011;118:1700 (Sep. 2011).
- Ho, "VEGF Trap-Eye in Wet AMD—CLEAR-IT 2: One-Year OCT and FA Outcomes" CLEAR-IT 2 Study Group, pp. 1-24 (Sep. 28, 2008).
- Ho et al., Slides entitled CLEAR IT 2 One-Year Key Results, Retina Society (2008).
- Holash et al., "Vessel Coooption, Regression, and Growth in Tumors Mediated by Angiopoietins and VEGF" *Science* (Jun. 18, 1999) 284(5422):1994-1998.
- Holash, "VEGF-Trap: A VEGF blocker with potent antitumor effects" *PNAS* 99(17):11393-11398 (Aug. 20, 2002).
- Holash, "Inhibitors of growth factor receptors, signaling pathways and angiogenesis as therapeutic molecular agents." *Cancer Metastasis* 25:243-252 (2006).
- Information from ClinicalTrials.gov archive History of Changes for Study: NCT00320775 "Safety and Tolerability of Intravitreal Administration of VEGF Trap in Patients With Neovascular Age-Related Macular Degeneration" 70 pages, Latest version submitted Jun. 8, 2011 on ClinicalTrials.gov (NCT00320775\_2006-2011).
- Information from ClinicalTrials.gov archive History of Changes for Study: NCT00320775 "Safety and Tolerability of Intravitreal Administration of VEGF Trap in Patients With Neovascular Age-Related Macular Degeneration" 10 pages, Latest version submitted Mar. 16, 2015 on ClinicalTrials.gov (NCT00320775\_2015).
- Information from ClinicalTrials.gov archive History of Changes for Study: NCT00320788 "Safety and Efficacy of Repeated Intravitreal Administration of Vascular Endothelial Growth Factor (VEGF) Trap in Patients With Wet Age-Related Macular Degeneration (AMD)" 71 pages, Latest version submitted Dec. 1, 2011 on ClinicalTrials.gov (NCT00320788\_2006-2011).
- Information from ClinicalTrials.gov archive History of Changes for Study: NCT00320788 "Safety and Efficacy of Repeated Intravitreal Administration of Vascular Endothelial Growth Factor (VEGF) Trap in Patients With Wet Age-Related Macular Degeneration (AMD)" 31 pages, Latest version submitted Jan. 27, 2012 on ClinicalTrials.gov (NCT00320788\_2012).
- Information from ClinicalTrials.gov archive History of Changes for Study: NCT00320814 "Phase 1 Study of VEGF Trap in Patients With Diabetic Macular Edema" 30 pages, Latest version submitted Jun. 8, 2011 on ClinicalTrials.gov (NCT00320814\_2006-2011).
- Information from ClinicalTrials.gov archive History of Changes for Study: NCT00509795 "Double-Masked Study of Efficacy and Safety of IVT VEGF Trap-Eye in Subjects With Wet AMD (VIEW 1)" 318 pages, Latest version submitted Dec. 1, 2011 on ClinicalTrials.gov (NCT00509795\_2007-2011).
- Information from ClinicalTrials.gov archive History of Changes for Study: NCT00509795 "Double-Masked Study of Efficacy and Safety of IVT VEGF Trap-Eye in Subjects With Wet AMD (VIEW 1)" 200 pages, Latest version submitted Dec. 20, 2012 on ClinicalTrials.gov (NCT00509795\_2012).
- Information from ClinicalTrials.gov archive History of Changes for Study: NCT00527423 "Randomized, Single-Masked, Long-Term,

(56)

**References Cited****OTHER PUBLICATIONS**

Safety and Tolerability Study of VEGF Trap-Eye in AMD" 64 pages, Latest version submitted Nov. 1, 2011 on ClinicalTrials.gov (NCT00527423\_2007-2011).

Information from ClinicalTrials.gov archive History of Changes for Study: NCT00527423 "Randomized, Single-Masked, Long-Term, Safety and Tolerability Study of VEGF Trap-Eye in AMD" 42 pages, Latest version submitted Jun. 10, 2013 on ClinicalTrials.gov (NCT00527423\_2012-2013).

Information from ClinicalTrials.gov archive History of Changes for Study: NCT00637377 "Vascular Endothelial Growth Factor (VEGF) Trap-Eye: Investigation of Efficacy and Safety in Wet Age-Related Macular Degeneration (AMD) (VIEW 2)" 667 pages, Latest version submitted Dec. 16, 2011 on ClinicalTrials.gov (NCT00637377\_2008-2011).

Information from ClinicalTrials.gov archive History of Changes for Study: NCT00637377 "Vascular Endothelial Growth Factor (VEGF) Trap-Eye: Investigation of Efficacy and Safety in Wet Age-Related Macular Degeneration (AMD) (VIEW 2)" 289 pages, Latest version submitted Nov. 28, 2014 on ClinicalTrials.gov (NCT00637377\_2012-2014).

Information from ClinicalTrials.gov archive History of Changes for Study: NCT00789477 "DME And VEGF Trap-Eye [Intravitreal Aflibercept Injection (IAI;EYLEA®;BAY86-5321)] INvestigation of Clinical Impact (DA VINCI)" 135 pages, Latest version submitted May 2, 2011 on ClinicalTrials.gov (NCT00789477\_2008-2011).

Information from ClinicalTrials.gov archive History of Changes for Study: NCT00789477 "DME And VEGF Trap-Eye [Intravitreal Aflibercept Injection (IAI;EYLEA®;BAY86-5321)] INvestigation of Clinical Impact (DA VINCI)" 53 pages, Latest version submitted Aug. 28, 2014 on ClinicalTrials.gov (NCT00789477\_2013-2014).

Information from ClinicalTrials.gov archive History of Changes for Study: NCT00943072 "Vascular Endothelial Growth Factor (VEGF) Trap-Eye: Investigation of Efficacy and Safety in Central Retinal Vein Occlusion (CRVO)" 98 pages, Latest version submitted May 9, 2011 on ClinicalTrials.gov (NCT00943072\_2009-2011).

Information from ClinicalTrials.gov archive History of Changes for Study: NCT00943072 "Vascular Endothelial Growth Factor (VEGF) Trap-Eye: Investigation of Efficacy and Safety in Central Retinal Vein Occlusion (CRVO)" 64 pages, Latest version submitted Apr. 16, 2013 on ClinicalTrials.gov (NCT00943072\_2012-2013).

Information from ClinicalTrials.gov archive View of NCT00637377 "Vascular Endothelial Growth Factor (VEGF) Trap-Eye: Investigation of Efficacy and Safety in Wet Age-Related Macular Degeneration (AMD) (VIEW 2)" ClinicalTrials.gov. Web. (Nov. 30, 2010). Information from ClinicalTrials.gov archive on the VIEW 2 study (NCT00637377) "VEGF Trap-Eye: Investigation of Efficacy and Safety in Wet AMD (VIEW 2)" version available (updated on Mar. 17, 2008).

Information from ClinicalTrials.gov archive on the view of NCT00509795 "Vascular Endothelial Growth Factor (VEGF) Trap-Eye: Investigation of Efficacy and Safety in Wet Age-Related Macular Degeneration (AMD)" (Dec. 1, 2009).

Information from ClinicalTrials.gov archive on the view of NCT00789477 "DME and VEGF Trap-Eye: Investigation of Clinical Impact" (Nov. 18, 2010).

Information from ClinicalTrials.gov archive on the view of NCT00509795 "Vascular Endothelial Growth Factor (VEGF) Trap-Eye: Investigation of Efficacy and Safety in Wet Age-Related Macular Degeneration (AMD)" (Jan. 7, 2011).

Information from ClinicalTrials.gov archive on the view of NCT01012973 Vascular Endothelial Growth Factor (VEGF) Trap-Eye: Investigation of Efficacy and Safety in Central Retinal Vein Occlusion (CRVO)(GALILEO) 7 pages, first posted Nov. 13, 2009; results first posted Nov. 22, 2012; last update posted Nov. 3, 2014; printed Dec. 4, 2019 (<https://clinicaltrials.gov/ct2/show/study/NCT01012973>) (NOTE: May correspond to "Vascular Endothelial Growth Factor Trap&#8208;Eye Investigation of Efficacy and Safety in Central Retinal Vein Occlusion title, 8 pages, Nov. 12, 2009, US [Cited in Third Party Observations filed in parent appli-

cation U.S. Appl. No. 16/055,847]" which was cited in the Third Party Observations dated May 1, 2019).

Karia, Niraj, "Retinal vein occlusion: pathophysiology and treatment options" *Clinical Ophthalmology*, 4:809-816 (2010).

Korobelnik et al., "Intravitreal Aflibercept Injection for Macular Edema Resulting from Central Retinal Vein Occlusion" *American Academy of Ophthalmology* (2014) 121 (1):202-208.

Korobelnik, "Intravitreal Aflibercept for Diabetic Macular Edema." *Ophthalmology*, 121(11):2247-54 (Nov. 2014).

Kuo, "Comparative evaluation of the antitumor activity of antiangiogenic proteins delivered by gene transfer" *PNAS* 98(8):4605-4610 (Apr. 10, 2001).

Krzystolik et al., "Prevention of Experimental Choroidal Neovascularization With Intravitreal Anti-Vascular Endothelial Growth Factor Antibody Fragment" *Arch Ophthalmol.*, 120:338-346 (Mar. 2002).

Lalwani, A Variable-dosing Regimen with Intravitreal Ranibizumab for Neovascular Age-related Macular Degeneration: Year 2 of the PRONTO Study. *Am J Ophthalmology*, 148(1):43-58 (Jul. 2009).

Levine, "Macular Hemorrhage In Neovascular Age-Related Macular Degeneration After Stabilization With Antiangiogenic Therapy." *Retina*, 29(8):1074-79 (2009).

Lucentis Label Title, 7 pages, Jun. 30, 2010 [Cited in Third Party Observations filed in parent application U.S. Appl. No. 16/055,847].

Major et al., "DA VINCI: DME and VEGF Trap-Eye: Investigation of Clinical Impact: Phase 2 Study in Patients with Diabetic Macular Edema (DME)" *ARVO Annual Meeting Abstract* (Apr. 2010).

Margolis, "Hemorrhagic Recurrence Of Neovascular Age-Related Macular Degeneration Not Predicted By Spectral Domain Optical Coherence Tomography." *Retinal Cases & Brief Reports*, 4:1-4 (2010).

Massin, "Safety and Efficacy of Ranibizumab in Diabetic Macular Edema (RESOLVE Study\*)." *Diabetes Care*, 33(11):2399-405 (Nov. 2010).

Mitchell, "The RESTORE Study, Ranibizumab Monotherapy or Combined with Laser versus Laser Monotherapy for Diabetic Macular Edema." *Ophthalmology*, 188(4):615-25 (Apr. 2011).

Mitchell, Edith P. "Targeted Therapy for Metastatic Colorectal Cancer: Role of Aflibercept" *Clinical Colorectal Cancer* (2013) 12(2):73-85.

N/A "Materials from Jun. 2011 FDA Committee Mtg" (Jun. 17, 2011).

N/A "Materials from Dec. 2011 FDA Committee Mtg"(Dec. 1, 2011).

Nguyen et al., "A Phase I Study of Intravitreal Vascular Endothelial Growth Factor Trap-Eye in Patients with Neovascular Age-Related Macular Degeneration" *Ophthalmology*, J.B. Lippincott Co., Philadelphia, PA, US, 116(11):2141-2148 (Nov. 1, 2009).

Nguyen et al., "Randomized, Double-masked, Active-controlled Phase 3 Trial of the Efficacy and Safety of Intravitreal VEGF Trap-Eye in Wet AMD: One-year Results of the VIEW 1 Study" *ARVO Annual Meeting Abstract* (Apr. 2011).

Nguyen, "Ranibizumab for Diabetic Macular Edema, Results from 2 Phase III Randomized Trials: RISE and RIDE." *Ophthalmology*, 119(4):789-801 (Apr. 2012).

Nguyen et al., "Results of a Phase I, Dose-Escalation, Safety, Tolerability, and Bioactivity Study of Intravitreal VEGF Trap in Patients with Neovascular Age-Related Macular Degeneration" *ARVO Annual Meeting Abstract* (May 2006).

Nichols, Earl R., "AAO: Ranibizumab (rhuRab) May Improve Vision in Age-Related Macular Degeneration" *Doctor's Guide Global Edition*, [www.pslgroup.com/dg/23f2aa.htm](http://www.pslgroup.com/dg/23f2aa.htm), pp. 1-2 (Nov. 24, 2003).

Noguera-Troise et al., "Blockade of D114 inhibits tumour growth by promoting non-productive angiogenesis" *Nature* (Dec. 2006) 444:1032-1037.

Ohr, "Aflibercept in wet age-related macular degeneration: a perspective review" *Ther. Adv. Chronic Dis.*, 3(4):153-161 (2012).

Olivera et al., "VEGF Trap R1R2 suppresses experimental corneal angiogenesis" *European Journal of Ophthalmology*, 20(1):48-54 (Jan. 1, 2010).

(56)

**References Cited****OTHER PUBLICATIONS**

Pai et al., "Current concepts in intravitreal drug therapy for diabetic retinopathy" *Saudi Journal of Ophthalmology* 24(4):143-149 (Jun. 30, 2010).

Papadopoulos, "Binding and neutralization of vascular endothelial growth factor (VEGF) and related ligands by VEGF Trap, ranibizumab and bevacizumab" *Angiogenesis*, 15:171-185 (2012).

Regeneron SEC Form 10-K (Feb. 27, 2008).

Regeneron SEC Form 10-K (Feb. 26, 2009).

Regeneron SEC Form 10-K (Feb. 17, 2011).

Regeneron SEC Form 10-Q (May 8, 2006).

Regeneron SEC Form 10-Q (Aug. 8, 2006).

Regeneron SEC Form 10-Q (Nov. 6, 2006).

Regeneron SEC Form 10-Q (May 4, 2007).

Regeneron SEC Form 10-Q (Aug. 3, 2007).

Regeneron SEC Form 10-Q (Apr. 30, 2009).

Regeneron SEC Form 10-Q (Nov. 3, 2009).

Regeneron SEC Form 10-Q (Apr. 29, 2010).

Regeneron SEC Form 10-Q (Jul. 28, 2010).

Regeneron SEC Form 10-Q (Oct. 28, 2010).

Regeneron SEC Form 10-Q (May 3, 2011).

Regeneron SEC Form 10-Q (Jul. 28, 2011).

Regeneron SEC Form 10-Q (Oct. 27, 2011).

Regeneron SEC Form 8-K Exhibit: "Press Release of Regeneron Pharmaceuticals, Inc. dated May 1, 2006" (May 2, 2006).

Regeneron SEC Form 8-K Exhibit: "Press Release of Regeneron Pharmaceuticals, Inc. dated May 3, 2006" (May 5, 2006).

Regeneron SEC Form 8-K Exhibit: "Slides presented at the Company's 2006 Annual Meeting of Shareholders held on Jun. 9, 2006" (Jun. 9, 2006).

Regeneron SEC Form 8-K Exhibit: "Press Release dated May 2, 2007" (May 3, 2007).

Regeneron SEC Form 8-K Exhibit: "Overheads for presentation at Regeneron's Annual Meeting of Shareholders to be held on Jun. 8, 2007" (Jun. 8, 2007).

Regeneron SEC Form 8-K Exhibit: "Press Release dated Oct. 1, 2007" (Oct. 1, 2007).

Regeneron SEC Form 8-K Exhibit: "Press Release dated Nov. 6, 2007" (Nov. 6, 2007).

Regeneron SEC Form 8-K Exhibit: "Press Release dated May 1, 2008" (May 2, 2008).

Regeneron SEC Form 8-K Exhibit: "Press Release dated Nov. 4, 2008" (Nov. 4, 2008).

Regeneron SEC Form 8-K Exhibit: "99(a) Slides that Regeneron Pharmaceuticals, Inc. intends to use in conjunction with meetings with investors at the J.P. Morgan 27th Annual Healthcare Conference in San Francisco on Jan. 12-15, 2009." (Jan. 9, 2009).

Regeneron SEC Form 8-K Exhibit: "Press Release dated Apr. 30, 2009" (May 1, 2009).

Regeneron SEC Form 8-K Exhibit: "Press Release dated Nov. 3, 2009." (Nov. 4, 2009).

Regeneron SEC Form 8-K Exhibit: "Press Release Reporting Positive Results for VEGF Trap-Eye in Phase 3 Study in Central Retinal Vein Occlusion (CRVO) and in Phase 2 Study in Diabetic Macular Edema (DME) dated Dec. 20, 2010." (Dec. 20, 2010).

Regeneron SEC Form 8-K Exhibit: "Press Release dated Feb. 17, 2011" (Feb. 18, 2011).

Regeneron SEC Form 8-K Exhibit: "Press Release Reporting Positive Results for VEGF Trap-Eye in Second Phase 3 Study in Central Retinal Vein Occlusion, dated Apr. 27, 2011" (Apr. 27, 2011).

Regeneron SEC Form 8-K Exhibit: "Press Release dated May 3, 2011." (May 3, 2011).

Regeneron SEC Form 8-K Exhibit: "Press Release, dated Jun. 17, 2011, Announcing that EYLEA™ (afibercept ophthalmic solution) Received Unanimous Recommendation for Approval for Treatment of Wet AMD from FDA Advisory Committee." (Jun. 21, 2011).

Regeneron SEC Form 8-K Exhibit: "Presentation entitled VEGF Trap-Eye in CRVO: 1-year Results of the Phase 3 COPERNICUS Study" (Aug. 22, 2011).

Regeneron SEC Form 8-K Exhibit: "Press Release Announcing FDA Approval of EYLEA™ (afibercept) Injection for the Treatment of Wet Age-Related Macular Degeneration, dated Nov. 18, 2011" (Nov. 21, 2011).

Regeneron Press Release "Positive Interim Phase 2 Data Reported For VEGF Trap-Eye In Age-Related Macular Degeneration" (Mar. 27, 2007).

Regeneron Press Release "VEGF TRAP-Eye Phase 2 Wet AMD Results Reported At Arvo Annual Meeting" (May 9, 2007).

Regeneron Press Release "Regeneron Reports Second Quarter Financial And Operating Results" (Aug. 1, 2007).

Regeneron Pharmaceuticals, Inc., "Regeneron and Bayer Healthcare Initiate Phase 3 Global Development Program for VEGF Trap-Eye In Wet Age-Related Macular Degeneration (AMD)" (Aug. 2, 2007).

Regeneron Pharmaceuticals, Inc. FORM 10-Q, published on Nov. 7, 2007 for the period ending Sep. 30, 2007.

Regeneron Press Release "Regeneron Announces Positive Primary Endpoint Results From A Phase 2 Study Of VEGF Trap-Eye In Age-Related Macular Degeneration" (Oct. 1, 2007).

Regeneron Press Release "Regeneron Reports Fourth Quarter And Full Year 2007 Financial And Operating Results" (Feb. 27, 2008).

Regeneron Pharmaceuticals, Inc., "Regeneron and Bayer Healthcare Announce Encouraging 32-Week Follow-up Results from a Phase 2 Study of VEGF Trap-Eye in Age-Related Macular Degeneration" (Apr. 28, 2008).

Regeneron, Press release "Regeneron Reports First Quarter 2008 Financial and Operating Results", May 1, 2008.

Regeneron Press Release, "Bayer and Regeneron Dose First Patient in Second Phase 3 Study for VEGF Trap-Eye in Wet Age-Related Macular Degeneration." May 8, 2008.

Regeneron Pharmaceuticals, Inc., "Regeneron and Bayer Healthcare Announce VEGF Trap-Eye Achieved Durable Improvement in Vision over 52 Weeks in a Phase 2 Study in Patients with Age-related Macular Degeneration" (Aug. 19, 2008).

Regeneron Pharmaceuticals Inc., "VIEW 1 Vascular Endothelial Growth Factor (VEGF) Trap-Eye 1-Year Results: Investigation of Efficacy and Safety in Wet Age-Related Macular Degeneration (AMD)" presented at Bascom Palmer Eye Institute's Angiogenesis, Exudation and Degeneration 2011 meeting in Miami, Florida (Feb. 12, 2011).

Regeneron Pharmaceuticals Inc., "VIEW 2 Vascular Endothelial Growth Factor (VEGF) Trap-Eye 1-Year Results: Investigation of Efficacy and Safety in Wet Age-Related Macular Degeneration (AMD)" presented at Bascom Palmer Eye Institute's Angiogenesis, Exudation and Degeneration 2011 meeting in Miami, Florida (Feb. 12, 2011).

Regeneron Pharmaceuticals Inc., "VEGF Trap-Eye CLEAR-IT 2 Final Primary Endpoint Results" presented at the 2007 Retina Society Conference in Boston, Massachusetts (Sep. 30, 2007).

Regeneron Pharmaceuticals Inc., "VEGF Trap-Eye Final Phase 2 Results in Age-related Macular Degeneration Presented at 2008 Retina Society Meeting" (Sep. 28, 2008) (XP-002770952).

Regeneron 2008 Annual Report.

Regeneron Pharmaceuticals, Inc. "Regeneron Reports Full Year and Fourth Quarter 2008 Financial and Operating Results" (Feb. 26, 2009).

Regeneron Pharmaceuticals, Inc. "Bayer and Regeneron Extend Development Program for VEGF Trap-Eye to Include Central Retinal Vein Occlusion" (Apr. 30, 2009).

Regeneron Press Release "First Patient Enrolled In Regeneron And Bayer Healthcare VEGF Trap-Eye Phase 3 Program In Central Retinal Vein Occlusion" (Jul. 23, 2009).

Regeneron Press Release "Enrollment Completed in Regeneron and Bayer Healthcare Phase 3 Studies of VEGF Trap-Eye in Neovascular Age-Related Macular Degeneration (Wet AMD)" Sep. 14, 2009.

Regeneron 2009 Annual Report and 10-K.

Regeneron Press Release, "VEGF Trap-Eye Shows Positive Results in a Phase 2 Study in Patients With Diabetic Macular Edema." Feb. 18, 2010.

(56)

**References Cited****OTHER PUBLICATIONS**

- Regeneron Press Release "Regeneron Schedules Nov. 22, 2010 Teleconference And Webcast To Discuss Results Of Two Phase 3 Studies With VEGF Trap-Eye In Wet Age-Related Macular Degeneration" (Nov. 19, 2010).
- Regeneron Press Release "Bayer and Regeneron Report Positive Top-Line Results of Two Phase 3 Studies with VEGF Trap-Eye in Wet Age-related Macular Degeneration" Nov. 22, 2010.
- Regeneron Press Release "Regeneron and Bayer Report Positive Results for VEGF Trap-Eye in Phase 3 Study in Central Retinal Vein Occlusion (CRVO) and in Phase 2 Study in Diabetic Macular Edema (DME)" Dec. 20, 2010.
- Regeneron 2010 Annual Report and 10-K.
- Regeneron Press Release "Regeneron And Bayer Start Phase 3 Trial To Extend Ophthalmology Research & Development Program For VEGF Trap-Eye In Asia" (Jan. 18, 2011).
- Regeneron Press Release "Regeneron To Webcast Investor Briefing On VEGF Trap-Eye Clinical Program On Sunday, Feb. 13 At 9 AM ET" (Feb. 9, 2011).
- Regeneron Press Release "Regeneron Submits Biologies License Application To FDA For VEGF Trap-Eye For Treatment Of Wet Age-Related Macular Degeneration" (Feb. 22, 2011).
- Regeneron Press Release "Regeneron And Bayer Announce Start Of Phase 3 Clinical Program In Diabetic Macular Edema" (Apr. 8, 2011).
- Regeneron Pharmaceuticals, Inc., "FDA Grants Priority Review for VEGF Trap-Eye for the Treatment of Wet Age-Related Macular Degeneration" (Apr. 18, 2011).
- Regeneron Press Release VEGF Trap-Eye Submitted for EU Marketing Authorization for Treatment of Wet Age-Related Macular Degeneration (Jun. 7, 2011).
- Regeneron Pharmaceuticals, Inc., "Regeneron Announces EYLEA™ (aflibercept ophthalmic solution) Receives Unanimous Recommendation for Approval for Treatment of Wet AMD from FDA Advisory Committee" (Jun. 17, 2011).
- Regeneron Press Release "Regeneron Announces Clinical Presentations at ASRS 2011 Annual Meeting" (Aug. 17, 2011).
- Regeneron Pharmaceuticals, Inc., "Regeneron Announces FDA Approval of EYLEA™ (aflibercept) Injection for the Treatment of Wet Age-Related Macular Degeneration: CORRECTED" (Nov. 18, 2011).
- Regeneron Pharmaceuticals, Inc., "Regeneron and Bayer Initiate Phase 3 Clinical Program for the Treatment of Wet Age-Related Macular Degeneration in China" (Nov. 28, 2011).
- Regeneron Pharmaceuticals, Inc., "Two Year Results of Phase 3 Studies with EYLEA™ (aflibercept) Injection in wet AMD Show Sustained Improvement in Visual Acuity" (Dec. 5, 2011).
- Rosenfeld, "Ranibizumab for Neovascular Age-Related Macular Degeneration." *N Engl J Med*, 355(14):1419-31 (Oct. 5, 2006).
- Rosenfeld, "Lessons Learned From Avastin and OCT—The Good, the Bad, and the Ugly: The LXXV Edward Jackson Memorial Lecture." *Am. J. Ophthalmology*, 164:26-45 (Aug. 2010).
- Rudge et al., "VEGF Trap complex formation measures production rates of VEGF, providing a biomarker for predicting efficacious angiogenic blockade" *PNAS* (Nov. 20, 2007) 104(47):18363-18370.
- Rudge et al. "Clinical Development of VEGF Trap" In: Figg W.D., Folkman J. (eds) *Angiogenesis* (2008).
- Schmidt-Erfurth, "Efficacy and Safety of Monthly versus Quarterly Ranibizumab Treatment in Neovascular Age-related Macular Degeneration: The EXCIE Study" *Ophthalmology*, 118(5):831-839 (2010).
- Schmidt-Erfurth et al. "Primary Results of an International Phase III Study Using Intravitreal VEGF Trap-Eye Compared to Ranibizumab in Patients with Wet AMD (VIEW 2)" *ARVO Annual Meeting Abstract* (Apr. 2011).
- Schmidt-Erfurth, "Three-Year Outcomes of Individualized Ranibizumab Treatment in Patients with Diabetic Macular Edema." *Ophthalmology*, 121(5):1045-53, (May 2014).
- Schmidt-Erfurth et al., "Intravitreal Aflibercept Injection for Neovascular Age-related Macular Degeneration" *Ophthalmology* (2014) 121:193-201.
- Schnichels, "Comparative toxicity and proliferation testing of aflibercept, bevacizumab and ranibizumab on different ocular cells." *Br. J. Ophthalmol.*, 97:917-923 (2013).
- Semeraro et al., "Aflibercept in wet AMD: specific role and optimal use" *Drug Design, Development and Therapy* (Aug. 2, 2013) 7:711-722.
- Sharma and S. and Kaiser, P. K., Update on VEGF TRAP-Eye Clinical Trials and Retinal. Physician, pp. 1-6 (Nov./Dec. 2010) <URL: <https://www.retinalphysician.com/issues/2010/nov-dec/update-on-vegf-trap-eye-clinical-trials>>.
- Slakter et al., "Influence of Baseline Angiographic Classification on Outcomes in the CLEAR-IT 2 Phase 2 Study of Intravitreal VEGF Trap-Eye in Neovascular Age-Related Macular Degeneration" *ARVO Annual Meeting Abstract* (Apr. 2010).
- Slakter et al., "A Phase 2, Randomized, Controlled Dose-and Interval-Ranging Study of Intravitreal VEGF Trap-Eye in Patients with Neovascular Age-Related Macular Degeneration: Optical Coherence Tomography (OCT) and Fluorescein Angiography (FA) Outcomes at 1 Year" *ARVO Annual Meeting Abstract* (Apr. 2009).
- Spaide, "Ranibizumab According to Need: A Treatment for Age-related Macular Degeneration." *Am J Ophthalmology*, 143(4):679-680 (Apr. 2007).
- Stewart, "The expanding role of vascular endothelial growth factor inhibitors in ophthalmology" *Mayo Clin Proc.* 87(1):77-88 (Jan. 2012).
- Stewart et al., "Predicted biological activity of intravitreal VEGF Trap" *British Journal of Ophthalmology*, 92(5):667-668 (2008).
- Stewart, "Aflibercept" *Nature Reviews: Drug Discovery* 11:269-270 (Apr. 1, 2012).
- Tannock et al., "Aflibercept versus placebo in combination with docetaxel and prednisone for treatment of men with metastatic castration-resistant prostate cancer (VENICE): a phase 3, double-blind randomized trial" *Lancet Oncol* (2013) 14:760-768.
- Thomas Reuters Integrity "VEGF Trap-Eye final phase II results in age-related macular degeneration presented at 2008 Retina Society Meeting" (Sep. 28, 2008).
- Thurston, Gavin "Complementary actions of VEGF and Angiopoietin-1 on blood vessel growth and leakage" *J. Anat.* (2002) 200:575-580.
- Thurston, "Vascular endothelial growth factor and other signaling pathways in developmental and pathologic angiogenesis." *International Journal of Hematology*, 80:7-20 (2004).
- Updated Information from ClinicalTrials.gov archive History of Changes for Study: NCT01012973 Vascular Endothelial Growth Factor (VEGF) Trap-Eye: Investigation of Efficacy and Safety in Central Retinal Vein Occlusion (CRVO)(GALILEO) 38 pages, Latest version submitted Oct. 27, 2014 on ClinicalTrials.gov (NCT01012973\_01182013\_27424.1).
- Updated Information from ClinicalTrials.gov archive History of Changes for Study: NCT01012973 Vascular Endothelial Growth Factor (VEGF) Trap-Eye: Investigation of Efficacy and Safety in Central Retinal Vein Occlusion (CRVO)(GALILEO) 10 pages, Latest version submitted Oct. 27, 2014 on ClinicalTrials.gov (NCT01012973\_01252011\_27433.1).
- Updated Information from ClinicalTrials.gov archive History of Changes for Study: NCT01012973 Vascular Endothelial Growth Factor (VEGF) Trap-Eye: Investigation of Efficacy and Safety in Central Retinal Vein Occlusion (CRVO)(GALILEO) 11 pages, Latest version submitted Oct. 27, 2014 on ClinicalTrials.gov (NCT01012973\_01262012\_27428.1).
- Updated Information from ClinicalTrials.gov archive History of Changes for Study: NCT01012973 Vascular Endothelial Growth Factor (VEGF) Trap-Eye: Investigation of Efficacy and Safety in Central Retinal Vein Occlusion (CRVO)(GALILEO) 38 pages, Latest version submitted Oct. 27, 2014 on ClinicalTrials.gov (NCT01012973\_01302013\_27423.1).
- Updated Information from ClinicalTrials.gov archive History of Changes for Study: NCT01012973 Vascular Endothelial Growth Factor (VEGF) Trap-Eye: Investigation of Efficacy and Safety in Central Retinal Vein Occlusion (CRVO)(GALILEO) 12 pages, Latest version submitted Oct. 27, 2014 on ClinicalTrials.gov (NCT01012973\_02092010\_27442.1).
- Updated Information from ClinicalTrials.gov archive History of Changes for Study: NCT01012973 Vascular Endothelial Growth

(56)

**References Cited****OTHER PUBLICATIONS**

Factor (VEGF) Trap-Eye: Investigation of Efficacy and Safety in Central Retinal Vein Occlusion (CRVO)(GALILEO) 11 pages, Latest version submitted Oct. 27, 2014 on ClinicalTrials.gov (NCT01012973\_02202012\_27427.1).

Updated Information from ClinicalTrials.gov archive History of Changes for Study: NCT01012973 Vascular Endothelial Growth Factor (VEGF) Trap-Eye: Investigation of Efficacy and Safety in Central Retinal Vein Occlusion (CRVO)(GALILEO) 12 pages, Latest version submitted Oct. 27, 2014 on ClinicalTrials.gov (NCT01012973\_03162010\_27441.1).

Updated Information from ClinicalTrials.gov archive History of Changes for Study: NCT01012973 Vascular Endothelial Growth Factor (VEGF) Trap-Eye: Investigation of Efficacy and Safety in Central Retinal Vein Occlusion (CRVO)(GALILEO) 10 pages, Latest version submitted Oct. 27, 2014 on ClinicalTrials.gov (NCT01012973\_04082011\_27432.1).

Updated Information from ClinicalTrials.gov archive History of Changes for Study: NCT01012973 Vascular Endothelial Growth Factor (VEGF) Trap-Eye: Investigation of Efficacy and Safety in Central Retinal Vein Occlusion (CRVO)(GALILEO) 12 pages, Latest version submitted Oct. 27, 2014 on ClinicalTrials.gov (NCT01012973\_04162010\_27440.1).

Updated Information from ClinicalTrials.gov archive History of Changes for Study: NCT01012973 Vascular Endothelial Growth Factor (VEGF) Trap-Eye: Investigation of Efficacy and Safety in Central Retinal Vein Occlusion (CRVO)(GALILEO) 10 pages, Latest version submitted Oct. 27, 2014 on ClinicalTrials.gov (NCT01012973\_06232011\_27431.1).

Updated Information from ClinicalTrials.gov archive History of Changes for Study: NCT01012973 Vascular Endothelial Growth Factor (VEGF) Trap-Eye: Investigation of Efficacy and Safety in Central Retinal Vein Occlusion (CRVO)(GALILEO) 12 pages, Latest version submitted Oct. 27, 2014 on ClinicalTrials.gov (NCT01012973\_07222010\_27439.1).

Updated Information from ClinicalTrials.gov archive History of Changes for Study: NCT01012973 Vascular Endothelial Growth Factor (VEGF) Trap-Eye: Investigation of Efficacy and Safety in Central Retinal Vein Occlusion (CRVO)(GALILEO) 12 pages, Latest version submitted Oct. 27, 2014 on ClinicalTrials.gov (NCT01012973\_08252010\_27438.1).

Updated Information from ClinicalTrials.gov archive History of Changes for Study: NCT01012973 Vascular Endothelial Growth Factor (VEGF) Trap-Eye: Investigation of Efficacy and Safety in Central Retinal Vein Occlusion (CRVO)(GALILEO) 10 pages, Latest version submitted Oct. 27, 2014 on ClinicalTrials.gov (NCT01012973\_08262010\_27437.1).

Updated Information from ClinicalTrials.gov archive History of Changes for Study: NCT01012973 Vascular Endothelial Growth Factor (VEGF) Trap-Eye: Investigation of Efficacy and Safety in Central Retinal Vein Occlusion (CRVO)(GALILEO) 10 pages, Latest version submitted Oct. 27, 2014 on ClinicalTrials.gov (NCT01012973\_09082010\_27436.1).

Updated Information from ClinicalTrials.gov archive History of Changes for Study: NCT01012973 Vascular Endothelial Growth Factor (VEGF) Trap-Eye: Investigation of Efficacy and Safety in Central Retinal Vein Occlusion (CRVO)(GALILEO) 10 pages, Latest version submitted Oct. 27, 2014 on ClinicalTrials.gov (NCT01012973\_09192011\_27430.1).

Updated Information from ClinicalTrials.gov archive History of Changes for Study: NCT01012973 Vascular Endothelial Growth Factor (VEGF) Trap-Eye: Investigation of Efficacy and Safety in Central Retinal Vein Occlusion (CRVO)(GALILEO) 10 pages, Latest version submitted Oct. 27, 2014 on ClinicalTrials.gov (NCT01012973\_10042010\_27435.1).

Updated Information from ClinicalTrials.gov archive History of Changes for Study: NCT01012973 Vascular Endothelial Growth Factor (VEGF) Trap-Eye: Investigation of Efficacy and Safety in Central Retinal Vein Occlusion (CRVO)(GALILEO) 38 pages,

Latest version submitted Oct. 27, 2014 on ClinicalTrials.gov (NCT01012973\_10232012\_27426.1).

Updated Information from ClinicalTrials.gov archive History of Changes for Study: NCT01012973 Vascular Endothelial Growth Factor (VEGF) Trap-Eye: Investigation of Efficacy and Safety in Central Retinal Vein Occlusion (CRVO)(GALILEO) 38 pages, Latest version submitted Oct. 27, 2014 on ClinicalTrials.gov (NCT01012973\_10272013\_27422.1).

Updated Information from ClinicalTrials.gov archive History of Changes for Study: NCT01012973 Vascular Endothelial Growth Factor (VEGF) Trap-Eye: Investigation of Efficacy and Safety in Central Retinal Vein Occlusion (CRVO)(GALILEO) 10 pages, Latest version submitted Oct. 27, 2014 on ClinicalTrials.gov (NCT01012973\_11012010\_27434.1).

Updated Information from ClinicalTrials.gov archive History of Changes for Study: NCT01012973 Vascular Endothelial Growth Factor (VEGF) Trap-Eye: Investigation of Efficacy and Safety in Central Retinal Vein Occlusion (CRVO)(GALILEO) 12 pages, Latest version submitted Oct. 27, 2014 on ClinicalTrials.gov (NCT01012973\_11132009\_27444.1).

Updated Information from ClinicalTrials.gov archive History of Changes for Study: NCT01012973 Vascular Endothelial Growth Factor (VEGF) Trap-Eye: Investigation of Efficacy and Safety in Central Retinal Vein Occlusion (CRVO)(GALILEO) 10 pages, Latest version submitted Oct. 27, 2014 on ClinicalTrials.gov (NCT01012973\_11292011\_27429.1).

Updated Information from ClinicalTrials.gov archive History of Changes for Study: NCT01012973 Vascular Endothelial Growth Factor (VEGF) Trap-Eye: Investigation of Efficacy and Safety in Central Retinal Vein Occlusion (CRVO)(GALILEO) 38 pages, Latest version submitted Oct. 27, 2014 on ClinicalTrials.gov (NCT01012973\_12182012\_27425.1).

Updated Information from ClinicalTrials.gov archive History of Changes for Study: NCT01012973 Vascular Endothelial Growth Factor (VEGF) Trap-Eye: Investigation of Efficacy and Safety in Central Retinal Vein Occlusion (CRVO)(GALILEO) 12 pages, Latest version submitted Oct. 27, 2014 on ClinicalTrials.gov (NCT01012973\_12212010\_27443.1).

Vascular Endothelial Growth Factor Trap&#8208;Eye Investigation of Efficacy and Safety in Central Retinal Vein Occlusion title, 8 pages, Nov. 12, 2009, US [Cited in Third Party Observations filed in parent application U.S. Appl. No. 16/055,847] Note: May correspond to "Information from ClinicalTrials.gov archive on the view of NCT01012973" Vascular Endothelial Growth Factor (VEGF) Trap-Eye: Investigation of Efficacy and Safety in Central Retinal Vein Occlusion (CRVO)(GALILEO) 7 pages, first posted Nov. 13, 2009; results first posted Nov. 22, 2012; last update posted Nov. 3, 2014; printed Dec. 4, 2019 (<https://clinicaltrials.gov/ct2/show/study/NCT01012973>) Office Action dated Dec. 10, 19 in U.S. Appl. No. 16/055,847.

Wachsberger, "Vegf trap in combination with radiotherapy improves tumor control in u87 glioblastoma." *Int. J. Radiation Oncology Biol Phys.* 67(5):1526-1537 (2007).

Wolfson, "Regeneron Focuses on Age-Related Macular Degeneration." *Chemistry & Biology* 15:303-304 (Apr. 2008).

Xia et al., "Transgenic delivery of VEGF to mouse skin leads to an inflammatory condition resembling human psoriasis" *Blood* (Jul. 1, 2003) 102(1):161 -168.

Yancopoulos, "Vascular-specific growth factors and blood vessel formation." *Nature* 407:242-48 (Sep. 14, 2000).

Yancopoulos, "Clinical Application of Therapies Targeting VEGF." *Cell* 143:13-16 (Oct. 1, 2010).

Yung, "Moving Toward the Next Steps in Angiogenesis Therapy?" *Society for Neuro-Oncology*, 10:939 (2008).

Eylea®, Highlights of Prescribing Information, Revised Nov. 2011. IPR2021-00880, Paper 1, Petition for IPR (May 5, 2021).

IPR2021-00880, Exhibit 1002, Albini Declaration (May 4, 2021).

IPR2021-00880, Exhibit 1003, Gerritsen Declaration (Apr. 30, 2021).

IPR2021-00880, Paper 10, Preliminary Response of Patent Owner (Aug. 16, 2021).

IPR2021-00881, Paper 1, Petition for IPR (May 5, 2021).

IPR2021-00881, Exhibit 1002, Albini Declaration (May 4, 2021).

(56)

**References Cited****OTHER PUBLICATIONS**

- IPR2021-00881, Exhibit 1003, Gerritsen Declaration (Apr. 26, 2021).
- IPR2021-00881, Paper 10, Preliminary Response of Patent Owner (Aug. 16, 2021).
- IPR2021-00881, Exhibit 2001, Do Declaration (Aug. 13, 2021).
- Mitchell et al., "Evaluating the Impact of Intravitreal Aflibercept on Diabetic Retinopathy Progression in the VIVID-DME and VISTA-DME Studies" *Ophthalmol Retina* 2(10):988-96 (2018).
- PGR2021-00035, Paper 2, Petition for PGR (Jan. 7, 2021).
- PGR2021-00035, Paper 6, Preliminary Response of Patent Owner (Apr. 15, 2021).
- PGR2021-00035, Exhibit 1003 Wu Declaration (Jan. 7, 2021).
- PGR2021-00035, Exhibit 2001 Do Declaration (Apr. 21, 2021).
- PGR2021-00035, Exhibit 2002 D. Brown Declaration (Apr. 14, 2021).
- Cao, J. R., R.; Wang, Q.; Yancopoulos, G.D.; Wiegand, S.J. (2002). Inhibition of Corneal Neovascularization and Inflammation by VEGF Trap. In "ARVO", Invest. Ophthalmol. Vis. Sci. vol. 43. E-Abstract 1863.
- Wang, Q. R., R.; Cao, J.; Yancopoulos, G.D.; and Wiegand, S.J. (2002). Anti-Angiogenic Properties of a New VEGF Antagonist, VEGF Trap, in a Mouse Model of Retinal Neovascularization. In "ARVO", Invest. Ophthalmol. Vis. Sci., vol. 43. E-Abstract. 3714.
- Saishin, Y., Saishin, Y., Takahashi, K., Lima e Silva, R., et al. (2003). VEGF-TRAP(R1R2) suppresses choroidal neovascularization and VEGF-induced breakdown of the blood-retinal barrier. *J Cell Physiol* 195:241-48.
- Cursiefen, C., Cao, J., Chen, L., Liu, Y., Maruyama, K., et al. (2004). Inhibition of hemangiogenesis and lymphangiogenesis after normal-risk corneal transplantation by neutralizing VEGF promotes graft survival. *Invest Ophthalmol Vis Sci* 45(8):2666-73.
- Cursiefen, C., Chen, L., Borges, L. P., Jackson, D., Cao, J., et al. (2004). VEGF-A stimulates lymphangiogenesis and hemangiogenesis in inflammatory neovascularization via macrophage recruitment. *J Clin Invest* 113(7):1040-50.
- Cao, J.; Song, H.; Renard, R.A.; Liu, Y.; Yancopoulos, G.D.; Wiegand, S.J. (2005). Systemic Administration of VEGF Trap Suppresses Vascular Leak and Leukostasis in the Retinas of Diabetic Rats. In "ARVO", vol. 46. Invest. Ophthalmol. Vis. Sci. E-Abstract 446.
- Nork, T. M., Dubielzig, R. R., Christian, B. J., Miller, P. E., Miller, J. M., et al. (2011). Prevention of experimental choroidal neovascularization and resolution of active lesions by VEGF trap in nonhuman primates. *Arch Ophthalmol* 129(8):1042-52.
- U.S. Pat. No. 7,374,758—Patent Term Extension Application submitted Dec. 22, 2011.
- Anderson et al., "Delivery of Anti-Angiogenic Molecular Therapies for Retinal Disease" *Drug Discovery Today* 15: 272 (2010).
- AVASTIN® label.
- Avery, R. L., D. J. Pieramici, M. D. Rabena, A. A. Castellarin, M. A. Nasir and M. J. Giust, "Intravitreal bevacizumab (Avastin) for neovascular age-related macular degeneration" *Ophthalmology* 113(3): 363-372 e365 (2006).
- Bashshur et al., "Intravitreal Bevacizumab for the Management of Choroidal Neovascularization in Age-Related Macular Degeneration" *Am J. Ophthalmology* 142: 1 (2006).
- Bontempo, "Preformulation Development of Parental Biopharmaceuticals" *Drugs and the Pharmaceutical Sciences* 85:91-108 (1997).
- Bressler, N. M. and G. Treatment of Age-Related Macular Degeneration with Photodynamic Therapy Study, "Photodynamic therapy of subfoveal choroidal neovascularization in age-related macular degeneration with verteporfin: two-year results of 2 randomized clinical trials—tap report 2." *Arch Ophthalmol* 119(2): 198-207 (2001)
- Brown & Regillo, "Anti-VEGF Agents in the Treatment of Neovascular Age-Related Macular Degeneration: Applying Clinical Trial Results to the Treatment of Everyday Patients" *Am J. Ophthalmology* 144: 627 (2007).
- Chi et al., "Physical Stability of Proteins in Aqueous Solution: Mechanism and Driving Forces in Nonnative Protein Aggregation" *Pharmaceutical Research* vol. 20, No. 9, 1325-1336 (Sep. 2003).
- Ciulla & Rosenfeld, "Antivascular Endothelial Growth Factor Therapy For Neovascular Age-Related Macular Degeneration" *Current Opinion Ophthalmology* 20: 158 (2009).
- Department of Health and Human Services, Office of Inspector General, "Questionable Billing for Medicare Ophthalmology Services" Sep. 2015 OEI-04-12-00280.
- Eylea® Prescribing Information, Revised May 2019.
- Ferrara, N. & Kerbel, R., "Angiogenesis as a Therapeutic Target" *Nature* 438: 967 (2005).
- Heier, J. S., P. A. Campochiaro, L. Yau, Z. Li, N. Saroj, R. G. Rubio and P. Lai "Ranibizumab for macular edema due to retinal vein occlusions: long-term follow-up in the HORIZON trial" *Ophthalmology* 119(4): 802-809 (2012).
- HERCEPTIN® label.
- Holz et al., "VEGF Trap-Eye for Macular Oedema Secondary to Central Retinal Vein Occlusion: 6-Month Results of the Phase III GALILEO Study" *British J. Ophthalmology* 97: 278 (2013).
- Ip, M. S., I. U. Scott, P. C. VanVeldhuisen, N. L. Oden, B. A. Blodi, M. Fisher, L. J. Singerman, M. Tolentino, C. K. Chan, V. H. Gonzalez and S. S. R. Group "A randomized trial comparing the efficacy and safety of intravitreal triamcinolone with observation to treat vision loss associated with macular edema secondary to central retinal vein occlusion: the Standard Care vs Corticosteroid for Retinal Vein Occlusion (SCORE) study report 5" *Arch Ophthalmol* 127(9): 1101-1114 (2009).
- Keane et al., "Effect of Ranibizumab Retreatment Frequency on Neurosensory Retinal Volume in Neovascular AMD" *Retina* 29: 592 (2009).
- Kim et al., "Potent VEGF Blockade Causes Regression of Coopted Vessels in a Model of Neuroblastoma" *Proc. Nat'l Acad. Sci.* 99: 11399 (2002).
- LUCENTIS Approval (2006).
- LUCENTIS® Label (14 pages).
- LUCENTIS® Prescribing Information (2006).
- Macular Photocoagulation Study, G., "Laser photocoagulation of subfoveal neovascular lesions in age-related macular degeneration. Results of a randomized clinical trial. Macular Photocoagulation Study Group" *Arch Ophthalmol* 109(9): 1220-1231 (1991).
- Mitchell et al., "Ranibizumab (Lucentis) in Neovascular Age-Related Macular Degeneration: Evidence from Clinical Trials" *Brit. J. Ophthalmology* 94: 2 (2009).
- Ni & Hui, "Emerging Pharmacologic Therapies for Wet Age-Related Macular Degeneration" *Ophthalmologica* 223: 401 (2009).
- Parkins & Lashmar, "The formulation of biopharmaceutical products" *Pharmaceutical Science & Technology Today* vol. 3, No. 4: 129-137 (Apr. 4, 2000).
- RAPTIVA® label.
- Regeneron Pharmaceuticals Inc. Regeneron Receives \$20 Million Milestone Payment for Initiation of Phase 3 Study of VEGF Trap-Eye in Wet AMD. Media Release: Aug. 14, 2007. Available from URL: <http://www.regeneron.com>.
- Regeneron Pharmaceuticals Inc. Regeneron Reports Fourth Quarter and Full Year 2004 Financial and Operating Results. Media Release: Feb. 22, 2005. Available from URL: <http://www.regeneron.com>.
- Regeneron Pharmaceuticals Inc. Regeneron Reports Fourth Quarter and Full Year 2005 Financial and Operating Results. Media Release: Feb. 24, 2006. Available from URL: <http://regeneron.com>.
- Regeneron Pharmaceuticals Inc. Regeneron Reports Positive Phase Data for the VEGF Trap in Age-Related Macular Degeneration; Preliminary Results Show Improvements in Vision and Reginal Swelling; VEGF Trap Was Well Tolerated at All Dose Levels. Media Release: May 1, 2006. Available from URL: <http://www.regeneron.com>.
- Reichert, "Antibody-Based Therapeutics To Watch In 2011" *MABS* 3: 76 (2011).
- REMICADE® label.
- Rogers et al., "The prevalence of retinal vein occlusion: pooled data from population studies from the United States, Europe, Asia, and Australia" *Ophthalmology* 117(2): 313-319 e311 (2010)

(56)

## References Cited

## OTHER PUBLICATIONS

- Rosenfeld, P. J., A. A. Moshfeghi and C. A. Puliafito, "Optical coherence tomography findings after an intravitreal injection of bevacizumab (avastin) for neovascular age-related macular degeneration" *Ophthalmic Surg Lasers Imaging* 36(4): 331-335 (2005).
- Schmidt-Erfurth "Current Concepts in the Management of Diabetic Macular Edema" *Proceedings* 7:52 (2010).
- Scott et al., "A randomized trial comparing the efficacy and safety of intravitreal triamcinolone with standard care to treat vision loss associated with macular edema secondary to branch retinal vein occlusion: the Standard Care vs Corticosteroid for Retinal Vein Occlusion (SCORE) study report 6" *Arch Ophthalmol* 127(9): 1115-1128 (2009).
- SIMULECT® label.
- Spaide et al., "Prospective Study of Intravitreal Ranibizumab as a Treatment for Decreased Visual Acuity Secondary to Central Retinal Vein Occlusion" *Am J. Ophthalmology* 147: 298 (2009).
- Spielberg, L. & Leys, A., "Intravitreal Bevacizumab for Myopic Choroidal Neovascularization: Short-Term and 1-Year Results" *Bulletin Societe Belge D'Ophthalmologie* 312: 17 (2009).
- Steinbrook, "The Price of Sight—Ranibizumab, Bevacizumab, and the Treatment of Macular Degeneration" *N. Eng. J. Med.* 355:1409 (2006).
- The Branch Vein Occlusion Study, G., "Argon laser photocoagulation for macular edema in branch vein occlusion" *Am J Ophthalmol* 98(3): 271-282 (1984).
- U.S. Dep't Health & Human Servs., Nat'l Inst. Health, Nat'l Eye Inst., "Age-Related Macular Degeneration: What You Should Know" (Sep. 2015) [https://www.nei.nih.gov/sites/default/files/healthpdfs/WYSK\\_AMD\\_English\\_Sep2015\\_PRINT.pdf](https://www.nei.nih.gov/sites/default/files/healthpdfs/WYSK_AMD_English_Sep2015_PRINT.pdf)
- U.S. Dep't Health & Human Servs., Nat'l Inst. Health, Nat'l Eye Inst., "Diabetic Retinopathy: What You Should Know" (Sep. 2015), <https://www.nei.nih.gov/sites/default/files/2019-06/Diabetic-Retinopathy-What-You-Should-Know-508.pdf>.
- Vascular Endothelial Growth Factor (VEGF) Trap-Eye: Investigation of Efficacy and Safety in Wet Age-Related Macular Degeneration (AMD) (VIEW1), NCT00509795, ClinicalTrials.gov (Apr. 28, 2009), <https://clinicaltrials.gov/ct2/show/NCT00509795> ("NCT-795").
- XOLAIR® label.
- Anonymous, Meeting Archive Titled "PA003 Eighteen-Month Results From an Extension Study of a Phase 2, Dose- and Interval-Ranging Study of VEGF Trap-Eye in Wet AMD," presented by David S Boyer, MD at Moscone Center (Oct. 2009).
- Anonymous, Meeting Archive Titled "PA040 One-Year Results of the DA VINCI Study of VEGF Trap-Eye in Diabetic Macular Edema," presented by Diana V Do, MD at Orange County Convention Center (Oct. 2011).
- Anonymous, Meeting Archive Titled "PA080 One-Year Results of a Phase 2 Study of Intravitreal VEGF Trap-Eye in Patients with Neovascular Age-Related Macular Degeneration," presented by David S Boyer, MD at Georgia World Congress Center (Nov. 2008).
- Anonymous, Meeting Archive Titled "PO259 OCT and Fluorescein Angiography Outcomes Through 1 Year for a Phase 2 Study of Intravitreal VEGF Trap-Eye in Neovascular AMD," presented by Peter K Kaiser, MD at Moscone Center (Oct. 2009).
- Anonymous, Meeting Archive Titled "PO260 VEGF Trap-Eye Vision-Specific Quality of Life Through 52 Weeks in Patients with Neovascular AMD in CLEAR-IT 2: A Phase 2 Clinical Trial," presented by Allen C Ho, MD at Moscone Center (Oct. 2009).
- Anonymous, Meeting Archive Titled "PO492 One-Year Results of the VIEW 1 and VIEW 2 Studies: VEGF Trap-Eye in Wet AMD," presented by David M Brown MD at Orange County Center (Oct. 2011).
- Anonymous, Meeting Archive Titled "PO549 The 6-Month (Primary Endpoint) Results of the Phase 3 GALILEO Study: VEGF Trap-Eye in Central Retinal Vein Occlusion," presented by Jean-Francois Korobelnik, MD at Orange County Convention Center (Oct. 2011).
- Anonymous, Meeting Archive Titled "PO571 OCT and Fluorescein Angiographic Outcomes Through 1 Year for the Phase 2 Study of Intravitreal VEGF Trap-Eye in Neovascular AMD," presented by Quan Dong Nguyen, MD at Georgia World Congress Center (Nov. 2008).
- Bontempo, "Preformulation Development of Parenteral Biopharmaceuticals," *Drugs and the Pharmaceutical Sciences*, 85:91-108 (1997).
- Bressler, N. M. Treatment of Age-Related Macular Degeneration with Photodynamic Therapy Study Group, "Photodynamic therapy of subfoveal choroidal neovascularization in age-related macular degeneration with verteporfin: two-year results of 2 randomized clinical trials—tap report 2," *Arch. Ophthalmol.*, 119(2):198-207 (2001).
- Brown et al., "Ranibizumab for Diabetic Macular Edema (DME): 24-Month Efficacy and Safety Results of RISE—a Phase 3 Randomized Controlled Trial," *ARVO Annual Meeting Abstract, Investigative Ophthalmology & Visual Science*, 52:6647 (Apr. 2011).
- Brown et al., "Sustained benefits from ranibizumab for macular edema following branch retinal vein occlusion: 12-month outcomes of a phase III study," *Ophthalmology*, 118(8):1594-2049 (2011).
- Cao et al., "VEGF Trap Promotes Regression of Choroidal Neovascularization (CNV) and Inhibits Fibrosis and Inflammation in the Subretinal Matrigel CNV Model," *ARVO Annual Meeting Abstract, Investigative Ophthalmology & Visual Science*, 50:2979 (Apr. 2009).
- Center for Drug Evaluation and Research Application No. 21-756 Medical Review(s) (Dec. 17, 2004) <URL:[https://www.accessdata.fda.gov/drugsatfda\\_docs/nda/2004/21-756\\_Macugen\\_medr.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/nda/2004/21-756_Macugen_medr.pdf)>.
- Center for Drug Evaluation and Research BLA Application No. 125156 Medical Review, (Jun. 2006) <URL:[https://www.accessdata.fda.gov/drugsatfda\\_docs/nda/2006/125156s0000\\_Lucentis\\_MedR.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/nda/2006/125156s0000_Lucentis_MedR.pdf)>.
- Cheung et al., "Combined anti-PIGF and anti-VEGF Therapy Ameliorates Pathological Neovascularization and Improves Retinal Revascularization in the Murine Model of Oxygen Induced Ischemic Retinopathy," *ARVO Annual Meeting Abstract, Investigative Ophthalmology & Visual Science*, 52:6064 (Apr. 2011).
- Dixon et al., "VEGF Trap-Eye for the treatment of neovascular age-related macular degeneration," *Expert Opin. Investig. Drugs*, 18(10):1573-1580 (2009).
- The Eyetech Study Group, "Anti-Vascular Endothelial Growth Factor Therapy for Subfoveal Choroidal Neovascularization Secondary to Age-related Macular Degeneration," *Ophthalmology*, 110(5):979-986 (May 2003).
- Hei er et al., "Ranibizumab for Choroidal Neovascularization Secondary to Causes Other Than Age-Related Macular Degeneration: A Phase I Clinical Trial," *Ophthalmology*, 118(1):111-118 (Jan. 2011).
- Hei er, "Intravitreal Aflibercept for Diabetic Macular Edema: 148-Week Results from the VISTA and VIVID Studies," *Ophthalmology*, 123(11):2376-2385 (2016).
- Hereceptin label, Sep. 1998.
- Information from ClinicalTrials.gov archive on the VIEW 2 study (NCT00637377) "VEGF Trap-Eye: Investigation of Efficacy and Safety in Wet AMD (VIEW 2)," v1 (Mar. 17, 2008).
- Ip et al., "A randomized trial comparing the efficacy and safety of intravitreal triamcinolone with observation to treat vision loss associated with macular edema secondary to central retinal vein occlusion: the Standard Care vs Corticosteroid for Retinal Vein Occlusion (SCORE) study report 5," *Arch. Ophthalmol.*, 127(9):1101-1114 (2009).
- Kaiser, "Vascular endothelial growth factor Trap-Eye for diabetic macular oedema," *Br. J. Ophthalmol.*, 93(2):135-36 (Feb. 2009).
- Korobelnik et al., "Intravitreal Aflibercept Injection for Macular Edema Resulting from Central Retinal Vein Occlusion," *Ophthalmology*, 121(1):202-208 (2014).
- Krzystolik et al., "Prevention of Experimental Choroidal Neovascularization With Intravitreal Anti-Vascular Endothelial Growth Factor Antibody Fragment," *Arch. Ophthalmol.*, 120(3):338-346 (Mar. 2002).
- Lalwani, "All About PrONTO: Study Yielded Good Results in AMD With Treatment Guided by OCT," *Retina Today* (May 2007).

(56)

## References Cited

## OTHER PUBLICATIONS


- Lobov et al., "VEGF Trap Treatment Regresses Pathological Neovessels, Improves Revascularization and Reduces Retinal Ischemia in the Murine Oxygen-Induced Retinopathy (OIR) Model," ARVO Annual Meeting Abstract, *Investigative Ophthalmology & Visual Science*, 52:3128 (Apr. 2011).
- Lucentis Approval (Jun. 30, 2006).
- Lucentis Label Title, 7 pages, Jun. 2010 [Cited in Third Party Observations filed in parent application U.S. Appl. No. 16/055,847].
- Macular Photocoagulation Study Group, "Laser photocoagulation of subfoveal neovascular lesions in age-related macular degeneration. Results of a randomized clinical trial," *Arc/z. Ophthalmol.*, 109(9):1220-1231 (1991).
- Mitchell et al., "Ranibizumab (Lucentis) in Neovascular Age-Related Macular Degeneration: Evidence from Clinical Trials," *Brit. J. Ophthalmology*, 94:2-13 (2010) (first online publication on May 20, 2009).
- Mitra et al., "Review of anti-vascular endothelial growth factor therapy in macular edema secondary to central retinal vein occlusions," *Expert Review in Ophthalmol.*, Taylor & Francis, GB 6(6):623-629 (Jan. 2011).
- Mousa and Mousa, "Current Status of Vascular Endothelial Growth Factor Inhibition in Age-Related Macular Degeneration," *Biodrugs*, 24(3):183-194 (2010).
- Nguyen et al., "A phase I trial of an IV-administered vascular endothelial growth factor trap for treatment in patients with choroidal neovascularization due to age-related macular degeneration," *Ophthalmology*, 113(9):1522e1-1522e14 (Sep. 2006) (epub Jul. 28, 2006).
- Regeneron Pharmaceuticals Inc., "Regeneron Receives \$20 Million Milestone Payment for Initiation of Phase 3 Study of VEGF Trap-Eye in Wet AMD," Media Release: Aug. 13, 2007. Available from URL: <http://www.regeneron.com>.
- Regeneron Pharmaceuticals Inc., "An Exploratory Study of the Safety, Tolerability and Biological Effect of a Single Intravitreal Administration of VEGF Trap in Patients with Diabetic Macular Edema," poster presented at the 2007 Association for Research in Vision and Ophthalmology meeting in Ft. Lauderdale, Florida (May 2007).
- Regeneron Pharmaceuticals Inc., "CLEAR-IT-2: Interim Results Of The Phase II, Randomized, Controlled Dose-and Interval-ranging Study Of Repeated Intravitreal VEGF Trap Administration In Patients With Neovascular Age-related Macular Degeneration (AMD)," poster presented at the 2007 Association for Research in Vision and Ophthalmology meeting in Ft. Lauderdale, Florida (May 2007).
- Regeneron Pharmaceuticals Inc., "Optical Coherence Tomography Outcomes of a Phase I, Dose-Escalation, Safety, Tolerability, and Bioactivity Study of Intravitreal VEGF Trap in Patients with Neovascular Age-Related Macular Degeneration: The CLEAR-IT 1 Study," poster presented at the 2007 Association for Research in Vision and Ophthalmology meeting in Ft. Lauderdale, Florida (May 2007).
- Regeneron Pharmaceuticals Inc., "Regeneron Reports First Quarter 2008 Financial and Operating Results," Press release May 1, 2008.
- Regeneron Pharmaceuticals Inc., Form 10-Q, published Nov. 7, 2007, for the period ending Sep. 30, 2007.
- Regillo et al., "Randomized, Double-Masked, Sham-Controlled Trial of Ranibizumab for Neovascular Age-related Macular Degeneration: OIER Study Year 1," *American Journal of Ophthalmology*, 145(2):239-248 (2008).
- Rosenfeld et al., "Optical coherence tomography findings after an intravitreal injection of bevacizumab (avastin) for neovascular age-related macular degeneration," *Ophthalmic Surg. Lasers Imaging*, 36(4):331-335 (2005).
- Scott et al., "A randomized trial comparing the efficacy and safety of intravitreal triamcinolone with standard care to treat vision loss associated with macular Edema secondary to branch retinal vein occlusion: the Standard Care vs Corticosteroid for Retinal Vein Occlusion (SCORE) study report 6," *Arch. Ophthalmol.*, 127(9):1115-1128 & 127(12):1653 (2009).
- Simo and Hernandez, "Advances in Medical Treatment of Diabetic Retinopathy," *Diabetes Care*, 32(8):1556-1562 (Aug. 2009).
- Slides for the 2008 Retina Society Meeting "VEGF Trap-Eye in Wet AMD CLEAR-IT 2: Summary of One-Year Key Results," Sep. 28, 2008.
- Tolentino et al., "One-year Results Of The Da Vinci Study of VEGF Trap-Eye In DME," ARVO Annual Meeting Abstract, *Investigative Ophthalmology & Visual Science*, 52:6646 (Apr. 2011).
- Van Bruggen et al., "VEGF antagonism reduces edema formation and tissue damage after ischemia/reperfusion injury in the mouse brain," *The Journal of clinical investigation*, 104(11):1613-1620 (1999).
- WHO Drug Information, "International Nonproprietary Names for Pharmaceutical Substances (INN)," 20(2):115-119 (2006).
- IPR2021-00880 dated Nov. 10, 2021, for U.S. Pat. No. 9,669,069 B2.
- IPR2021-00881 dated Nov. 10, 2021, for U.S. Pat. No. 9,254,338 B2.
- U.S. Pat. No. 7,374,758—Patent Term Extension Application dated Dec. 22, 2011.
- ADIS R&D Profile, "Aflibercept: AVE 0005, AVE 005, AVE0005, VEGF Trap—Regeneron, VEGF Trap (R1R2), VEGF Trap-Eye," *Drugs R. D.*, 9(4):261-269 (2008).
- Andersen & Krummen, "Recombinant protein expression for therapeutic applications," *Current Opinion in Biotechnology*, 13:117-123 (2002).
- Anderson et al., "Delivery of Anti-Angiogenic Molecular Therapies for Retinal Disease," *Drug Discovery Today*, 15(7/8):272-282 (2010).
- Article in Retinal Physician, "Subspecialty News," available online at <http://www.retinalphysician.com/printarticle.aspx?articleID=104007> (Mar. 2010).
- Ass'n for Res. Vision & Ophthalmology, ARVO News (Summer 2007).
- Ass'n for Res. Vision & Ophthalmology, ARVO News (Winter/Spring 2008).
- Avastin Label (Revised Dec. 2017).
- Avery, R. L., D. J. Pieramici, M. D. Rabena, A. A. Castellarin, M. A. Nasir and M. J. Giust, "Intravitreal bevacizumab (Avastin) for neovascular age-related macular degeneration," *Ophthalmology*, 113(3):363-372e5 (2006).
- Bashshur et al., "Intravitreal Bevacizumab for the Management of Choroidal Neovascularization in Age-Related Macular Degeneration," *Am. J. Ophthalmology*, 142(1):1-9 (2006).
- Bayer Press Release, "Bayer and Regeneron Dose First Patient in Second Phase 3 Study for VEGF Trap-Eye in Wet Age-Related Macular Degeneration," May 8, 2008.
- Bayer Press Release, "Bayer HealthCare and Regeneron Announce Encouraging 32-Week Follow Up Results From A Phase 2 Study of VEGF Trap-Eye in Age-Related Macular Degeneration," Apr. 28, 2008.
- Bayer Press Release, "Bayer HealthCare and Regeneron Announce VEGF Trap-Eye Achieved Durable Improvement in Vision Over 52 Weeks in a Phase 2 Study in Patients with Age-Related Macular Degeneration," Aug. 19, 2008.
- Bayer Press Release, "VEGF Trap-Eye Shows Positive Results in Phase II Study in Patients with Diabetic Macular Edema," Feb. 18, 2010.
- BMJ Publishing Group Ltd., "Review: Ranibizumab (Lucentis) In Neovascular Age-Related Macular Degeneration: Evidence From Clinical Trials," *British J. Ophthalmology*, (Dec. 2020), <https://bjoo.bmj.com/content/94/1/2.altmetrics>.
- Brown & Regillo, "Anti-VEGF Agents in the Treatment of Neovascular Age-Related Macular Degeneration: Applying Clinical Trial Results to the Treatment of Everyday Patients," *Am J. Ophthalmology*, 144(4):627-637e2 (2007).
- Chi et al., "Physical Stability of Proteins in Aqueous Solution: Mechanism and Driving Forces in Nonnative Protein Aggregation" *Pharmaceutical Research*, 20(9):1325-1336 (Sep. 2003).
- Ciulla & Rosenfeld, "Antivascular Endothelial Growth Factor Therapy For Neovascular Age-Related Macular Degeneration," *Current Opinion Ophthalmology*, 20:158-165 (2009).

(56)

## References Cited

## OTHER PUBLICATIONS

- Clinicaltrials.gov. I-SPY 2 TRIAL: Neoadjuvant and Personalized Adaptive Novel Agents to Treat Breast Cancer, Accessed 2010; <http://clinicaltrials.gov/ct2/show/NCT01042379?term=NCT01042379&rank=1>.
- CMS, Local Coverage Determination (LCD) for Ranibizumab (Lucentis) (L29266, First Coast Service Options, Inc Jun. 14, 2011). Controls in SCI experiments, RegenBase. Retrieved Jan. 6, 2021, from <http://regenbase.org/control-groups.html>.
- Drug Vehicle (Code C927), National Cancer Institute (NCI). Retrieved Jan. 6, 2021, from [https://ncithesaurus.nci.nih.gov/ncitbrowser/ConceptReport.jsp?dictionary=NCI\\_Thesaurus&code=C927&ns=ncit](https://ncithesaurus.nci.nih.gov/ncitbrowser/ConceptReport.jsp?dictionary=NCI_Thesaurus&code=C927&ns=ncit).
- EP 2 663 325 File History.
- Eylea Prescribing Information, Revised May 2019.
- Ferrara, N. & Kerbel, R., "Angiogenesis as a Therapeutic Target," *Nature*, 438:967-974 (2005).
- Fraser et al., "Single Injections of Vascular Endothelial Growth Factor Trap Block Ovulation in the Macaque and Produce a Prolonged, Dose-Related Suppression of Ovarian Function." *J. Clin. Endocrinol & Metab.* 90(2): 1114-1122 (Feb. 2005).
- Genentech, "FDA Approves Lucentis for the Treatment of Wet Age-Related Macular Degeneration," News Release dated Jun. 30, 2006 (Jun. 30, 2006).
- Gupta, O. P., G. Shienbaum, A. H. Patel, C. Fecarotta, R. S. Kaiser and C. D. Regillo, "A treat and extend regimen using ranibizumab for neovascular age-related macular degeneration clinical and economic impact," *Ophthalmology*, 117(11): 2134-2140 (2010).
- Heier et al., "Ranibizumab for macular edema due to retinal vein occlusions: long-term follow-up in the HORIZON trial," *Ophthalmology*, 119(4):802-809 (2012).
- Heier, "Intravitreal VEGF Trap for AMD: An Update," *Retina Today* 44 (Oct. 2009).
- Holz et al., "VEGF Trap-Eye for Macular Oedema Secondary to Central Retinal Vein Occlusion: 6-Month Results of the Phase III GALILEO Study," *British J. Ophthalmology*, 97:278-284 (2013).
- Information from ClinicalTrials.gov archive on the VIEW 1 study (NCT00509795) "Vascular Endothelial Growth Factor (VEGF) Trap-Eye: Investigation of Efficacy and Safety in Wet Age-Related Macular Degeneration (AMD) (VIEW1)," changes from v8 (Mar. 3, 2009) to v9 (Apr. 28, 2009).
- Janeeway et al., "The structure of a typical antibody molecule" *Immunobiology: The Immune System in Health and Disease*. 5th edition. New York: Garland Science (2001).
- Keane et al., "Effect of Ranibizumab Retreatment Frequency on Neurosensory Retinal Volume in Neovascular AMD," *Retina*, 29(5):592-600 (2009).
- Kim et al., "Potent VEGF Blockade Causes Regression of Coopted Vessels in a Model of Neuroblastoma," *Proc. Nat'l Acad. Sci.*, 99(17):11399-11404 (2002).
- Lucentis Label (Revised 2006).
- Lucentis Label (Revised 2014).
- Massin, "Anti-VEGF Therapy for Diabetic Macular Edema: An Update," *Retina Today* 54 (Sep./Oct. 2008).
- Michels, S., P. J. Rosenfeld, C. A. Puliafito, E. N. Marcus and A. S. Venkatraman, "Systemic bevacizumab (Avastin) therapy for neovascular age-related macular degeneration twelve-week results of an uncontrolled open-label clinical study," *Ophthalmology*, 112(6):1035-1047 (2005).
- Ni & Hui, "Emerging Pharmacologic Therapies for Wet Age-Related Macular Degeneration," *Ophthalmologica*, 223:401-410 (Published Online First May 20, 2009).
- Parkins & Lashmar, "The formulation of biopharmaceutical products," *Pharmaceutical Science & Technology Today*, 3(4):129-137 (Apr. 4, 2000).
- Phosphate buffer. Cold Spring Harbor Protocols 2006: pdb.rec8543 (2006).
- Randolph & Jones, "Surfactant-Protein Interactions" *Rational Design of Stable Protein Formulations* pp. 159-175, Springer, Boston, MA (2002).
- Raptiva Label (Final Labelling Mar. 13, 2009).
- Regeneron Pharmaceuticals Inc., Regeneron Reports Fourth Quarter and Full Year 2004 Financial and Operating Results. Media Release: Feb. 22, 2005.
- Regeneron Pharmaceuticals Inc., Regeneron Reports Fourth Quarter and Full Year 2005 Financial and Operating Results. Media Release: Feb. 24, 2006.
- Regeneron Pharmaceuticals Inc., Regeneron Reports Positive Phase 1 Data for the VEGF Trap in Age-Related Macular Degeneration; Preliminary Results Show Improvements in Vision and Reginal Swelling; VEGF Trap Was Well Tolerated at All Dose Levels. Media Release: May 1, 2006.
- Regeneron SEC Form 10-Q (Sep. 30, 2009).
- Reichert, "Antibody-Based Therapeutics To Watch In 2011," *MABS*, 3(1):76-99 (2011).
- Remicade Label (Revised Nov. 2013).
- Retina Coding Q & A, Retinal Physician, 16: 18, 54 (Jul./Aug. 2019).
- Rogers et al., "The prevalence of retinal vein occlusion: pooled data from population studies from the United States, Europe, Asia, and Australia," *Ophthalmology*, 117(2):313-319e1 (2010).
- Rudge et al., "VEGF Trap as a Novel Antiangiogenic Treatment Currently in Clinical Trials for Cancer and Eye Diseases, and VelociGene-based Discovery of the Next Generation of Angiogenesis Targets," *Cold Spring Harbor Symposia on Quantitative Biology*, 70:411-418 (2005).
- Schmidt-Erfurth, "Current Concepts in the Management of Diabetic Macular Edema," *Johns Hopkins Advanced Studies in Ophthalmology*, 7(2):52-59 (2010).
- Simulect Label (May 1998).
- Spaide et al., "Prospective Study of Intravitreal Ranibizumab as a Treatment for Decreased Visual Acuity Secondary to Central Retinal Vein Occlusion," *Am. J. Ophthalmology*, 147(2):298-306 (2009).
- Spielberg, L. & Leys, A., "Intravitreal Bevacizumab for Myopic Choroidal Neovascularization: Short-Term and 1-Year Results," *Bulletin Societe Belge D'Ophthalmologie*, 312:17-27 (2009).
- Steinbrook, "The Price of Sight—Ranibizumab, Bevacizumab, and the Treatment of Macular Degeneration," *N. Eng. J. Med.*, 355(14):1409-1412 (2006).
- The Branch Vein Occlusion Study, G., "Argon laser photocoagulation for macular edema in branch vein occlusion," *Am. J. Ophthalmology*, 98(3):271-282 (1984).
- The Central Vein Occlusion Study, G., "Evaluation of grid pattern photocoagulation for macular edema in central vein occlusion. The Central Vein Occlusion Study Group M report," *Ophthalmology*, 102(10):1425-1433 (1995).
- U.S. Department of Health and Human Services, Food and Drug Administration, "Guidance for industry Q1A(R2) stability testing of new drug substances and products," Rockville, MD (Nov. 2003).
- U.S. Department of Health and Human Services, National Institute of Health, National Eye Institute, "Age-Related Macular Degeneration: What You Should Know," (Sep. 2015) [https://www.nei.nih.gov/sites/default/files/healthpdfs/WYSK\\_AMD\\_English\\_Sept2015\\_PRINT.pdf](https://www.nei.nih.gov/sites/default/files/healthpdfs/WYSK_AMD_English_Sept2015_PRINT.pdf).
- U.S. Department of Health and Human Services, National Institute of Health, National Eye Institute, "Diabetic Retinopathy: What You Should Know," (Sep. 2015) <https://www.nei.nih.gov/sites/default/files/2019-06/Diabetic-Retinopathy-What-You-Should-Know-508.pdf>.
- U.S. Department of Health and Human Services, Office of Inspector General, "Questionable Billing for Medicare Ophthalmology Services" Sep. 2015 OEI-04-12-00280.
- Wall Street Journal, "Genentech's Big Drug for Eyes Faces a Rival" (2007).
- Wulff et al., "Prevention of Thecal Angiogenesis, Antral Follicular Growth, and Ovulation in the Primate by Treatment with Vascular Endothelial Growth Factor Trap R1R2" *Endocrinology* 143(7): 2797-2807 (Jul. 2002).
- Xolair Label (2003).
- Zarbin & Rosenfeld, "Pathway-Based Therapies for Age-Related Macular Degeneration: An Integrated Survey of Emerging Treatment Alternatives" *Retina* 30: 1350 (2010).



1

## USE OF A VEGF ANTAGONIST TO TREAT ANGIOGENIC EYE DISORDERS

### CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a continuation of Ser. No. 17/350,958 filed Jun. 17, 2021 which is a continuation of Ser. No. 17/112,404 filed Dec. 4, 2020 which is a continuation of Ser. No. 17/072,417 filed Oct. 16, 2020 which is a continuation of Ser. No. 16/055,847 filed Aug. 6, 2018, now U.S. Pat. No. 10,857,205 issued Dec. 8, 2020 and is a continuation of Ser. No. 16/397,267 filed Apr. 29, 2019, which is a continuation of Ser. No. 16/159,282 filed Oct. 12, 2018, now U.S. Pat. No. 10,828,345 issued Nov. 10, 2020, which is a continuation of Ser. No. 15/471,506 filed Mar. 28, 2017, now U.S. Pat. No. 10,130,681 issued Nov. 20, 2018, which is a continuation of Ser. No. 14/972,560 filed Dec. 17, 2015, now U.S. Pat. No. 9,669,069 issued Jun. 6, 2017, which is a continuation of Ser. No. 13/940,370 filed Jul. 12, 2013, now U.S. Pat. No. 9,254,338 issued Feb. 9, 2016, which is a continuation-in-part of International Patent Application No. PCT/US2012/020855, filed on Jan. 11, 2012, which claims the benefit of U.S. Provisional Application No. 61/432,245, filed on Jan. 13, 2011, 61/434,836, filed on Jan. 21, 2011, and 61/561,957, filed on Nov. 21, 2011, the contents of which are hereby incorporated by reference in their entireties.

### FIELD OF THE INVENTION

The present invention relates to the field of therapeutic treatments of eye disorders. More specifically, the invention relates to the administration of VEGF antagonists to treat eye disorders caused by or associated with angiogenesis.

### BACKGROUND

Several eye disorders are associated with pathological angiogenesis. For example, the development of age-related macular degeneration (AMD) is associated with a process called choroidal neovascularization (CNV). Leakage from the CNV causes macular edema and collection of fluid beneath the macula resulting in vision loss. Diabetic macular edema (DME) is another eye disorder with an angiogenic component. DME is the most prevalent cause of moderate vision loss in patients with diabetes and is a common complication of diabetic retinopathy, a disease affecting the blood vessels of the retina. Clinically significant DME occurs when fluid leaks into the center of the macula, the light-sensitive part of the retina responsible for sharp, direct vision. Fluid in the macula can cause severe vision loss or blindness. Yet another eye disorder associated with abnormal angiogenesis is central retinal vein occlusion (CRVO). CRVO is caused by obstruction of the central retinal vein that leads to a back-up of blood and fluid in the retina. The retina can also become ischemic, resulting in the growth of new, inappropriate blood vessels that can cause further vision loss and more serious complications. Release of vascular endothelial growth factor (VEGF) contributes to increased vascular permeability in the eye and inappropriate new vessel growth. Thus, inhibiting the angiogenic-promoting properties of VEGF appears to be an effective strategy for treating angiogenic eye disorders.

FDA-approved treatments of angiogenic eye disorders such as AMD and CRVO include the administration of an

2

anti-VEGF antibody called ranibizumab (Lucentis®, Genentech, Inc.) on a monthly basis by intravitreal injection.

Methods for treating eye disorders using VEGF antagonists are mentioned in, e.g., U.S. Pat. Nos. 7,303,746; 7,306,799; 7,300,563; 7,303,748; and US 2007/0190058. Nonetheless, there remains a need in the art for new administration regimens for angiogenic eye disorders, especially those which allow for less frequent dosing while maintaining a high level of efficacy.

### BRIEF SUMMARY OF THE INVENTION

The present invention provides methods for treating angiogenic eye disorders. The methods of the invention comprise sequentially administering multiple doses of a VEGF antagonist to a patient over time. In particular, the methods of the invention comprise sequentially administering to the patient a single initial dose of a VEGF antagonist, followed by one or more secondary doses of the VEGF antagonist, followed by one or more tertiary doses of the VEGF antagonists. The present inventors have surprisingly discovered that beneficial therapeutic effects can be achieved in patients suffering from angiogenic eye disorders by administering a VEGF antagonist to a patient at a frequency of once every 8 or more weeks, especially when such doses are preceded by about three doses administered to the patient at a frequency of about 2 to 4 weeks. Thus, according to the methods of the present invention, each secondary dose of VEGF antagonist is administered 2 to 4 weeks after the immediately preceding dose, and each tertiary dose is administered at least 8 weeks after the immediately preceding dose. An example of a dosing regimen of the present invention is shown in FIG. 1. One advantage of such a dosing regimen is that, for most of the course of treatment (i.e., the tertiary doses), it allows for less frequent dosing (e.g., once every 8 weeks) compared to prior administration regimens for angiogenic eye disorders which require monthly administrations throughout the entire course of treatment. (See, e.g., prescribing information for Lucentis® [ranibizumab], Genentech, Inc.).

The methods of the present invention can be used to treat any angiogenic eye disorder, including, e.g., age related macular degeneration, diabetic retinopathy, diabetic macular edema, central retinal vein occlusion, corneal neovascularization, etc.

The methods of the present invention comprise administering any VEGF antagonist to the patient. In one embodiment, the VEGF antagonist comprises one or more VEGF receptor-based chimeric molecule(s), (also referred to herein as a “VEGF-Trap” or “VEGFT”). An exemplary VEGF antagonist that can be used in the context of the present invention is a multimeric VEGF-binding protein comprising two or more VEGF receptor-based chimeric molecules referred to herein as “VEGFR1R2-FcΔC1(a)” or “aflibercept.”

Various administration routes are contemplated for use in the methods of the present invention, including, e.g., topical administration or intraocular administration (e.g., intravitreal administration).

Aflibercept (EYLEA™, Regeneron Pharmaceuticals, Inc) was approved by the FDA in November 2011, for the treatment of patients with neovascular (wet) age-related macular degeneration, with a recommended dose of 2 mg administered by intravitreal injection every 4 weeks for the first three months, followed by 2 mg administered by intravitreal injection once every 8 weeks.

Other embodiments of the present invention will become apparent from a review of the ensuing detailed description.

#### BRIEF DESCRIPTION OF THE FIGURE

FIG. 1 shows an exemplary dosing regimen of the present invention. In this regimen, a single “initial dose” of VEGF antagonist (“VEGFT”) is administered at the beginning of the treatment regimen (i.e. at “week 0”), two “secondary doses” are administered at weeks 4 and 8, respectively, and at least six “tertiary doses” are administered once every 8 weeks thereafter, i.e., at weeks 16, 24, 32, 40, 48, 56, etc.).

#### DETAILED DESCRIPTION

Before the present invention is described, it is to be understood that this invention is not limited to particular methods and experimental conditions described, as such methods and conditions may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting, since the scope of the present invention will be limited only by the appended claims.

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. As used herein, the term “about,” when used in reference to a particular recited numerical value, means that the value may vary from the recited value by no more than 1%. For example, as used herein, the expression “about 100” includes 99 and 101 and all values in between (e.g., 99.1, 99.2, 99.3, 99.4, etc.).

Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials are now described.

#### Dosing Regimens

The present invention provides methods for treating angiogenic eye disorders. The methods of the invention comprise sequentially administering to a patient multiple doses of a VEGF antagonist. As used herein, “sequentially administering” means that each dose of VEGF antagonist is administered to the patient at a different point in time, e.g., on different days separated by a predetermined interval (e.g., hours, days, weeks or months). The present invention includes methods which comprise sequentially administering to the patient a single initial dose of a VEGF antagonist, followed by one or more secondary doses of the VEGF antagonist, followed by one or more tertiary doses of the VEGF antagonist.

The terms “initial dose,” “secondary doses,” and “tertiary doses,” refer to the temporal sequence of administration of the VEGF antagonist. Thus, the “initial dose” is the dose which is administered at the beginning of the treatment regimen (also referred to as the “baseline dose”); the “secondary doses” are the doses which are administered after the initial dose; and the “tertiary doses” are the doses which are administered after the secondary doses. The initial, secondary, and tertiary doses may all contain the same amount of VEGF antagonist, but will generally differ from one another in terms of frequency of administration. In certain embodiments, however, the amount of VEGF antagonist contained in the initial, secondary and/or tertiary doses will vary from one another (e.g., adjusted up or down as appropriate) during the course of treatment.

In one exemplary embodiment of the present invention, each secondary dose is administered 2 to 4 (e.g., 2, 2½, 3,

3½, or 4) weeks after the immediately preceding dose, and each tertiary dose is administered at least 8 (e.g., 8, 8½, 9, 9½, 10, 10½, 11, 11½, 12, 12½, 13, 13½, 14, 14½, or more) weeks after the immediately preceding dose. The phrase “the immediately preceding dose,” as used herein, means, in a sequence of multiple administrations, the dose of VEGF antagonist which is administered to a patient prior to the administration of the very next dose in the sequence with no intervening doses.

In one exemplary embodiment of the present invention, a single initial dose of a VEGF antagonist is administered to a patient on the first day of the treatment regimen (i.e., at week 0), followed by two secondary doses, each administered four weeks after the immediately preceding dose (i.e., at week 4 and at week 8), followed by at least 5 tertiary doses, each administered eight weeks after the immediately preceding dose (i.e., at weeks 16, 24, 32, 40 and 48). The tertiary doses may continue (at intervals of 8 or more weeks) indefinitely during the course of the treatment regimen. This exemplary administration regimen is depicted graphically in FIG. 1.

The methods of the invention may comprise administering to a patient any number of secondary and/or tertiary doses of a VEGF antagonist. For example, in certain embodiments, only a single secondary dose is administered to the patient. In other embodiments, two or more (e.g., 2, 3, 4, 5, 6, 7, 8, or more) secondary doses are administered to the patient. Likewise, in certain embodiments, only a single tertiary dose is administered to the patient. In other embodiments, two or more (e.g., 2, 3, 4, 5, 6, 7, 8, or more) tertiary doses are administered to the patient.

In embodiments involving multiple secondary doses, each secondary dose may be administered at the same frequency as the other secondary doses. For example, each secondary dose may be administered to the patient 4 weeks after the immediately preceding dose. Similarly, in embodiments involving multiple tertiary doses, each tertiary dose may be administered at the same frequency as the other tertiary doses. For example, each tertiary dose may be administered to the patient 8 weeks after the immediately preceding dose. Alternatively, the frequency at which the secondary and/or tertiary doses are administered to a patient can vary over the course of the treatment regimen. For example, the present invention includes methods which comprise administering to the patient a single initial dose of a VEGF antagonist, followed by one or more secondary doses of the VEGF antagonist, followed by at least 5 tertiary doses of the VEGF antagonist, wherein the first four tertiary doses are administered 8 weeks after the immediately preceding dose, and wherein each subsequent tertiary dose is administered from 8 to 12 (e.g., 8, 8½, 9, 9½, 10, 10½, 11, 11½, 12) weeks after the immediately preceding dose. The frequency of administration may also be adjusted during the course of treatment by a physician depending on the needs of the individual patient following clinical examination.

#### VEGF Antagonists

The methods of the present invention comprise administering to a patient a VEGF antagonist according to specified dosing regimens. As used herein, the expression “VEGF antagonist” means any molecule that blocks, reduces or interferes with the normal biological activity of VEGF.

VEGF antagonists include molecules which interfere with the interaction between VEGF and a natural VEGF receptor, e.g., molecules which bind to VEGF or a VEGF receptor and prevent or otherwise hinder the interaction between VEGF and a VEGF receptor. Specific exemplary VEGF antagonists include anti-VEGF antibodies, anti-VEGF receptor antibod-

ies, and VEGF receptor-based chimeric molecules (also referred to herein as “VEGF-Traps”).

VEGF receptor-based chimeric molecules include chimeric polypeptides which comprise two or more immunoglobulin (Ig)-like domains of a VEGF receptor such as VEGFR1 (also referred to as Flt1) and/or VEGFR2 (also referred to as Flk1 or KDR), and may also contain a multimerizing domain (e.g., an Fc domain which facilitates the multimerization [e.g., dimerization] of two or more chimeric polypeptides). An exemplary VEGF receptor-based chimeric molecule is a molecule referred to as VEGFR1R2-FcΔC1(a) which is encoded by the nucleic acid sequence of SEQ ID NO:1. VEGFR1R2-FcΔC1(a) comprises three components: (1) a VEGFR1 component comprising amino acids 27 to 129 of SEQ ID NO:2; (2) a VEGFR2 component comprising amino acids 130 to 231 of SEQ ID NO:2; and (3) a multimerization component (“FcΔC1(a)”) comprising amino acids 232 to 457 of SEQ ID NO:2 (the C-terminal amino acid of SEQ ID NO:2 [i.e., K458] may or may not be included in the VEGF antagonist used in the methods of the invention; see e.g., U.S. Pat. No. 7,396,664). Amino acids 1-26 of SEQ ID NO:2 are the signal sequence.

The VEGF antagonist used in the Examples set forth herein below is a dimeric molecule comprising two VEGFR1R2-FcΔC1(a) molecules and is referred to herein as “VEGFT.” Additional VEGF receptor-based chimeric molecules which can be used in the context of the present invention are disclosed in U.S. Pat. Nos. 7,396,664, 7,303,746 and WO 00/75319.

#### Angiogenic Eye Disorders

The methods of the present invention can be used to treat any angiogenic eye disorder. The expression “angiogenic eye disorder,” as used herein, means any disease of the eye which is caused by or associated with the growth or proliferation of blood vessels or by blood vessel leakage. Non-limiting examples of angiogenic eye disorders that are treatable using the methods of the present invention include age-related macular degeneration (e.g., wet AMD, exudative AMD, etc.), retinal vein occlusion (RVO), central retinal vein occlusion (CRVO; e.g., macular edema following CRVO), branch retinal vein occlusion (BRVO), diabetic macular edema (DME), choroidal neovascularization (CNV; e.g., myopic CNV), iris neovascularization, neovascular glaucoma, post-surgical fibrosis in glaucoma, proliferative vitreoretinopathy (PVR), optic disc neovascularization, corneal neovascularization, retinal neovascularization, vitreal neovascularization, pannus, pterygium, vascular retinopathy, and diabetic retinopathies.

#### Pharmaceutical Formulations

The present invention includes methods in which the VEGF antagonist that is administered to the patient is contained within a pharmaceutical formulation. The pharmaceutical formulation may comprise the VEGF antagonist along with at least one inactive ingredient such as, e.g., a pharmaceutically acceptable carrier. Other agents may be incorporated into the pharmaceutical composition to provide improved transfer, delivery, tolerance, and the like. The term “pharmaceutically acceptable” means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly, in humans. The term “carrier” refers to a diluent, adjuvant, excipient, or vehicle with which the antibody is administered. A multitude of appropriate formulations can be found in the formulary known to all pharmaceutical chemists: Remington’s Pharmaceutical Sciences (15th ed, Mack Publishing Company, Easton, Pa., 1975), particularly Chapter

87 by Blaug, Seymour, therein. These formulations include, for example, powders, pastes, ointments, jellies, waxes, oils, lipids, lipid (cationic or anionic) containing vesicles (such as LIPOFECTIN™), DNA conjugates, anhydrous absorption pastes, oil-in-water and water-in-oil emulsions, emulsions carbowax (polyethylene glycols of various molecular weights), semi-solid gels, and semi-solid mixtures containing carbowax. Any of the foregoing mixtures may be appropriate in the context of the methods of the present invention, provided that the VEGF antagonist is not inactivated by the formulation and the formulation is physiologically compatible and tolerable with the route of administration. See also Powell et al. PDA (1998) J Pharm Sci Technol. 52:238-311 and the citations therein for additional information related to excipients and carriers well known to pharmaceutical chemists.

Pharmaceutical formulations useful for administration by injection in the context of the present invention may be prepared by dissolving, suspending or emulsifying a VEGF antagonist in a sterile aqueous medium or an oily medium conventionally used for injections. As the aqueous medium for injections, there are, for example, physiological saline, an isotonic solution containing glucose and other auxiliary agents, etc., which may be used in combination with an appropriate solubilizing agent such as an alcohol (e.g., ethanol), a polyalcohol (e.g., propylene glycol, polyethylene glycol), a nonionic surfactant [e.g., polysorbate 80, HCO-50 (polyoxyethylene (50 mol) adduct of hydrogenated castor oil)], etc. As the oily medium, there may be employed, e.g., sesame oil, soybean oil, etc., which may be used in combination with a solubilizing agent such as benzyl benzoate, benzyl alcohol, etc. The injection thus prepared can be filled in an appropriate ampoule if desired.

#### Modes of Administration

The VEGF antagonist (or pharmaceutical formulation comprising the VEGF antagonist) may be administered to the patient by any known delivery system and/or administration method. In certain embodiments, the VEGF antagonist is administered to the patient by ocular, intracocular, intravitreal or subconjunctival injection. In other embodiments, the VEGF antagonist can be administered to the patient by topical administration, e.g., via eye drops or other liquid, gel, ointment or fluid which contains the VEGF antagonist and can be applied directly to the eye. Other possible routes of administration include, e.g., intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, and oral.

#### Amount of VEGF Antagonist Administered

Each dose of VEGF antagonist administered to the patient over the course of the treatment regimen may contain the same, or substantially the same, amount of VEGF antagonist. Alternatively, the quantity of VEGF antagonist contained within the individual doses may vary over the course of the treatment regimen. For example, in certain embodiments, a first quantity of VEGF antagonist is administered in the initial dose, a second quantity of VEGF antagonist is administered in the secondary doses, and a third quantity of VEGF antagonist is administered in the tertiary doses. The present invention contemplates dosing schemes in which the quantity of VEGF antagonist contained within the individual doses increases over time (e.g., each subsequent dose contains more VEGF antagonist than the last), decreases over time (e.g., each subsequent dose contains less VEGF antagonist than the last), initially increases then decreases, initially decreases then increases, or remains the same throughout the course of the administration regimen.

The amount of VEGF antagonist administered to the patient in each dose is, in most cases, a therapeutically effective amount. As used herein, the phrase “therapeutically effective amount” means a dose of VEGF antagonist that results in a detectable improvement in one or more symptoms or indicia of an angiogenic eye disorder, or a dose of VEGF antagonist that inhibits, prevents, lessens, or delays the progression of an angiogenic eye disorder. In the case of an anti-VEGF antibody or a VEGF receptor-based chimeric molecule such as VEGFR1R2-FcΔC1(a), a therapeutically effective amount can be from about 0.05 mg to about 5 mg, e.g., about 0.05 mg, about 0.1 mg, about 0.15 mg, about 0.2 mg, about 0.25 mg, about 0.3 mg, about 0.35 mg, about 0.4 mg, about 0.45 mg, about 0.5 mg, about 0.55 mg, about 0.6 mg, about 0.65 mg, about 0.7 mg, about 0.75 mg, about 0.8 mg, about 0.85 mg, about 0.9 mg, about 1.0 mg, about 1.05 mg, about 1.1 mg, about 1.15 mg, about 1.2 mg, about 1.25 mg, about 1.3 mg, about 1.35 mg, about 1.4 mg, about 1.45 mg, about 1.5 mg, about 1.55 mg, about 1.6 mg, about 1.65 mg, about 1.7 mg, about 1.75 mg, about 1.8 mg, about 1.85 mg, about 1.9 mg, about 2.0 mg, about 2.05 mg, about 2.1 mg, about 2.15 mg, about 2.2 mg, about 2.25 mg, about 2.3 mg, about 2.35 mg, about 2.4 mg, about 2.45 mg, about 2.5 mg, about 2.55 mg, about 2.6 mg, about 2.65 mg, about 2.7 mg, about 2.75 mg, about 2.8 mg, about 2.85 mg, about 2.9 mg, about 3.0 mg, about 3.5 mg, about 4.0 mg, about 4.5 mg, or about 5.0 mg of the antibody or receptor-based chimeric molecule.

The amount of VEGF antagonist contained within the individual doses may be expressed in terms of milligrams of antibody per kilogram of patient body weight (i.e., mg/kg). For example, the VEGF antagonist may be administered to a patient at a dose of about 0.0001 to about 10 mg/kg of patient body weight.

#### Treatment Population and Efficacy

The methods of the present invention are useful for treating angiogenic eye disorders in patients that have been diagnosed with or are at risk of being afflicted with an angiogenic eye disorder. Generally, the methods of the present invention demonstrate efficacy within 104 weeks of the initiation of the treatment regimen (with the initial dose administered at “week 0”), e.g., by the end of week 16, by the end of week 24, by the end of week 32, by the end of week 40, by the end of week 48, by the end of week 56, etc. In the context of methods for treating angiogenic eye disorders such as AMD, CRVO, and DME, “efficacy” means that, from the initiation of treatment, the patient exhibits a loss of 15 or fewer letters on the Early Treatment Diabetic Retinopathy Study (ETDRS) visual acuity chart. In certain embodiments, “efficacy” means a gain of one or more (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or more) letters on the ETDRS chart from the time of initiation of treatment.

#### EXAMPLES

The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how to make and use the methods and compositions of the invention, and are not intended to limit the scope of what the inventors regard as their invention. Efforts have been made to ensure accuracy with respect to numbers used (e.g., amounts, temperature, etc.) but some experimental errors and deviations should be accounted for. Unless indicated otherwise, parts are parts by weight, molecular weight is average molecular weight, temperature is in degrees Centigrade, and pressure is at or near atmospheric.

The exemplary VEGF antagonist used in all Examples set forth below is a dimeric molecule having two functional VEGF binding units. Each functional binding unit is comprised of Ig domain 2 from VEGFR1 fused to Ig domain 3 from VEGFR2, which in turn is fused to the hinge region of a human IgG1 Fc domain (VEGFR1R2-FcΔC1(a); encoded by SEQ ID NO:1). This VEGF antagonist is referred to in the examples below as “VEGFT”. For purposes of the following Examples, “monthly” dosing is equivalent to dosing once every four weeks.

#### Example 1: Phase I Clinical Trial of Intravitreally Administered VEGF Receptor-Based Chimeric Molecule (VEGFT) in Subjects with Neovascular AMD

In this Phase I study, 21 subjects with neovascular AMD received a single intravitreal (IVT) dose of VEGFT. Five groups of three subjects each received either 0.05, 0.15, 0.5, 2 or 4 mg of VEGFT, and a sixth group of six subjects received 1 mg. No serious adverse events related to the study drug, and no identifiable intraocular inflammation was reported. Preliminary results showed that, following injection of VEGFT, a rapid decrease in foveal thickness and macular volume was observed that was maintained through 6 weeks. At Day 43 across all dose groups, mean excess retinal thickness [excess retinal thickness=(retinal thickness–179p)] on optical coherence tomography (OCT) was reduced from 119p to 27p as assessed by Fast Macular Scan and from 194p to 60p as assessed using a single Posterior Pole scan. The mean increase in best corrected visual acuity (BCVA) was 4.75 letters, and BCVA was stable or improved in 95% of subjects. In the 2 highest dose groups (2 and 4 mg), the mean increase in BCVA was 13.5 letters, with 3 of 6 subjects demonstrating improvement of  $\geq 3$  lines.

#### Example 2: Phase II Clinical Trial of Repeated Doses of Intravitreally Administered VEGF Receptor-Based Chimeric Molecule (VEGFT) in Subjects with Neovascular AMD

This study was a double-masked, randomized study of 3 doses (0.5, 2, and 4 mg) of VEGFT tested at 4-week and/or 12-week dosing intervals. There were 5 treatment arms in this study, as follows: 1) 0.5 mg every 4 weeks, 2) 0.5 mg every 12 weeks, 3) 2 mg every 4 weeks, 4) 2 mg every 12 weeks and 5) 4 mg every 12 weeks. Subjects were dosed at a fixed interval for the first 12 weeks, after which they were evaluated every 4 weeks for 9 months, during which additional doses were administered based on pre-specified criteria. All subjects were then followed for one year after their last dose of VEGFT. Preliminary data from a pre-planned interim analysis indicated that VEGFT met its primary endpoint of a statistically significant reduction in retinal thickness after 12 weeks compared with baseline (all groups combined, decrease of 135 $\mu$ ,  $p<0.0001$ ). Mean change from baseline in visual acuity, a key secondary endpoint of the study, also demonstrated statistically significant improvement (all groups combined, increase of 5.9 letters,  $p<0.0001$ ). Moreover, patients in the dose groups that received only a single dose, on average, demonstrated a decrease in excess retinal thickness ( $p<0.0001$ ) and an increase in visual acuity ( $p=0.012$ ) at 12 weeks. There were no drug-related serious adverse events, and treatment with the VEGF antagonists was generally well-tolerated. The

most common adverse events were those typically associated with intravitreal injections.

Example 3: Phase I Clinical Trial of Systemically Administered VEGF Receptor-Based Chimeric Molecule (VEGFT) in Subjects with Neovascular AMD

This study was a placebo-controlled, sequential-group, dose-escalating safety, tolerability and bioeffect study of VEGFT by IV infusion in subjects with neovascular AMD. Groups of 8 subjects meeting eligibility criteria for subfoveal choroidal neovascularization (CNV) related to AMD were assigned to receive 4 IV injections of VEGFT or placebo at dose levels of 0.3, 1, or 3 mg/kg over an 8-week period.

Most adverse events that were attributed to VEGFT were mild to moderate in severity, but 2 of 5 subjects treated with 3 mg/kg experienced dose-limiting toxicity (DLT) (one with Grade 4 hypertension and one with Grade 2 proteinuria); therefore, all subjects in the 3 mg/kg dose group did not enter the study. The mean percent changes in excess retinal thickness were: -12%, -10%, -66%, and -60% for the placebo, 0.3, 1, and 3 mg/kg dose groups at day 15 (ANOVA  $p < 0.02$ ), and -5.6%, +47.1%, and -63.3% for the placebo, 0.3, and 1 mg/kg dose groups at day 71 (ANOVA  $p < 0.02$ ). There was a numerical improvement in BCVA in the subjects treated with VEGFT. As would be expected in such a small study, the results were not statistically significant.

Example 4: Phase III Clinical Trials of the Efficacy, Safety, and Tolerability of Repeated Doses of Intravitreal VEGFT in Subjects with Neovascular Age-Related Macular Degeneration

A. Objectives, Hypotheses and Endpoints

Two parallel Phase III clinical trials were carried out to investigate the use of VEGFT to treat patients with the neovascular form of age-related macular degeneration (Study 1 and Study 2). The primary objective of these studies was to assess the efficacy of IVT administered VEGFT compared to ranibizumab (Lucentis®, Genentech, Inc.), in a non-inferiority paradigm, in preventing moderate vision loss in subjects with all subtypes of neovascular AMD.

The secondary objectives were (a) to assess the safety and tolerability of repeated IVT administration of VEGFT in subjects with all sub-types of neovascular AMD for periods up to 2 years; and (b) to assess the effect of repeated IVT administration of VEGFT on Vision-Related Quality of Life (QOL) in subjects with all sub-types of neovascular AMD.

The primary hypothesis of these studies was that the proportion of subjects treated with VEGFT with stable or improved BCVA (<15 letters lost) is similar to the proportion treated with ranibizumab who have stable or improved BCVA, thereby demonstrating non-inferiority.

The primary endpoint for these studies was the prevention of vision loss of greater than or equal to 15 letters on the ETDRS chart, compared to baseline, at 52 weeks. Secondary endpoints were as follows: (a) change from baseline to Week 52 in letter score on the ETDRS chart; (b) gain from baseline to Week 52 of 15 letters or more on the ETDRS chart; (c) change from baseline to Week 52 in total NEI VFQ-25 score; and (d) change from baseline to Week 52 in CNV area.

B. Study Design

For each study, subjects were randomly assigned in a 1:1:1:1 ratio to 1 of 4 dosing regimens: (1) 2 mg VEGFT administered every 4 weeks (2Q4); (2) 0.5 mg VEGFT administered every 4 weeks (0.5Q4); (3) 2 mg VEGFT administered every 4 weeks to week 8 and then every 8 weeks (with sham injection at the interim 4-week visits

when study drug was not administered (2Q8); and (4) 0.5 mg ranibizumab administered every 4 weeks (RQ4). Subjects assigned to (2Q8) received the 2 mg injection every 4 weeks to week 8 and then a sham injection at interim 4-week visits (when study drug is not to be administered) during the first 52 weeks of the studies. (No sham injection were given at Week 52).

The study duration for each subject was scheduled to be 96 weeks plus the recruitment period. For the first 52 weeks (Year 1), subjects received an IVT or sham injection in the study eye every 4 weeks. (No sham injections were given at Week 52). During the second year of the study, subjects will be evaluated every 4 weeks and will receive IVT injection of study drug at intervals determined by specific dosing criteria, but at least every 12 weeks. (During the second year of the study, sham injections will not be given.) During this period, injections may be given as frequently as every 4 weeks, but no less frequently than every 12 weeks, according to the following criteria: (i) increase in central retinal thickness of  $\geq 100 \mu\text{m}$  compared to the lowest previous value as measured by optical coherence tomography (OCT); or (ii) a loss from the best previous letter score of at least 5 ETDRS letters in conjunction with recurrent fluid as indicated by OCT; or (iii) new or persistent fluid as indicated by OCT; or (iv) new onset classic neovascularization, or new or persistent leak on fluorescein angiography (FA); or (v) new macular hemorrhage; or (vi) 12 weeks have elapsed since the previous injection. According to the present protocol, subjects must receive an injection at least every 12 weeks.

Subjects were evaluated at 4 weeks intervals for safety and best corrected visual acuity (BCVA) using the 4 meter ETDRS protocol. Quality of Life (QOL) was evaluated using the NEI VFQ-25 questionnaire. OCT and FA examinations were conducted periodically.

Approximately 1200 subjects were enrolled, with a target enrollment of 300 subjects per treatment arm.

To be eligible for this study, subjects were required to have subfoveal choroidal neovascularization (CNV) secondary to AMD. "Subfoveal" CNV was defined as the presence of subfoveal neovascularization, documented by FA, or presence of a lesion that is juxtafoveal in location angiographically but affects the fovea. Subject eligibility was confirmed based on angiographic criteria prior to randomization.

Only one eye was designated as the study eye. For subjects who met eligibility criteria in both eyes, the eye with the worse VA was selected as the study eye. If both eyes had equal VA, the eye with the clearest lens and ocular media and least amount of subfoveal scar or geographic atrophy was selected. If there was no objective basis for selecting the study eye, factors such as ocular dominance, other ocular pathology and subject preference were considered in making the selection.

Inclusion criteria for both studies were as follows: (i) signed Informed consent; (ii) at least 50 years of age; (iii) active primary subfoveal CNV lesions secondary to AMD, including juxtafoveal lesions that affect the fovea as evidenced by FA in the study eye; (iv) CNV at least 50% of total lesion size; (v) early treatment diabetic retinopathy study (ETDRS) best-corrected visual acuity of: 20/40 to 20/320 (letter score of 73 to 25) in the study eye; (vi) willing, committed, and able to return for all clinic visits and complete all study-related procedures; and (vii) able to read, understand and willing to sign the informed consent form (or, if unable to read due to visual impairment, be read to verbatim by the person administering the informed consent or a family member).

Exclusion criteria for both studies were as follows: 1. Any prior ocular (in the study eye) or systemic treatment or surgery for neovascular AMD except dietary supplements or vitamins. 2. Any prior or concomitant therapy with another investigational agent to treat neovascular AMD in the study

eye, except dietary supplements or vitamins. 3. Prior treatment with anti-VEGF agents as follows: (a) Prior treatment with anti-VEGF therapy in the study eye was not allowed; (b) Prior treatment with anti-VEGF therapy in the fellow eye with an investigational agent (not FDA approved, e.g. bevacizumab) was allowed up to 3 months prior to first dose in the study, and such treatments were not allowed during the study. Prior treatment with an approved anti-VEGF therapy in the fellow eye was allowed; (c) Prior systemic anti-VEGF therapy, investigational or FDA/Health Canada approved, was only allowed up to 3 months prior to first dose, and was not allowed during the study. 4. Total lesion size >12 disc areas (30.5 mm<sup>2</sup>, including blood, scars and neovascularization) as assessed by FA in the study eye. 5. Subretinal hemorrhage that is either 50% or more of the total lesion area, or if the blood is under the fovea and is 1 or more disc areas in size in the study eye. (If the blood is under the fovea, then the fovea must be surrounded 270 degrees by visible CNV.) 6. Scar or fibrosis, making up >50% of total lesion in the study eye. 7. Scar, fibrosis, or atrophy involving the center of the fovea. 8. Presence of retinal pigment epithelial tears or rips involving the macula in the study eye. 9. History of any vitreous hemorrhage within 4 weeks prior to Visit 1 in the study eye. 10. Presence of other causes of CNV, including pathologic myopia (spherical equivalent of -8 diopters or more negative, or axial length of 25 mm or more), ocular histoplasmosis syndrome, angioid streaks, choroidal rupture, or multifocal choroiditis in the study eye. 11. History or clinical evidence of diabetic retinopathy, diabetic macular edema or any other vascular disease affecting the retina, other than AMD, in either eye. 12. Prior vitrectomy in the study eye. 13. History of retinal detachment or treatment or surgery for retinal detachment in the study eye. 14. Any history of macular hole of stage 2 and above in the study eye. 15. Any intraocular or periocular surgery within 3 months of Day 1 on the study eye, except lid surgery, which may not have taken place within 1 month of day 1, as long as it was unlikely to interfere with the injection. 16. Prior trabeculectomy or other filtration surgery in the study eye. 17. Uncontrolled glaucoma (defined as intraocular pressure greater than or equal to 25 mm Hg despite treatment with anti-glaucoma medication) in the study eye. 18. Active intraocular inflammation in either eye. 19. Active ocular or periocular infection in either eye. 20. Any ocular or periocular infection within the last 2 weeks prior to Screening in either eye. 21. Any history of uveitis in either eye. 22. Active scleritis or episcleritis in either eye. 23. Presence or history of scleromalacia in either eye. 24. Aphakia or pseudophakia with absence of posterior capsule (unless it occurred as a result of a yttrium aluminum garnet [YAG] posterior capsulotomy) in the study eye. 25. Previous therapeutic radiation in the region of the study eye. 26. History of corneal transplant or corneal dystrophy in the study eye. 27. Significant media opacities, including cataract, in the study eye which might interfere with visual acuity, assessment of safety, or fundus photography. 28. Any concurrent intraocular condition in the study eye (e.g. cataract) that, in the opinion of the investigator, could require either medical or surgical intervention during the 96 week study period. 29. Any concurrent ocular condition in the study eye which, in the opinion of the investigator, could either increase the risk to the subject beyond what is to be expected from standard procedures of intraocular injection, or which otherwise may interfere with the injection procedure or with evaluation of efficacy or safety. 30. History of other disease, metabolic dysfunction, physical examination finding, or clinical laboratory finding giving reasonable suspicion of a disease or condition that contraindicates the use of an investigational drug or that might affect interpretation of the results of the study or render the subject at high risk for treatment complications. 31. Participation as a subject in any clinical study within the 12 weeks prior to

Day 1. 32. Any systemic or ocular treatment with an investigational agent in the past 3 months prior to Day 1. 33. The use of long acting steroids, either systemically or intraocularly, in the 6 months prior to day 1. 34. Any history of allergy to povidone iodine. 35. Known serious allergy to the fluorescein sodium for injection in angiography. 36. Presence of any contraindications indicated in the FDA Approved label for ranibizumab (Lucentis®). 37. Females who were pregnant, breastfeeding, or of childbearing potential, unwilling to practice adequate contraception throughout the study. Adequate contraceptive measures include oral contraceptives (stable use for 2 or more cycles prior to screening); IUD; Depo-Provera®; Norplant® System implants; bilateral tubal ligation; vasectomy; condom or diaphragm plus either contraceptive sponge, foam or jelly.

Subjects were not allowed to receive any standard or investigational agents for treatment of their AMD in the study eye other than their assigned study treatment with VEGFT or ranibizumab as specified in the protocol until they completed the Completion/Early Termination visit assessments. This includes medications administered locally (e.g., IVT, topical, juxtascleral or periorbital routes), as well as those administered systemically with the intent of treating the study and/or fellow eye.

The study procedures are summarized as follows:

Best Corrected Visual Acuity: Visual function of the study eye and the fellow eye were assessed using the ETDRS protocol (The Early Treatment Diabetic Retinopathy Study Group) at 4 meters. Visual Acuity examiners were certified to ensure consistent measurement of BCVA. The VA examiners were required to remain masked to treatment assignment.

Optical Coherence Tomography: Retinal and lesion characteristics were evaluated using OCT on the study eye. At the Screen Visit (Visit 1) images were captured and transmitted for both eyes. All OCT images were captured using the Zeiss Stratus OCT™ with software Version 3 or greater. OCT images were sent to an independent reading center where images were read by masked readers at visits where OCTs were required. All OCTs were electronically archived at the site as part of the source documentation. A subset of OCT images were read. OCT technicians were required to be certified by the reading center to ensure consistency and quality in image acquisition. Adequate efforts were made to ensure that OCT technicians at the site remained masked to treatment assignment.

Fundus Photography and Fluorescein Angiography (FA): The anatomical state of the retinal vasculature of the study eye was evaluated by funduscopic examination, fundus photography and FA. At the Screen Visit (Visit 1) funduscopic examination, fundus photography and FA were captured and transmitted for both eyes. Fundus and angiographic images were sent to an independent reading center where images were read by masked readers. The reading center confirmed subject eligibility based on angiographic criteria prior to randomization. All FAs and fundus photographs were archived at the site as part of the source documentation. Photographers were required to be certified by the reading center to ensure consistency and quality in image acquisition. Adequate efforts were made to ensure that all photographers at the site remain masked to treatment assignment.

Vision-Related Quality of Life: Vision-related QOL was assessed using the National Eye Institute 25-Item Visual Function Questionnaire (NEI VFQ-25) in the interviewer-administered format. NEI VFQ-25 was administered by certified personnel at a contracted call center. At the screening visit, the sites assisted the subject and initiated the first call to the call center to collect all of the subject's contact information and to complete the first NEI VFQ-25 on the phone prior to randomization and IVT injection. For all subsequent visits, the call center called the subject on the phone, prior to IVT injection, to complete the questionnaire.

Intraocular Pressure: Intraocular pressure (IOP) of the study eye was measured using applanation tonometry or Tonopen. The same method of IOP measurement was used in each subject throughout the study.

#### C. Results Summary (52 Week Data)

The primary endpoint (prevention of moderate or severe vision loss as defined above) was met for all three VEGFT groups (2Q4, 0.5Q4 and 2Q8) in this study. The results from both studies are summarized in Table 1.

TABLE 1

	Ranibizumab 0.5 mg monthly (RQ4)	VEGFT 0.5 mg monthly (0.5Q4)	VEGFT 2 mg monthly (2Q4)	VEGFT 2 mg every 8 weeks <sup>[a]</sup> (2Q8)
Maintenance of vision* (% patients losing <15 letters) at week 52 versus baseline				
Study 1	94.4%	95.9%**	95.1%**	95.1%**
Study 2	94.4%	96.3%**	95.6%**	95.6%**
Mean improvement in vision* (letters) at 52 weeks versus baseline (p-value vs RQ4)***				
Study 1	8.1	6.9 (NS)	10.9 (p <0.01)	7.9 (NS)
Study 2	9.4	9.7 (NS)	7.6 (NS)	8.9 (NS)

<sup>[a]</sup>Following three initial monthly doses

\*Visual acuity was measured as the total number of letters read correctly on the Early Treatment Diabetic Retinopathy Study (ETDRS) eye chart.

\*\*Statistically non-inferior based on a non-inferiority margin of 10%, using confidence interval approach (95.1% and 95% for Study 1 and Study 2, respectively)

\*\*\*Test for superiority

NS = non-significant

In Study 1, patients receiving VEGFT 2 mg monthly (2Q4) achieved a statistically significant greater mean improvement in visual acuity at week 52 versus baseline (secondary endpoint), compared to ranibizumab 0.5 mg monthly (RQ4); patients receiving VEGFT 2 mg monthly on average gained 10.9 letters, compared to a mean 8.1 letter gain with ranibizumab 0.5 mg dosed every month (p<0.01). All other dose groups of VEGFT in Study 1 and all dose groups in Study 2 were not statistically different from ranibizumab in this secondary endpoint.

A generally favorable safety profile was observed for both VEGFT and ranibizumab. The incidence of ocular treatment emergent adverse events was balanced across all four treatment groups in both studies, with the most frequent events associated with the injection procedure, the underlying disease, and/or the aging process. The most frequent ocular adverse events were conjunctival hemorrhage, macular degeneration, eye pain, retinal hemorrhage, and vitreous floaters. The most frequent serious non-ocular adverse events were typical of those reported in this elderly population who receive intravitreal treatment for wet AMD; the most frequently reported events were falls, pneumonia,

myocardial infarction, atrial fibrillation, breast cancer, and acute coronary syndrome. There were no notable differences among the study arms.

#### Example 5: Phase II Clinical Trial of VEGFT in Subjects with Diabetic Macular Edema (DME)

In this study, 221 patients with clinically significant DME with central macular involvement were randomized, and 219 patients were treated with balanced distribution over five groups. The control group received macular laser therapy at baseline, and patients were eligible for repeat laser treatments, but no more frequently than at 16 week intervals. The remaining four groups received VEGFT by intravitreal injection as follows: Two groups received 0.5 or 2 mg of VEGFT once every four weeks throughout the 12-month dosing period (0.5Q4 and 2Q4, respectively). Two groups received three initial doses of 2 mg VEGFT once every four weeks (i.e., at baseline, and weeks 4 and 8), followed through week 52 by either once every 8 weeks dosing (2Q8) or as needed dosing with very strict repeat dosing criteria (PRN). Mean gains in visual acuity versus baseline were as shown in Table 2:

TABLE 2

	n	Mean change in visual acuity at week 24 versus baseline (letters)	Mean change in visual acuity at week 52 versus baseline (letters)
Laser	44	2.5	-1.3
VEGFT 0.5 mg monthly (0.5Q4)	44	8.6**	11.0**
VEGFT 2 mg monthly (2Q4)	44	11.4**	13.1**
VEGFT 2 mg every 8 weeks <sup>[a]</sup> (2Q8)	42	8.5**	9.7**
VEGFT 2 mg as needed <sup>[a]</sup> (PRN)	45	10.3**	12.0**

<sup>[a]</sup>Following three initial monthly doses

\*\*p < 0.01 versus laser

In this study, the visual acuity gains achieved with VEGFT administration at week 24 were maintained or numerically improved up to completion of the study at week 52 in all VEGFT study groups, including 2 mg dosed every other month

As demonstrated in the foregoing Examples, the administration of VEGFT to patients suffering from angiogenic eye disorders (e.g., AMD and DME) at a frequency of once every 8 weeks, following a single initial dose and two secondary doses administered four weeks apart, resulted in significant prevention of moderate or severe vision loss or improvements in visual acuity.

Example 6: A Randomized, Multicenter,  
Double-Masked Trial in Treatment Naïve Patients  
with Macular Edema Secondary to CRVO

In this randomized, double-masked, Phase 3 study, patients received 6 monthly injections of either 2 mg intravitreal VEGFT (114 patients) or sham injections (73 patients). From Week 24 to Week 52, all patients received 2 mg VEGFT as-needed (PRN) according to retreatment criteria. Thus, “sham-treated patients” means patients who received sham injections once every four weeks from Week 0 through Week 20, followed by intravitreal VEGFT as needed from Week 24 through Week 52. “VEGFT-treated patients” means patients who received VEGFT intravitreal injections once every four weeks from Week 0 through Week 20, followed by intravitreal VEGFT as needed from Week 24 through Week 52. The primary endpoint was the proportion of patients who gained  $\geq 15$  ETDRS letters from baseline at Week 24. Secondary visual, anatomic, and Quality of Life NEI VFQ-25 outcomes at Weeks 24 and 52 were also evaluated.

At Week 24, 56.1% of VEGFT-treated patients gained ETDRS letters from baseline vs 12.3% of sham-treated patients ( $P < 0.0001$ ). Similarly, at Week 52, 55.3% of VEGFT-treated patients gained  $\geq 15$  letters vs 30.1% of sham-treated patients ( $P < 0.01$ ). At Week 52, VEGFT-treated patients gained a mean of 16.2 letters vs 3.8 letters for sham-treated patients ( $P < 0.001$ ). Mean number of injections was 2.7 for VEGFT-treated patients vs 3.9 for sham-treated patients. Mean change in central retinal thickness was  $-413.0 \mu\text{m}$  for VEGFT-treated patients vs  $-381.8 \mu\text{m}$  for sham-treated patients. The proportion of patients with ocular neovascularization at Week 24 were 0% for VEGFT-treated patients and 6.8% for sham-treated patients, respectively; at Week 52 after receiving VEGFT PRN, proportions were 0% and 6.8% for VEGFT-treated and sham-treated. At Week 24, the mean change from baseline in the VFQ-25 total score was 7.2 vs 0.7 for the VEGFT-treated and sham-treated groups; at Week 52, the scores were 7.5 vs 5.1 for the VEGFT-treated and sham-treated groups.

This Example confirms that dosing monthly with 2 mg intravitreal VEGFT injection resulted in a statistically significant improvement in visual acuity at Week 24 that was maintained through Week 52 with PRN dosing compared with sham PRN treatment. VEGFT was generally well tolerated and had a generally favorable safety profile.

Example 7: Dosing Regimens

Specific, non-limiting examples of dosing regimens within the scope of the present invention are as follows:

VEGFT 2 mg (0.05 mL) administered by intravitreal injection once every 4 weeks (monthly).

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 8 weeks, followed by 2 mg (0.05 mL) via intravitreal injection once every 8 weeks.

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 8 weeks, followed by 2 mg (0.05 mL) via intravitreal injection on a less frequent basis based on visual and/or anatomical outcomes (as assessed by a physician or other qualified medical professional).

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 8 weeks, followed by 2 mg (0.05 mL) via intravitreal injection administered pro re nata (PRN) based on visual and/or anatomical outcomes (as assessed by a physician or other qualified medical professional).

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 12 weeks, followed by 2 mg (0.05 mL) via intravitreal injection once every 8 weeks.

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 12 weeks, followed by 2 mg (0.05 mL) via intravitreal injection on a less frequent basis based on visual and/or anatomical outcomes (as assessed by a physician or other qualified medical professional).

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 12 weeks, followed by 2 mg (0.05 mL) via intravitreal injection administered pro re nata (PRN) based on visual and/or anatomical outcomes (as assessed by a physician or other qualified medical professional).

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 16 weeks, followed by 2 mg (0.05 mL) via intravitreal injection once every 8 weeks.

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 16 weeks, followed by 2 mg (0.05 mL) via intravitreal injection on a less frequent basis based on visual and/or anatomical outcomes (as assessed by a physician or other qualified medical professional).

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 16 weeks, followed by 2 mg (0.05 mL) via intravitreal injection administered pro re nata (PRN) based on visual and/or anatomical outcomes (as assessed by a physician or other qualified medical professional).

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 20 weeks, followed by 2 mg (0.05 mL) via intravitreal injection once every 8 weeks.

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 20 weeks, followed by 2 mg (0.05 mL) via intravitreal injection on a less frequent basis based on visual and/or anatomical outcomes (as assessed by a physician or other qualified medical professional).

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 20 weeks, followed by 2 mg (0.05 mL) via intravitreal injection administered pro re nata (PRN) based on visual and/or anatomical outcomes (as assessed by a physician or other qualified medical professional).

17

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 24 weeks, followed by 2 mg (0.05 mL) via intravitreal injection once every 8 weeks.

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 24 weeks, followed by 2 mg (0.05 mL) via intravitreal injection on a less frequent basis based on visual and/or anatomical outcomes (as assessed by a physician or other qualified medical professional).

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 24 weeks, followed by 2 mg (0.05 mL) via intravitreal injection administered pro re nata (PRN) based on visual and/or anatomical outcomes (as assessed by a physician or other qualified medical professional).

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 28 weeks, followed by 2 mg (0.05 mL) via intravitreal injection once every 8 weeks.

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 28 weeks, followed by 2 mg (0.05 mL) via intravitreal injection on a less frequent basis based on visual and/or anatomical outcomes (as assessed by a physician or other qualified medical professional).

VEGFT 2 mg (0.5 mL) administered by intravitreal injection once every 4 weeks for the first 28 weeks, followed by 2 mg (0.05 mL) via intravitreal injection administered pro re nata (PRN) based on visual and/or anatomical outcomes (as assessed by a physician or other qualified medical professional).

VEGFT 2 mg (0.05 mL) administered by intravitreal injection as a single initial dose, followed by additional doses administered pro re nata (PRN) based on visual and/or anatomical outcomes (as assessed by a physician or other qualified medical professional).

Variations on the above-described dosing regimens would be appreciated by persons of ordinary skill in the art and are also within the scope of the present invention. For example, the amount of VEGFT and/or volume of formulation administered to a patient may be varied based on patient characteristics, severity of disease, and other diagnostic assessments by a physician or other qualified medical professional.

Any of the foregoing administration regimens may be used for the treatment of, e.g., age-related macular degeneration (e.g., wet AMD, exudative AMD, etc.), retinal vein occlusion (RVO), central retinal vein occlusion (CRVO; e.g., macular edema following CRVO), branch retinal vein occlusion (BRVO), diabetic macular edema (DME), choroidal neovascularization (CNV; e.g., myopic CNV), iris neovascularization, neovascular glaucoma, post-surgical fibrosis in glaucoma, proliferative vitreoretinopathy (PVR),

18

optic disc neovascularization, corneal neovascularization, retinal neovascularization, vitreal neovascularization, panus, pterygium, vascular retinopathy, etc.

## SEQUENCES

SEQ ID NO: 1 (DNA sequence having 1377 nucleotides):

ATGGTCAGCTACTGGGACACCGGGCTCTGTGTGCGCGCTCAGCTG  
TCTGCTTCTCACAGGATCTAGTTCCGGAAGTGATACCGGTAGACCTTTCG  
TAGAGATGTACAGTGAATCCCCGAAATTATACACATGACTGAAGGAAGG  
GAGCTCGTCATTCCTGCGGGTTACGTCACTAACATCACTGTTACTTT  
AAAAAAGTTTCCACTTGACACTTTGATCCCTGATGGAAAAACGCATAATCT  
GGGACAGTAGAAAGGGCTTCATCATATCAAATGCAACGTACAAGAAAAATA  
GGGCTTCTGACCTGTGAAGCAACAGTCAATGGGCATTTGTATAAGACAAA  
CTATCTCACACATCGACAAACCAATACATCATAGATGTGGTCTGAGTCT  
CGTCTCATGGAATTGAATCTCTGTTGGAGAAAAGCTTGTCTTAATTTGT  
ACAGCAAGAACTGAACATAATGTGGGATTGACTTCAACTGGGAATACCC  
TTCTTCGAAGCATCAGCATAAGAACTTGTAAACCGAGACCTAAAAACCC  
AGTCTGGGAGTGAGATGAAGAAATTTTGTAGCACCTTAATATAGATGGT  
GTAACCCGGAGTGACCAAGGATTGTACACCTGTGCAGCATCCAGTGGGCT  
GATGACCAAGAAGAACAGCACATTGTTCAGGGTCCATGAAAAGGACAAAA  
CTCACACATGCCACCGTGGCCAGCACCTGAACTCCTGGGGGACCGTCA  
GTCTTCTCTTCCCCCAAAACCCAGGACACCTCATGATCTCCCGGAC  
CCCTGAGGTCACATGCGTGGTGGTGGACGTGAGCCACGAAGACCTGAGG  
TCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAATGCCAAGACA  
AAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTCAAGCTCCT  
CACCGTCTGCACCAAGGACTGGCTGAATGGCAAGGAGTACAAGTCAAGG  
TCTCCAACAAGCCCTCCAGCCCCATCGAGAAAACCATCTCCAAAGCC  
AAAGGGCAGCCCCGAGAACACAGGTGTACACCTGCCCCCATCCCGGGA  
TGAGCTGACCAAGAACAGGTCAGCCTGACCTGCCTGGTCAAAGGCTTCT  
ATCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAAC  
AACTACAAGACCACGCTCCCGTGTGGACTCCGACGGCTCCTCTTCTCT  
CTCAGCAGCTCACCCTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCT  
TTCATGCTCCGTGATGATGAGGCTCTGCACAACCACTACACGCAAGAG  
AGCCTCTCCCTGTCTCCGGTAAATGA

SEQ ID NO: 2 (polypeptide sequence having 458 amino acids):

MVSYWDTGVLLCALLSCLLLTGSSSGSDTGRPFVEMYSEIPEI IHMTEGR  
ELVIPCRVTSNPITVTLKKFPLDTLIPDGKRI IWSRKGFI ISNATYKEI  
GLLTCEATVNGHLYKNTYLTNRQTNTIIDVVLSPSHGIELSVGEKLVLC  
TARTELNVGIDFNWEYPSKHKHKLVRDLKTSQSGSEMKKFLSTLTIDG  
VTRSDQGLYTCAASSGLMTKKNSTFVRVHEKDKHTTCCPPAPELLGGPS  
VFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVVHNAKT  
KPREQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPI EKTISKAK  
KGQPREPVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPEN  
NYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVSCFVSVMHEALHNYTQK  
SLSLSPGK

The present invention is not to be limited in scope by the specific embodiments described herein. Indeed, various modifications of the invention in addition to those described herein will become apparent to those skilled in the art from the foregoing description and the accompanying FIGURES. Such modifications are intended to fall within the scope of the appended claims.

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 2

<210> SEQ ID NO 1

<211> LENGTH: 1377

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 1

atgggtcagct	actggggacac	cggggtcctg	ctgtgcgcgc	tgctcagctg	tctgcttctc	60
acaggatcta	gttcgggaag	tgataccggg	agacctttcg	tagagatgta	cagtgaaatc	120
cccgaatta	tacacatgac	tgaagggaag	gagctcgtca	ttccctgccg	ggttacgtca	180
cctaacaatca	ctgttacttt	aaaaaagtgt	ccacttgaca	ctttgatccc	tgatggaaaa	240
cgcataatct	gggacagtag	aaagggcttc	atcatatcaa	atgcaacgta	caaagaaata	300
gggcttctga	cctgtgaagc	aacagtcaat	gggcatttgt	ataagacaaa	ctatctcaca	360
catcgacaaa	ccaataacaat	catagatgtg	gttctgagtc	cgtctcatgg	aattgaacta	420
tctgttggag	aaaagcttgt	cttaaattgt	acagcaagaa	ctgaactaaa	tgtggggatt	480
gacttcaact	gggaataacc	ttcttcgaag	catcagcata	agaaacttgt	aaaccgagac	540
ctaaaaaacc	agtctgggag	tgagatgaag	aaatttttga	gcaccttaac	tatagatggg	600
gtaacccgga	gtgaccaagg	attgtacacc	tgtgcagcat	ccagtgggct	gatgaccaag	660
aagaacagca	catttgtcag	gggccatgaa	aaggacaaaa	ctcacacatg	cccaccgtgc	720
ccagcacctg	aactcctggg	gggacgcgtc	gtcttctctc	tcccccaaaa	acccaaggac	780
accctcatga	tctcccgga	cctgaggtc	acatgcgtgg	tggtggacgt	gagccacgaa	840
gacctgagg	tcaagttaa	ctgtacatg	gacggcggtg	aggtgcataa	tgccaagaca	900
aagccgctg	aggagcagta	caacagcacg	taccgtgtgg	tcagcgtcct	caccgtcctg	960
caccaggact	ggctgaatgg	caaggagtag	aagtgaagg	tctccaacaa	agccctccca	1020
gcccccatcg	agaaaaccat	ctccaaagcc	aaagggcagc	cccgagaacc	acaggtgtac	1080
accctgcccc	catcccgga	tgagctgacc	aagaaccagg	tcagcctgac	ctgectggtc	1140
aaaggcttct	atcccagcga	catcgccgtg	gagtgggaga	gcaatgggca	gccggagaa	1200
aactacaaga	ccacgcctcc	cgtgctggac	tccgacggct	ccttcttctc	ctacagcaag	1260
ctcaccgtgg	acaagagcag	tgggcagcag	gggaacgtct	tctcatgctc	cgtgatgcat	1320
gaggctctgc	acaaccacta	cacgcagaag	agcctctccc	tgtctccggg	taaatga	1377

```
<210> SEQ ID NO 2
<211> LENGTH: 458
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic
```

<400> SEQUENCE: 2

Met	Val	Ser	Tyr	Trp	Asp	Thr	Gly	Val	Leu	Leu	Cys	Ala	Leu	Leu	Ser	
1				5					10					15		
Cys	Leu	Leu	Leu	Thr	Gly	Ser	Ser	Ser	Gly	Ser	Asp	Thr	Gly	Arg	Pro	
			20					25					30			
Phe	Val	Glu	Met	Tyr	Ser	Glu	Ile	Pro	Glu	Ile	Ile	His	Met	Thr	Glu	
		35					40					45				
Gly	Arg	Glu	Leu	Val	Ile	Pro	Cys	Arg	Val	Thr	Ser	Pro	Asn	Ile	Thr	
	50					55					60					
Val	Thr	Leu	Lys	Lys	Phe	Pro	Leu	Asp	Thr	Leu	Ile	Pro	Asp	Gly	Lys	
65					70					75					80	
Arg	Ile	Ile	Trp	Asp	Ser	Arg	Lys	Gly	Phe	Ile	Ile	Ser	Asn	Ala	Thr	
			85						90					95		
Tyr	Lys	Glu	Ile	Gly	Leu	Leu	Thr	Cys	Glu	Ala	Thr	Val	Asn	Gly	His	
			100					105					110			
Leu	Tyr	Lys	Thr	Asn	Tyr	Leu	Thr	His	Arg	Gln	Thr	Asn	Thr	Ile	Ile	
		115					120					125				

-continued

---

Asp Val Val Leu Ser Pro Ser His Gly Ile Glu Leu Ser Val Gly Glu		
130	135	140
Lys Leu Val Leu Asn Cys Thr Ala Arg Thr Glu Leu Asn Val Gly Ile		
145	150	155 160
Asp Phe Asn Trp Glu Tyr Pro Ser Ser Lys His Gln His Lys Lys Leu		
	165	170 175
Val Asn Arg Asp Leu Lys Thr Gln Ser Gly Ser Glu Met Lys Lys Phe		
	180	185 190
Leu Ser Thr Leu Thr Ile Asp Gly Val Thr Arg Ser Asp Gln Gly Leu		
	195	200 205
Tyr Thr Cys Ala Ala Ser Ser Gly Leu Met Thr Lys Lys Asn Ser Thr		
	210	215 220
Phe Val Arg Val His Glu Lys Asp Lys Thr His Thr Cys Pro Pro Cys		
225	230	235 240
Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro		
	245	250 255
Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys		
	260	265 270
Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp		
	275	280 285
Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu		
	290	295 300
Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu		
305	310	315 320
His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn		
	325	330 335
Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly		
	340	345 350
Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu		
	355	360 365
Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr		
	370	375 380
Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn		
385	390	395 400
Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe		
	405	410 415
Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn		
	420	425 430
Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr		
	435	440 445
Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys		
450	455	

---

23

What is claimed is:

1. A method of treating an angiogenic eye disorder in a patient in need thereof comprising sequentially administering to the patient by intravitreal injection a single initial dose of 2 mg of aflibercept, followed by one or more secondary doses of 2 mg of aflibercept, followed by one or more tertiary doses of 2 mg of aflibercept;

wherein each secondary dose is administered approximately 4 weeks following the immediately preceding dose; and

wherein each tertiary dose is administered approximately 8 weeks following the immediately preceding dose;

wherein the patient achieves a gain in visual acuity within 52 weeks following the initial dose.

2. The method of claim 1 wherein the patient achieves a gain in Best Corrected Visual Acuity (BCVA) according to Early Treatment Diabetic Retinopathy Study (ETDRS) letter score.

3. The method of claim 2 wherein the patient gains at least 7 letters Best Corrected Visual Acuity (BCVA) according to Early Treatment Diabetic Retinopathy Study (ETDRS) letter score.

4. The method of claim 3 wherein the patient achieves the gain in visual acuity within 24 weeks following the initial dose.

5. The method of claim 3 wherein only two secondary doses are administered to the patient.

6. The method of claim 3 wherein the aflibercept is formulated as an isotonic solution.

7. The method of claim 3 wherein the aflibercept is formulated with a nonionic surfactant.

8. The method of claim 2 wherein the patient gains at least 8 letters Best Corrected Visual Acuity (BCVA) according to Early Treatment Diabetic Retinopathy Study (ETDRS) letter score.

9. The method of claim 8 wherein the patient achieves the gain in visual acuity within 24 weeks following the initial dose.

10. The method of claim 2 wherein the patient gains at least 9 letters Best Corrected Visual Acuity (BCVA) according to Early Treatment Diabetic Retinopathy Study (ETDRS) letter score.

11. The method of claim 10 wherein only two secondary doses are administered to the patient.

12. The method of claim 10 wherein the aflibercept is formulated as an isotonic solution.

13. The method of claim 10 wherein the aflibercept is formulated with a nonionic surfactant.

14. The method of claim 1 wherein exclusion criteria for the patient include both of:

- (1) active ocular inflammation; and
- (2) active ocular or periocular infection.

15. A method of treating diabetic macular edema in a patient in need thereof comprising sequentially administering to the patient a single initial dose of 2 mg of aflibercept, followed by one or more secondary doses of 2 mg of aflibercept, followed by one or more tertiary doses of 2 mg of aflibercept;

wherein each secondary dose is administered to the patient by intravitreal injection approximately 4 weeks following the immediately preceding dose; and

wherein each tertiary dose is administered to the patient by intravitreal injection approximately 8 weeks following the immediately preceding dose.

16. The method of claim 15 wherein the patient achieves a gain in visual acuity within 52 weeks following the initial dose.

24

17. The method of claim 16 wherein the patient gains at least 9 letters Best Corrected Visual Acuity (BCVA) according to Early Treatment Diabetic Retinopathy Study (ETDRS) letter score.

18. The method of claim 17 wherein the aflibercept is formulated as an isotonic solution.

19. The method of claim 17 wherein the aflibercept is formulated with a non-ionic surfactant.

20. The method of claim 17 wherein the patient achieves a gain in visual acuity within 24 weeks following the initial dose.

21. The method of claim 16 wherein the patient gains at least 8 letters Best Corrected Visual Acuity (BCVA) according to Early Treatment Diabetic Retinopathy Study (ETDRS) letter score.

22. The method of claim 21 wherein the aflibercept is formulated as an isotonic solution.

23. The method of claim 21 wherein the aflibercept is formulated with a nonionic surfactant.

24. The method of claim 15 wherein only two secondary doses are administered to the patient.

25. The method of claim 15 wherein four secondary doses are administered to the patient.

26. A method of treating age related macular degeneration in a patient in need thereof comprising sequentially administering to the patient a single initial dose of 2 mg of aflibercept, followed by one or more secondary doses of 2 mg of aflibercept, followed by one or more tertiary doses of 2 mg of aflibercept;

wherein each secondary dose is administered to the patient by intravitreal injection approximately 4 weeks following the immediately preceding dose; and

wherein each tertiary dose is administered to the patient by intravitreal injection approximately 8 weeks following the immediately preceding dose;

wherein the method is as effective in achieving a gain in visual acuity as monthly administration of 0.5 mg of ranibizumab by intravitreal injection in human subjects with age-related macular degeneration at 52 weeks following the initial dose.

27. The method of claim 26 wherein only two secondary doses are administered to the patient.

28. The method of claim 26 wherein the gain in visual acuity is measured using the Early Treatment Diabetic Retinopathy Study (ETDRS) letter score.

29. A method of treating age-related macular degeneration in a patient in need thereof comprising sequentially administering to the patient a single initial dose of 2 mg of aflibercept, followed by one or more secondary doses of 2 mg of aflibercept, followed by one or more tertiary doses of 2 mg of aflibercept;

wherein each secondary dose is administered to the patient by intravitreal injection approximately 4 weeks following the immediately preceding dose; and

wherein each tertiary dose is administered to the patient by intravitreal injection approximately 8 weeks following the immediately preceding dose;

wherein the method is as effective in maintaining visual acuity as monthly administration of 0.5 mg of ranibizumab by intravitreal injection in human subjects with age-related macular degeneration at 52 weeks following the initial dose.

**25**

**30.** The method of claim **29** wherein maintenance of visual acuity means loss of less than 15 letters Best Corrected Visual Acuity (BCVA) as measured by using the Early Treatment Diabetic Retinopathy Study (ETDRS) letter score.

5

\* \* \* \* \*

**26**